



## **Effect of Irrigation and N-fertigation Levels on Broccoli Performance in a Polynet House**

**Sathyamurthy Naik <sup>a\*</sup>, K. S. Kumar <sup>b#</sup>, Sai Kumar Rondla <sup>ct</sup> and K. Kishan <sup>a</sup>**

<sup>a</sup> Department of Soil and Water Engineering, College of Agricultural Engineering, Sangareddy, PJTSAU, Telangana, India.

<sup>b</sup> College of Agricultural Engineering, Sangareddy, PJTSAU, Telangana, India.

<sup>c</sup> Agricultural College, Jagtial, PJTSAU, Telangana, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2021/v11i1230576

#### Editor(s):

(1) Dr. Wen-Cheng Liu, National United University, Taiwan.

#### Reviewers:

(1) Edinson Fabian Monroy Avila, National Open and Distance University – UNAD, Colombia.

(2) Erzsebet Buta, University of Agricultural Sciences of Veterinary Medicine Cluj – Napoca, Romania.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/80192>

**Original Research Article**

**Received 08 October 2021**

**Accepted 16 December 2021**

**Published 17 December 2021**

### **ABSTRACT**

The purpose of this study is to see how the microclimate of the polynet house, irrigation, and fertigation levels affect broccoli output. The experiment was set up in a randomized block design within bounds beneath a polynet house with 320 m<sup>2</sup> floor area covered with 50% perforated green colour net. Plant height, dry matter production yield, and its attributes were found to be maximum in drip irrigation (1.0Epan) + 125 % N at all growth stages, as per effect of scheduled irrigation and fertigation levels on crop biometric performance. Drip irrigation (1.0Epan) + 125% N resulted in a high higher yield (22.02 t ha<sup>-1</sup>); while treatments drip irrigation (1.5 Epan) + 75% N resulted as in lowest yield (18.02 t ha<sup>-1</sup>). The treatment drip irrigation (1.0Epan) + 125% N had a maximum NPK uptake. Polynet house covered with 50% perforation of green colour is suitable for realizing favorable microclimate for broccoli cultivation. Drip irrigation scheduled at 1.0 E pan with 125%N in the poly net house is recommended for broccoli production in central Telangana agro-climatic conditions as it maintained high yield and water use efficiency.

<sup>#</sup>Associate Dean,

<sup>†</sup>Assistant Professor,

\*Corresponding author: E-mail: [sathyamurthy.nayak@gmail.com](mailto:sathyamurthy.nayak@gmail.com);

**Keywords:** Broccoli; Nitrogen fertigation; irrigation; poly net house; yield and nutrient uptake.

## 1. INTRODUCTION

Broccoli (*Brassica oleracea L var. italica*) is a recently introduced exotic vegetable to India. The earliest sprouting broccoli selection was most likely made in pre-Christian Greece [1]. Broccoli is a vegetable that is a cabbage-like edible green plant with a large blossoming head. Broccoli is one of the most nutritious cole crops since it is high in vitamins and minerals, and boiling broccoli lowers the quantities of anti-carcinogenic substances like sulforaphane. It contains the same amount of calcium as milk, making it a vital source of nourishment for persons suffering from osteoporosis or calcium deficiency [2].

Polyhouse technology is an effective crop production intervention, especially under difficult climatic circumstances. In comparison to open field farming, it has the ability to provide year-round production of high-quality produce from modest landholdings. By maximizing productivity per unit area, it can address the challenge of year-round climatic variations as well as price variations and production. The theory behind a polyhouse is that crop output is controlled not only by genetics but also by the environment surrounding the plant. Temperature, light, relative humidity, air composition, and the nature of the root medium can all be regulated to give the plant the greatest possible environment for development.

Optimizing the irrigation and fertigation levels for highly nitrous and medicinal crop like broccoli under poly net house will improve the production potential and thereby improving the living standards of individuals. Such studies are more pertinent in semiarid tropical climate of Telangana, as the polyhouse industry is growing fast because of existence of favorable climate, resulting in high economic returns. Keeping the concept in back drop, a field experiment entitled "Optimization of irrigation and N-fertigation levels for broccoli (*Brassica oleracea var. L italica*) under poly net house conditions" was conducted at College of Agricultural Engineering, Kandi, Sangareddy during *rabi*, 2019-20.

## 2. MATERIALS METHODS

The experimental polynet house is located at the College of Agricultural Engineering's Precision

Farming Section in Kandi, Sangareddy district, Telangana. It is situated at a latitude of 17°19' N and a longitude of 78°28'E, at a height of 535 meters above sea level. The climate in the area is semiarid tropical, with an average annual rainfall of 800 mm, with 66 percent of that falling between June and October.

## 2.1 Experimental Design and Details

The soil of the experimental site was sandy loam in texture, low in available nitrogen (243.2), high in available phosphorus (32.6 kg ha<sup>-1</sup>) and low in available potassium K (243.2) K<sub>2</sub>O kg ha<sup>-1</sup>). The experiment was conducted during *rabi* 2019-20 and laid out in a randomized block design with plot size of 320 m<sup>2</sup> area. A total of nine irrigation accompanied fertigation levels were set out under poly net house condition. All the treatments were replicated thrice for assessing the level of significance and keeping the error degree of freedom in limits. The data on plant growth parameters *viz.*, plant height, dry matter accumulation at 30, 60, 90 days after transplanting and at harvest were recorded. The data on yield and its attributes consisted of curd weight (gm) and broccoli yield (t ha<sup>-1</sup>) was also recorded. In addition, nutrient analysis on the available nitrogen, phosphorus and potassium uptake at harvest with post harvest soil nutrient status pertaining to N, P and K was also performed.

## 3. RESULTS AND DISCUSSION

### 3.1 Observation of Microclimate Parameters

On an average, daily variation of air temperature in the poly net house ranged from 15°C to 28°C degree centigrade throughout the season, which is favorable for vegetable crop growth. Relative humidity variations in the poly net house revealed that, it was always found more at 12:30 PM and 4:00 PM during the experimental season and is favorable for broccoli cultivation. Solar radiation levels vary significantly in the poly net house and ranged from 4100.0 to 13221.3lux which is favorable for poly net house vegetable crop growth.

### 3.2 Evaluation of Broccoli Crop Biometric Performance under Semiarid Tropical Conditions

#### 3.2.1 Plant height (cm)

The main crop biometric growth factor i.e., plant height response for different irrigation and fertigation levels is presented in Table 1. It has been observed that the crop height progressively increased with the age of the crop and up to harvest, irrespective of the treatments at 30, 60 and 90 Days after transplanting and at harvest. Plant height significantly varied with different dates of transplanting and an overview of the results indicated that significantly superior plant height (37.3) was observed in T6 (polynet house condition (PNH) +Drip fertigation of 1.0 Epan +

125% N) at all growth stages and at harvest followed by T5 (PNH + Drip fertigation of 1.0 Epan + 100% N) of 36.2 cm, which is comparable. Significantly lower plant height (32.8 cm) was recorded in T7 (PNH + Drip fertigation of 1.5 Epan+ 75% N, when compared with all the levels of irrigation and fertigation. The increased plant height in the treatment of PNH + Drip fertigation (1.0Epan) + 125% N treatment is due to maintenance of optimum soil moisture coupled with availability nutrients in the soil profile due to frequent irrigation favoring higher nutrient uptake, rapid cell division and cell enlargement. The decrease in plant height at harvest in all the treatments is due to senescence of plants. These results are in quite agreement with the findings of Locasio and Smajstrala [3] and Candido et al. [4].




Fig. 1. Experimental layout of broccoli through drip irrigation in polynet house

Table 1. Plant height (cm) as influenced by different Irrigation and N-Fertigation levels

Treatments	Plant height (cm)			
	30DAT	60 DAT	90 DAT	At harvest
T <sub>1</sub> - PNH + drip fertigation (0.50 Epan) + 75%N	20.8	29.6	35.8	34.0
T <sub>2</sub> - PNH+drip fertigation (0.50 Epan) + 100%N	21.9	30.2	36.5	35.0
T <sub>3</sub> - PNH + drip fertigation (0.50 Epan) + 125%N	22.1	31.0	36.9	35.2
T <sub>4</sub> - PNH + drip fertigation (1.0Epan) + 75%N	21.5	30.0	36.2	34.2
T <sub>5</sub> - PNH + drip fertigation (1.0 Epan) + 100%N	23.0	31.4	37.3	36.2
T <sub>6</sub> - PNH + drip fertigation (1.0 Epan) + 125%N	24.1	32.0	38.2	37.3
T <sub>7</sub> - PNH + drip fertigation (1.5Epan) + 75%N	19.1	27.9	34.8	32.8
T <sub>8</sub> - PNH + drip fertigation (1.5 Epan) + 100%N	19.8	28.1	35.1	33.0
T <sub>9</sub> - PNH + drip fertigation (1.5 Epan) + 125%N	20.0	28.9	35.5	33.5
SEm ±	0.14	0.13	0.14	0.12
CD @5%	0.42	0.86	0.39	0.35
CV (%)	1.17	0.76	0.64	0.60
	NS	S	S	S

### 3.2.2 Number of leaves

Table 2 shows the crop's biometric performance in terms of leaf number. Among the various irrigation and fertigation optimization treatments applied to broccoli, treatment T6, i.e., PNH + Drip fertigation at 1.0 Epan+ 125 percent N at harvest, resulted in a greater number of leaves (32 per plant). T5 of PNH + Drip fertigation at 1.0 Epan) + 100 percent N, on the other hand, produced a comparable number of leaves (29 per plant). The treatment T7, i.e., PNH + Drip fertigation at 1.5 Epan+ 75 percent N (17 per plant), has the fewest leaves per plant, which is significantly inferior. The increase in vegetative growth parameters in broccoli is due to increased nutrient availability and fertigation, which results in better nutrient uptake and utilization in plant organ growth and development [5]. Reported similar results in a broccoli crop that are in line with the current study's findings.

### 3.2.3 Yield and yield attributes

Data pertaining to broccoli yield ( $t\ ha^{-1}$ ) at all levels of irrigation and fertigation, as well as at all growth stages, had been significantly influenced, and the results are presented in Table 3. Among the various levels of irrigation and fertigation, the treatment of PNH + Drip fertigation (1.0 Epan) +125 percent N (22.02 t  $ha^{-1}$ ) produced the highest yield of 22.02kg/ha, which was found to be significantly superior to all other treatments. PNH + Drip fertigation (1.5 Epan) +75 percent N had the lowest yield (18.02 t  $ha^{-1}$ ). PNH + Drip fertigation (1.5 Epan) +75 percent N yields were lowered. The higher head yield obtained at 1.0 Epan with 125% RDN of water-soluble fertigation might be due to increased nutrient dynamics in the root zone and in plants under drip fertigation. Similar results were also reported by Erdem et al. [6] and Vasu and Reddy [7] which are in quite agreement.

**Table 2. Number of leaves of broccoli as influenced by different Irrigation and N-Fertigation levels**

Treatments	No. of leaves			
	30 DAT	60 DAT	90 DAT	At harvest
T <sub>1</sub> -PNH + drip fertigation (0.50 Epan) + 75%N	10.0	16.0	27.0	22.0
T <sub>2</sub> -PNH + drip fertigation (0.50 Epan) + 100%N	10.0	18.0	30.0	25.0
T <sub>3</sub> -PNH + drip fertigation (0.50 Epan) + 125%N	10.0	20.0	31.0	26.0
T <sub>4</sub> -PNH + drip fertigation (1.0 Epan) + 75%N	9.0	18.0	28.0	23.0
T <sub>5</sub> -PNH + drip fertigation (1.0 Epan) + 100%N	11.0	21.0	33.0	29.0
T <sub>6</sub> -PNH + drip fertigation (1.0 Epan) + 125%N	13.0	23.0	35.0	32.0
T <sub>7</sub> -PNH + drip fertigation (1.5Epan) + 75%N	9.0	12.0	21.0	17.0
T <sub>8</sub> -PNH + drip fertigation (1.5 Epan) + 100%N	8.0	14.0	23.0	19.0
T <sub>9</sub> -PNH + drip fertigation (1.5 Epan) + 125%N	9.0	15.0	25.0	20.0
SEm ±	0.12	0.16	0.15	0.05
CD @5%	0.37	0.47	0.44	0.16
CV (%)	2.26	1.61	0.94	0.42
	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>

**Table 3. Broccoli head weight, yield per plot and yield (t/ha) as influenced by irrigation and N-fertigation levels**

Treatments	Average head weight(g)	Head Yield per plot (Kg)	Yield (t/ha)
T <sub>1</sub> -PNH + drip fertigation (0.50 Epan) + 75%N	399.00	1.97	19.70
T <sub>2</sub> -PNH + drip fertigation (0.50 Epan) + 100%N	411.90	2.03	20.34
T <sub>3</sub> -PNH + drip fertigation (0.50 Epan) + 125%N	421.60	2.08	20.82
T <sub>4</sub> -PNH + drip fertigation (1.0 Epan) + 75%N	404.80	2.00	19.99
T <sub>5</sub> -PNH + drip fertigation (1.0 Epan) + 100%N	430.00	2.12	21.23
T <sub>6</sub> -PNH + drip fertigation (1.0 Epan) + 125%N	446.00	2.20	22.02
T <sub>7</sub> -PNH+ drip fertigation (1.5Epan) + 75%N	365.00	1.80	18.02
T <sub>8</sub> -PNH + drip fertigation (1.5 Epan) + 100%N	378.00	1.87	18.67
T <sub>9</sub> -PNH + drip fertigation (1.5 Epan) + 125%N	389.00	1.92	19.21
SEm ±	0.12	0.06	0.02
CD @5%	0.36	0.19	0.05
CV (%)	0.05	5.00	0.14
	<b>S</b>	<b>S</b>	<b>S</b>

**Table 4. N, P and K uptake (kg ha<sup>-1</sup>) of the broccoli leaves excluding head as influenced by irrigation and N-fertigation levels**

Treatments	% N in leaves	N uptake (Kg/ha)	% P in leaves	P uptake (Kg/ha)	% K in leaves	K uptake (Kg/ha)
T <sub>1</sub> -PNH + drip fertigation (0.50 Epan) + 75%N	1.88	66.94	0.45	16.14	2.65	94.48
T <sub>2</sub> -PNH+ drip fertigation (0.50 Epan) + 100%N	2.01	75.53	0.52	19.63	2.86	107.17
T <sub>3</sub> -PNH+ drip fertigation (0.50 Epan) + 125%N	2.13	82.77	0.59	22.96	2.90	112.99
T <sub>4</sub> -PNH + drip fertigation (1.0 Epan) + 75%N	1.88	67.64	0.47	17.06	2.79	102.13
T <sub>5</sub> -PNH + drip fertigation (1.0 Epan) + 100%N	2.32	99.22	0.65	27.90	2.96	126.40
T <sub>6</sub> -PNH + drip fertigation (1.0 Epan) + 125%N	2.63	118.04	0.67	30.11	3.12	140.36
T <sub>7</sub> -PNH+ drip fertigation (1.5Epan) + 75%N	1.85	58.84	0.44	14.07	2.45	77.89
T <sub>8</sub> -PNH + drip fertigation (1.5 Epan) + 100%N	1.95	62.41	0.50	16.13	2.63	84.21
T <sub>9</sub> -PNH + drip fertigation (1.5 Epan) + 125%N	1.96	66.56	0.55	18.67	3.05	105.24
SEm ±	0.01	0.45	0.09	0.40	0.09	3.31
CD @5%	0.03	1.31	0.02	1.18	0.27	9.00
CV (%)	1.14	1.05	3.13	3.49	5.00	5.00

### 3.3 Plant Nutrient Content and Uptake of Nutrients

Nutrient content of the plant parts needs to be optimum for normal functioning of the plant. The level of nutrient uptake is better indicated by the nutrient content of the plant parts particularly leaves and dry matter production. The results obtained during the process of experiment on the nutrient content of the wrapper leaves in broccoli at harvest are presented in Table 4. The amount of nitrogen, phosphorus, and potassium in the leaf, as well as absorption, were all affected by fertigation. The treatment of PNH + Drip fertigation (1.0 Epan) + 125 percent N had the greatest N, P, and K content in leaf and uptake of 2.63 percent, 118.04 Kg ha<sup>-1</sup>, 0.67 percent and 30.11 Kg ha<sup>-1</sup>, and 3.12 percent and 140.36 Kg ha<sup>-1</sup>, respectively, at harvest.

Nutrient uptake is a function of dry matter production, available nutrient status of the soil. The right combination of water and nutrients are essential for obtaining higher yields along with quality. The method of fertilizer application is also important in improving the use efficiency of the nutrients. Drip fertigation method enables the adequate supplies of water and nutrients with precise timing and uniform distribution to meet the crop nutrient demand [8-9]. Better uptake of N, P and K under PHC + Drip irrigation (1.0 Epan) + 125% N might be due to higher dry matter production. This could be attributed to the synergistic effects of higher water content and available nutrients in the soil and crop growth. These results are conformity with the findings of Gupta et al. [10].

### 4. CONCLUSION

Optimizing the irrigation and fertigation levels for highly nitrous and medicinal crop like broccoli under poly net house will improve the production potential and thereby improving the living standards of individuals. Growth parameters viz., plant height (at 30, 60 and 90 DAT) and dry matter accumulation (at 30, 60, 90 DAT and at harvest) were significantly higher with the SNHC + Drip irrigation (1.0 x Epan) + 125% N at all the growth stages. Significantly lower yields have been observed in 22.02 t/ha and Curd weight 446 grams and yield 22.02 t ha<sup>-1</sup> were significantly higher with PNH + drip fertigation (1.0 Epan) + 125%N. Drip irrigation scheduled at 1.0 E pan with 125% N in the poly net house is recommended for broccoli production in central

Telangana agro-climatic conditions i.e tropical to semiarid as it maintained high yield.

### ACKNOWLEDGEMENT

The authors are grateful to Professor Jayashankar Telangana State Agricultural University, Rajendranagar, and Hyderabad for their support.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCE

1. Heywood VH. Flowering plant of the world. Mayflower Books, New York. 1978;2-3.
2. Nirmal D, Singh KP, Benerjee MK, Rai M. Exotic Vegetables-Technical Bulletin Indian Institute of Vegetable Research, Varanasi, India. 2004;2(1):4-6.
3. Locasio SJ, Smajstrala AG. Water application scheduling by pan evaporation for dry irrigated tomato. Journal of American Society of Horticultural Science. 1996;121(1): 63-68.
4. Candido V, Miccolis V, Perniola M. Effects of irrigation regime on yield and quality of processing tomato (*Lycopersicon-esculentum* Mill.) cultivars.III International Symposium on Irrigation of Horticultural Crops.Acta Horticulture. 2000;537:779-788.
5. Biradar MS, Mantur SM, Dhotre M. Influence of fertigation on growth and yield of broccoli and red cabbage under net house conditions. Acta Horticulturae. 2018;(1227):485-490. DOI:10.17660/actahortic.2018.1227.61.
6. Erdem T, Arin L, Erdem Y, Polat S, Devenci M, Okursoy H and Gultas HT. 2010. Yield and quality response of drip irrigated broccoli (*Brassica oleracea* L. var. italica) under different irrigation regimes, nitrogen applications and cultivation periods. Agricultural Water Management. 2010;97: 681-688.
7. Vasu D, Reddy MS. Effect of fertigation on yield, quality, nutrient uptake, fertilizer and water use efficiency in cabbage (*Brassica oleracea*).Agropedology (Nagpur). 2013; 23(2):106-112.
8. Patel N, Rajput TBS. Effect of fertigation on growth and yield of onion. In: Micro

- Irrigation, CBIP publication no. 2000;282: 451-454.
9. Narda NK, Chawla JK. A simple nitrate sub-model for trickle fertigated potatoes. Irrigation and Drainage. 2002;51:361-371.
10. Gupta AJ, Ahmed N, Lal Singh. Response of lettuce to drip irrigation and fertigation. Indian Journal of Fertilizers. 2010;6(5):12-16.

---

© 2021 Naik et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80192>