



Anti – Androgenic Activity of *Caesalpinia bonducella* in Androgen-induced Polycystic Ovarian Syndrome Rats

M. Sumithra ^{a*}, V. Chitra ^a, Murali Krishna Moka ^a, Padaleeswaran ^a and Safeeq Ahamed ^a

^a *Department of Pharmacology, SRM College of Pharmacy, SRMIST Kattankulathur, Tamil Nadu, 603203, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MS designed the study. Author VC did the protocol framing. Authors MKM and Padaleeswaran managed the analysis of the study. Author SA did the statistical evaluation. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i55A33826

Editor(s):

(1) Dr. Mohamed Fawzy Ramadan Hassanien, Zagazig University, Egypt.

Reviewers:

(1) Leticia Morales-Ledesma, UNAM, Mexico.

(2) Valquíria Miwa Hanai Yoshida, University of Sorocaba, Brazil.

(3) Akinseye Olanrewaju Roland, Nigerian Institute for Trypanosomiasis & Onchocerciasis Research, Nigeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/77611>

Original Research Article

Received 02 October 2021

Accepted 06 December 2021

Published 13 December 2021

ABSTRACT

Aim: Our goal was to use the androgen and anabolic steroid (AAS) Testosterone Propionate (TP) to create a novel animal model to study polycystic ovaries.

Materials and Methods: Thirty albino female Wistar rats weighing 200 – 250 g were split into five groups with six rats in each group. Group I was treated as standard control, Group II was treated as the PCOS induced group, Group III was treated as a standard group, Clomiphene citrate (20 mg per kg body weight), intraperitoneally (I.P) along with a 4 mg TP injection intramuscularly (I.M) in an oily solution, Group IV was treatment control group treated with a hydro-alcoholic extract of *Caesalpinia bonducella* (CBHAE) at a dose of 200 mg per kg body weight. Group V was given CBHAE at 400 mg per kg of body weight. Blood collected from animals examined for hormonal parameters and ovaries is subjected to histopathological studies.

Results: There was a fall in testosterone volume and an increase in the efflux of female hormone constituents, which improved the ovarian development and helped to regulate menses.

Conclusion: Significantly lower levels of Testosterone were produced as a result of the extract, which also showed superior anti-androgenic properties. However, additional intervention is recommended in order to investigate the therapeutic effect of *Caesalpinia bonducella* seed in further detail.

Keywords: Anti-androgens; *Caesalpinia bonducella*; follicles; PCOS; tubular injury.

1. INTRODUCTION

Androgens in normal levels are beneficial, but in abnormal range (women), it leads to severe consequences i.e. forbid ovaries from releasing egg (ovulation), excessive hair growth, acne, Insulin resistance etc., eventually leads to PCOS.

Luteinizing hormone (LH) is a pituitary hormone released in response to Luteinizing hormone-Releasing hormone. It regulates the length and duration of the menstrual cycle, ovulation, implantation of a fertilized an egg in the uterus, secretion of estrogen and progesterone. Theca cells secrete testosterone in response to LH surge and in turn convert to estrogen by adjoining granulose cells. Excess LH stimulates ovulation, but a relative FSH shortage inhibits follicular development. FSH (3:1) stimulates the proliferation of ovarian theca cells, resulting in enhanced steroidogenesis and, ultimately, hyperandrogenism in PCOS women. *Caesalpinia bonducella* is abundant in furanoditerpenes beneficial in the management of PCOS [1]. Seeds comprised of phytochemicals including β -caesalpin, γ - caesalpin, caesalpin, citrulline and bonducellin, as well as fatty acid; stearic acid, palmitic acid, and oleic acid; amino acids such as arginine and aspartic acid [2-3]. Phytochemicals in *Caesalpinia bonducella* potent against a variety of PCOS attributes [4]. There are numerous documented medicinal properties like [5] anti-oxidant [6], adaptogenic [7], anti-inflammatory [8], anti-estrogen [9], and Immunomodulatory [10]. In order to appease the consequences anti-androgens came to light. In present study by using anti-androgenic property of *Caesalpinia bonducella* to appease PCOS in rodents.

2. MATERIALS AND METHODS

2.1 Collection of Plant Material

The marketed *Caesalpinia bonducella* was collected from an herbal production factory named ANNAIARAVINDHHERBALSlocated in 1, 2 & 3, Janaki Nagar, Maduravoyal, Chennai, and Tamil Nadu 600095.

2.2 Plant Authentication

The roots were recognized by professor P Jayaraman, Ph.D. from the plant anatomy and research center located at Tambaram (west). The authentication ID is PARC/2019/4111.

2.3 Extraction Procedure

Caesalpinia bonducella fresh fruits were collected and authenticated. After the seeds were rubbed out, the skin was left to dry in the shade. The shade-dried seeds mashed into a coarse powder and sieved through mesh no 60, were subjected to extraction.

500mg of coarsely powdered dried Grains of *Caesalpinia bonducella* was placed in a maceration flask with 70% of 2-liter Ethanol at 70°C temperature for 72 hours with occasional shaking. Then it was filtered and the macerate was evaporated to attain the yield. Finally the percentage of yield was calculated.

2.4 Experimental Model

2.4.1 Grouping of animal's

Female albino rats (180-220gm) were purchased from VETERINARY AND ANIMAL SCIENCES UNIVERSITY, TAMIL NADU, Laboratory Animal Medicine, and Centre for Animal Health Studies, Chennai- 51 the study was taken from an experimental facility, they were kept in a poly glycol boxes in a temperature-controlled (25±20°C) environment with a 12 hours dark/light cycle with standard ad libitum. All animals were acclimatized to the laboratory condition for a period of 15 days before the commencement of the study. The animals were divided into five groups, each of which had six animals.

2.4.2 PCOS Induction in the animals

The treatment of PCOS is evaluated in this study using TP-caused PCOS. [11] Thirty female virgin Wistar rats, weighing 180-220g of 10-12 weeks aged, having regular estrus cycles as determined by vaginal smear, were employed in the study.

[12]. Excluding the standard control group, all the other groups were treated with 4mg TP intramuscular injection in oily solution per rat. [13, 14] each animal had subjected to vaginal smear examination daily. The absence of cyclicity was shown as criteria for induction of PCOS by prolonged cornification of vaginal smears.

2.4.3 Vaginal smears

The stage of cyclicity was determined via microscopic examination of the primary cell type in vaginal smears. Estrous cycle cyclicity was assessed with 800 to 1200 hrs, and the relative components of epithelium, leukocytes, and cornified cells collected in daily vaginal lavages were studied using light microscopy. Which are known to alter at different stages of the estrous cycle. The control group or PCOS rats (proestrus, estrus, diestrus1, and diestrus2,) usually lasts about 4 days. [15]

2.4.4 Vaginal cytology

Every day vaginal smears were checked, while having lumbar support, the rats were confined at the upper ventral side and thorax. Cotton soaked in a drop of physiological saline solution was used to collect vaginal secretions. The cotton swab was inserted almost one or two inches into the female rat's vagina and rotated for two to three rotations. (Allowing the cotton swab end to collect an acceptable volume of cells), and the cotton tip was rolled along with the glass slide and the swab was gently removed. The dried smear was preserved by soaking it in 70% alcohol. After that, the slides were stained with methylene blue 0.5 % solution, cleansed with tap water and inspected under a light microscope with a $\times 10$ objective lens (without the condenser lens) [16].

2.4.5 Treatment Protocol

Except for I Group, which receives TP intramuscularly, all rats in Groups IV and V were given oral gavage for 15 days after induction. Whereas group 3 rats were treated with Clomiphene citrate Standard drug for a period of 15 days. On 16th day, Retro orbital puncture was performed to collect blood samples and serum was subjected to hormonal analysis (LH, FSH, estradiol, progesterone and testosterone). Animal in each group was selected and euthanized in carbon dioxide incubator.

Finally incision was made to collect the ovaries then weighed and subjected to histopathological examination.

2.4.6 Serum hormonal assay

LH, FSH, Serum testosterone, progesterone and estradiol, were measured using Enzyme-Linked immunoassay kit for quantitative estimation of respective hormones.

2.4.7 Histopathological examination

Bouin's solution was used to repair the removed ovaries and the ovaries were drained in a series of alcohol, and then cleansed in xylene before being immersed in heated paraffin wax at 60 degrees Celsius. They were dehydrated in a series of alcohol, and then cleaned in xylene by being embedded in paraffin wax melted at 60°C. Consecutive slices were placed on slide with 3-aminopropyl triethylsilane-coated sections were dried at 37°C for 24 hours. Before being mounted for histology, the slides were deparaffinized, hydrated, and stained with Mayer's hematoxylin & eosin dyes. Scope picture 3.0 imaging instrument (ScopeTek DCM 200) was used to examine the ovaries at a magnification of 40X (Hangzhou Scope tekOpto-Electric Co Ltd, USB 2.0). The thickness and diameter of cystic follicles were measured. Cystic follicles have a thicker, fibrotic cortex with a visible surface theca and interior layer. [17].

2.5 Statistical Analysis

The results are expressed as Mean \pm SEM. Data were evaluated using ONE WAY ANOVA followed by Newman – Keul's multiple range test. Probability values less than ($p < 0.01$) were considered significant.

3. RESULTS

3.1 Hormonal Parameter Evaluation

3.1.1 The effect of CBHAE on LH levels in rats with TP induced PCOS rats

When compared standard control group (G1), TP causes a considerable increase in LH levels & decrease in toxic control FSH (G2) levels ($P < 0.001$). The levels of LH/FSH differed considerably between control groups. When compared to toxic and a control group, extracted treated animals i.e., and both doses of CBHAE 200mg/kg & 400mg/kg exhibited a low level or

low levels of LH/FSH ratio and a remarkable drop in LH (P<0.01) & ascent in FSH levels (Tab: 2).

3.1.2 The effect of CBHAE in TP induced PCOS rats

Estradiol levels decline markedly when TP injections are given. A considerable rise in estradiol levels (P<0.01) result from simultaneous treatment for 15 days of CBHAE extracts. In standard group animals, estradiol levels are also significantly increased (Tab: 2)

3.1.3 The Effect of CBHAE on progesterone in TP induced PCOS rats

The drug induced group, which was given TP, has considerably decreased in progesterone levels. CBHAE treatment at both dosages in accordance with TP (200mg/kg & 400mg/kg) caused a substantial rise in progesterone (P < 0,001) levels to nearly normal levels. The control group found similar findings with CBHAE (Table: 2).

3.1.4 The effect of CBHAE on ovarian morphology

There was more cystic follicle in the ovaries of the positive control group (TP), but in 200mg/kg & 400mg/kg CBHAE shows near to normal follicles. Atretic follicles were seen and present at 200mg/kg. There were several healthy follicles in the 400 mg/kg group (Tab: 3)

3.1.5 The Effect of CBHAE on follicular diameter & thickness

The diameter and thickness of the cysts in the PCOS treatment group have increased but the standard group and extract treated group has fallen. (Tab: 3)

3.1.6 The Effect of CBHAE on ovarian weight

The TP. control group's ovarian weight fell significantly (P<0.01) as compared to the other groups, but it was restored to normal levels in the treatment groups 200mg/kg & 400mg/kg. (Table: 3).

G1 Antral follicles, Corpus luteum, Oocyte are surrounded by granulose cells. Theca cells is visible in this section, G2 Many cystic degenerating follicle and atretic follicles with degenerated granulosa layer is observed in an ovarian section from PCOS rat, G3 A normal follicle with clear antrum, and Oocyte in granulosa layer is visible in ovarian section of PCOS rat, treated with clomiphene citrate, G4 The ovary of a PCOS rat treated with a low dose of CBHAE 200mg/kg reveals mild degenerative follicle and absence of cystic, atretic follicle, G5 The ovary of a PCOS rat treated with a low dose of CBHAE 400mg/kg reveals the existence of growing regenerating follicles and corpus luteum, as well as Oocytes within granulosa and thicker theca.

Table 1. Percentage yield (w/w) of Plant extract

Sr No.	Extract	Color	Percentage yield (w/w)
1.	Hydro-alcoholic extract of <i>Caesalpinia bonducella</i>	Brownish Black	3.70

Table 2. Effect of CBHAE on serum hormone in TP induced PCOS

GROUP	FSH	LH	Estradiol	TSH	PRGSN
G1	8.27±0.25	6.146±0.33	54.12±2.31	0.28±0.04	14.15±0.61
G2	2.26±0.235**a	11.36±0.75**a	14.32±0.82**a	0.37±0.02**a	7.055±0.712**a
G3	7.02±0.46	5.25±0.38	46.49±1.74	0.32±0.02	12.12±0.29
G4	6.63±0.53**b	3.70±0.18**b	38.17±1.33**b	0.34±0.01**b	10.8±0.72*b
G5	6.13±0.56**b	4.57±0.21**b	41.18±0.92**b	0.32±0.01**b	11.60±0.76*b

All data's were analyzed G1-Normal, G2-Toxic, G3-Standard, G4-Low dose (CBHAE), G5-High dose (CBHAE)

For each group of 6 animals, all values represented as means ± SEM.

**a- values are significantly differ from Normal control (G1) at P<0.001

**b- values are significantly differ from PCOS control (G2) at P<0.001

*b- values are significantly differ from PCOS control (G2) at P<0.01

Table 3. Effect of CBHAE on ovarian morphology of PCOS rats

Dose mg.kg ovarian feature	Normal	Std control	Toxic control	Low dose	High dose
Atrectic follicle	0.00±0.00	1.12±0.05	4.43±0.307	3.06±0.17**b	0.05±0.11*b
Cystic follicle	0.00±0.00	3.7±10.58	10.55±1.21	0.00±0.00	0.00±0.00
Cystic follicle diameter	0.00±0.00	71.19±2.35	87.73±2.367	0.00±0.00	0.00±0.00
Cystic follicle thickness	0.00±0.00	34.63±2.18	42.35±1.48	0.00±0.00	0.00±0.00

G1-Normal, G2-Toxic, G3-Standard, G4-Lowdose, G5-High dose
 For each group of 6 animals, all values represented as means ± SEM.
 **b- values are considerably different from PCOS control (G2) at P<0.001
 *b- values are considerably different from PCOS control (G2) at P<0.01

3.2 Histopathological Examination




Plate. 1. G1: Normal Control (10ml/kg Normal Saline), G2: Drug Induced Group (TP 4mg/kg), G3: Standard Control (Clomiphene Citrate 20mg/ml), G4: Treatment Control (LOW DOSE of CBHAE 200mg/kg), G5: Treatment Control (High Dose of CBHAE 400mg/kg)

4. DISCUSSION

Although many models can be used to study PCOS, induction of PCOS by Testosterone Propionate can also be considered as one of the best model for studying PCOS. Hence, in terms of exhibiting the most reproductive and endocrine symptoms associated with PCOS, rodent PCOS models appear to parallel the human condition closely. This study investigated the effect CBHAE on the serum levels of LH, FSH, estradiol, testosterone & progesterone in TP induced PCOS. After 30 days of PCO induction, animals were analyzed for both hormonal & histologically, on 16th day after treatment with hydroalcoholic

extract, animals were also analyzed irrespective of their estrous cycle. In PCOS condition, normal gonadotropin-ovarian axis is disturbed results in hormonal imbalance reflected by the higher levels of LH, lower FSH levels and reversal of LH: FSH ratio. An elevated LH/FSH ratio and anovulation are typical findings in women with PCOS [22,23]. The extract treated groups show better reduction in this LH/ FSH ratio, indicating that extract could reverse PCOS condition. Oestrogen similar to other steroids becomes altered in PCOS [24]. Repetitive administration of CBHAE led to significant rise in estradiol. Similarly, the reduction in the level of progesterone in the PCOS-induced animals could be responsible for

the persistent oestrus phase. Elevation in the concentration of serum progesterone by CBHAE may be responsible for the reversal of the luteal phase dysfunction and restoration of normalcy of the estrous cycle.

Our study showed that CBHAE induced an increase in serum estradiol implies that plant causes marked improvement in endocrine function and recovery of ovulatory functions in the rats. Hyperandrogenism (as a result of high testosterone levels) which is evident in human PCOS.[25, 26] was not present in this animal model of TP induced PCOS [27]. Therefore no effect of the extract on androgen levels was observed using this model of PCOS induction. Ovarian weight in PCOS induced rats was more than the normal rats which is in accordance with earlier findings. [28-30] Treatment with CBHAE prevented further increase in ovarian weight & returned to normalcy. The biochemical results are also supported by histopathological observation of light microscopy. The histomorphometry of PCOS was a suitable measurement for describing the cystic status because differences were observed in the morphological characters and in the presence or absence of follicular cysts. It is reported that the histopathological study of PCOS induced rats shows the formation of poly cysts in the ovary.[31,32]. Ovaries exhibited increased follicle atresia and multiple cysts with thin granulosa cell layers and thickened theca cell layers.[33] After treatment with extract of *Caesalpinia bonducella* PCOS condition was reversed, number of cystic follicles reduced & found numerous healthy follicles at a different stages of development. CBHAE indicates that treatment group shows marked recovery of ovarian tissue and the animals may probably be preparing for ovulation.

After treatment with CBHAE, the incidence of cystic follicles was reduced, and many healthy follicles at various stages of development were detected. CBHAE indicates that the therapy group has made substantial progress, and the ovarian tissue and animals are most likely preparing for ovulation. In addition, CBHAE therapy restores gonadotropin feedback inhibition (LH & FSH), resulting in a rise in estradiol and progesterone. The extracts capability for reducing the histological, clinical, and biochemical aspects of the PCOS indicates all of these variables. The presence of flavonoids may explain the pharmacological effect of CBHAE extracts. Aside from the flavonoids discovered to have an antioxidant function in PCOS rats, the PCOS

illness has been linked to a reducing in the quantity of antioxidant enzyme/molecule. Antioxidants aid in protecting the human body from the negative effects of reactive oxygen species (ROS) (ROS). Plants with a strong antioxidant characteristic, particularly flavonoids, have shown themselves. The well-known usage of plants in the past in the management of gynecological disorders may be responsible for these plant phytochemicals.

5. CONCLUSION

PCOS is a prevalent endocrine disorder of women, which can result in infertility. The prospective function of Herbal medicine in the treatment of PCOS is consistent with few adverse effects. Herbal medicine improves body immunity without changing hormonal level and also regularise menstrual cycle. In conclusion CBHAE reveals a considerable recovery of PCOS animals from, LH, FSH, estradiol & progesterone as well as the irregular cycle and ovarian physiology. TP injection enhances the ovary weight and reproductive system of normal rats, which returned to normal, after treating with CBHAE. Polycyst's were seen as a group exception of the normal control group, which has no polycysts at all. CBHAE was found to correct hormonal imbalance caused by TP in PCOS rats, indicating that it might be used to treat infertility.

SUPPLEMENTARY MATERIALS

Supplementary Materials available in this link: <https://www.journaljpri.com/index.php/JPRI/libraryFiles/downloadPublic/23>.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The research has been reviewed and approved by Institutional animal ethical committee and

was carried out in accordance with CPCSEA guidelines (approval number IAEC/219/2019).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Pascoe, Keith O, Basil A. Burke, and Wilfred R. Chan. "Caesalpin F: a new furanoditerpene from *Caesalpinia bonducella*." *Journal of Natural Products*. 1986;49.5:913-915. Available: <https://doi.org/10.1021/np50047a025>
- Prasad, GoliPenchala, et al. "Analytical study of Kuberaksha/ Kantakikaranjapatra Churna [*Caesalpinia bonduc* (L.) Roxb. leaf powder]." *Ayu*. 2010;31.2:251.
- Ramadurai, Sivasankari, and Usha Balasundaram. "Rhizomicrobiomics of *Caesalpinia bonducella*, a wonder plant for PCOS treatment." *Physiology and Molecular Biology of Plants*. 2020;26.12:2453-2463.
- Kandasamy, Veerapandiyan, and Usha Balasundaram. "*Caesalpinia bonduc* (L.) Roxb. As a promising source of pharmacological compounds to treat Polycystic Ovary Syndrome (PCOS): A review." *Journal of Ethnopharmacology*. 2021:114375.
- Shukla, Shruti, et al. "Antioxidant activity and total phenolic content of ethanolic extract of *Caesalpinia bonducella* seeds." *Food and Chemical Toxicology*. 2009;47.8:1848-1851.
- Kannur DM, Hukkeri VI, Akki KS. "Adaptogenic activity of *Caesalpinia bonducella* seed extracts in rats." *Journal of Ethnopharmacology*. 2006;108.3: 327-331.
- Shukla, Shruti, et al. "Studies on anti-inflammatory, antipyretic and analgesic properties of *Caesalpinia bonducella* F. seed oil in experimental animal models." *Food and Chemical Toxicology*. 2010;48.1:61-64.
- Biswas TK, et al. "Oral hypoglycemic effect of *Caesalpinia bonducella*." *International Journal of Pharmacognosy*. 1997;35.4: 261-264.
- Kannur, Dayanand M. "Antidiabetic and antihyperlipidemic activity of *Bonducella* (*Caesalpinia bonducella*) seeds." *Nuts and Seeds in Health and Disease Prevention*. Academic Press. 2011:237-244.
- Shukla, Shruti, et al. "In vivo immunomodulatory activities of the aqueous extract of bonduc nut *Caesalpinia bonducella* seeds." *Pharmaceutical Biology*. 2010;48.2:227-230.
- Alaa El-Din AB, Vern JC, Michelle M, Jenny HZ, Richard S. Antioxidant activities, sensory and anti-influenza activity of grape skin tea infusion. *Food Chemistry*. 2011;129: 837–845.
- Dehghani F, Panjehshahin MR, Vojdani Z. The Effect of hydroalcoholic extract of caraway on thyroid gland structure and hormones in female rat. *Iranian Journal of Veterinary Research, Shiraz University*. 2010;11(4):337-341.
- Sunanda P, Anand K. studied the anti-thyroid action of leaf extract of *Annona squamosa* in albino rats. *Current Science*. 2003;84(11):1402-1404.
- Kar A, Panda S, Bharti S. The Relative efficacy of three medicinal plant extracts in the alteration of thyroid hormone concentrations in male mice. *Journal of Ethnopharmacology*, 2002;81:281-285.
- Panda S, Jafri M, Kar A Meheta BK. The Thyroid inhibitory, antiperoxidative and hypoglycemic effects of stigmasterol isolated from *Butea monosperma*. *Fitoterapia*. 2009;80:123-126.
- Jean-Marc S, Franck B, Michel M. The Polymeric proanthocyanidins from grape skins. *Phytochemistry*, 1996;43(2):509-512.
- Sourgens H, Winterhoff H, Gumbinger HG, Kemper FH. Antihormonal effects of plant extracts, TSH and prolactin-suppressing properties of *Lithospermum officinale* and other plants. *Planta Med*. 1982;45:78-86.
- Kim SM. Effects of ginseng radix on the rat hypothyroidism induced by PTU (6-n-propyl-2-thiouracil). *Korean Journal of Herbology*. 2010;25(3):11–18.
- Kavutcu MO, Canbolat ztu"rk SO" et al. Reduced enzymatic antioxidant defense mechanism in kidney tissues from gentamicin-treated guinea pigs: effects of vitamins E and C. *Nephron*. 1996;72(2): 269–274.
- Jamall IS, Smith JC. Effects of cadmium on glutathione peroxidase, superoxide dismutase, and lipid peroxidation in the rat heart: a possible mechanism of cadmium cardiotoxicity. *Toxicology and Applied Pharmacology*. 1985;80(1):33–42.

21. Sedlak J, Lindsay RH. Estimation of total, proteinbound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical Biochemistry*. 1968;25(1):192–205.22.
22. Ivana N, Patricia J, Marijan S, Ana MOB. Ultrasoundextracted flavonoids from four varieties of Portuguese red grape skinsdetermined by reverse-phase high-performance liquid chromatographywith electrochemical detection. *AnalyticaChimica Acta*. 2008;630: 107–115.
23. Mukesh Y, Shalini J, Aarti B, Ravinder N et al. Biologicaland Medicinal Properties of Grapes and Their Bioactive Constituents:An Update. *Journal Of Medicinal Food*, 2009;12(3):473–484.
24. Panda S, Kar A. Amelioration of L-thyroxine-inducedhyperthyroidism by coumarin (1, 2-benzopyrone) in female rats.*Clinical and Experimental Pharmacology and Physiology*, 2007;34(11):1217–1219.
25. Seong-Mo K, Sang-Chan K, In-Kwon C, Woo-Hyun C, SaeK. Antioxidant and Protective Effects of Bupleurum falcatum on the L-Thyroxine-Induced Hyperthyroidism in Rats. *Evidence-Based Complementary and Alternative Medicine*. 2012;1- 12.
26. Lei Z, Shelly H, Jianrong L, Shi S, Corene C, Shao JZ, andKequan Z. Grape skin extract inhibits mammalian intestinal α -glucosidase activity and suppresses postprandial glycemic response instreptozocin-treated mice. *Food Chemistry*. 2011;126:466–471.
27. Tahiliani P, Kar A. The combined effects of Trigonella andAllium extracts in the regulation of hyperthyroidism in rats.*Phytomedicine*. 2003;10(8): 665–668.
28. Subudhi KUD, Paital B, Bhanja S, Chainy GBN. Alleviationof enhanced oxidative stress and oxygen consumption of L-thyroxineinduced hyperthyroid rat liver mitochondria by vitamin E andcurcumin,. *Chemico- Biological Interactions*; 2008,
29. Kim SM. Effects of ginseng radix on the rat hypothyroidism inducedby PTU (6-n-propyl-2-thiouracil). *Korean Journal of Herbology*. 2010;25(3):11–18.
30. Kavutcu MO, Canbolat, ztu'rk SO" et al. Reduced enzymaticantioxidant defense mechanism in kidney tissues from gentamicin-treated guinea pigs: effects of vitamins E and C. *Nephron*, 1996; 72(2):269–274.
31. Jamall IS, Smith JC. Effects of cadmium on glutathione peroxidase, superoxide dismutase, and lipid peroxidation in the rat heart: a possiblemechanism of cadmium cardiotoxicity. *Toxicology and Applied Pharmacology*.1985;80(1):33–42.
32. Sedlak J, Lindsay RH. Estimation of total, proteinbound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical Biochemistry*. 1968;25(1): 192–205.
33. Aebi H. Catalase, in *Methods in Enzymatic Analysis*, H. U.Bergmeyer, Ed., Academic Press, New York, NY USA. 1974:673–686.
34. Sun LY, Oberley W, Li Y. A simple method for clinical assay ofsuperoxide dismutase. *Clinical Chemistry*. 1988;34(3): 497–500..
35. Balu MSP, Haripriya D, Panneerselvam C. Rejuvenation ofantioxidant system in central nervous system of aged rats by grape seedextract. *Neurosci. Lett*. 2005; 383:295-300.

© 2021 Sumithra et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77611>