



Level of Interferon - Gamma, Interlukin – 6, Frequency of Monocyte and Lymphocyte Subsets among Newly Diagnosed Tuberculosis Patients in Northwest Ethiopia

Mulualem Lemma^{1,2*}, Tadelo wondmagegn^{1,2}, Bewketu Mengesha³, Sheferaw Yehuala⁴, Meseret Workineh¹ and Ebba Abate^{1,5}

¹Department of Immunology and Molecular Biology, School of Biomedical and Laboratory Science, College of Medicine and Health Science, University of Gondar, Ethiopia.

²Department of Medical Laboratory Services, University of Gondar Hospital, Ethiopia.

³Leishmaniasis Research and Treatment Center, University of Gondar Hospital, Ethiopia.

⁴Department of Statistics, University of Gondar, Ethiopia.

⁵Ethiopian Public Health Institute, Ethiopia.

Authors' contributions

This work was carried out in collaboration between all authors. Authors ML and EA conceived and designed the experiments. Authors ML, MW and BM carried out data collection and laboratory analysis. Author EA supervised the data collection and laboratory analysis. Authors ML and SY analyzed the data and prepared the first manuscript draft. Authors EA, MW and ML reviewed the draft. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMB/2019/44844

Editor(s):

(1) Dr. Adekunle Sanyaolu, Epidemiology Division, Nigeria Center for Disease Control, Federal Ministry of Health, Abuja, Nigeria.

Reviewers:

(1) Alok Nahata, Ying Zhi Agricultural and Industries Sdn Bhd, Malaysia.

(2) Nagahito Saito, Nemuro City Hospital, Japan.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/44844>

Data Article

Received 02 October 2018
Accepted 15 December 2018
Published 15 January 2019

ABSTRACT

Background: Tuberculosis is one of the fundamental health problems in the world. During TB infection immune cells and cytokines play the major role for the pathogenesis of the disease and also for host defence. Cells of immune system play a great role to decrease the progress of the disease with the coordination of cytokines. The aim of our study was to determine the level of immune cell and certain cytokine in TB patients and apparently health controls.

*Corresponding author: E-mail: mulualemi@yahoo.com;

Methods: A comparative cross-sectional study was conducted among newly diagnosed TB patients in Northwest Ethiopia. Blood samples were collected from all study participants. Serum level of interferon-gamma and interleukin-6 were determined by using an Enzyme Linked Immunosorbent Assay (Ready-SET-Go! Kit), Frequency of Monocyte and T cell sub population were measured with Flow cytometry using BD Multi TEST™ reagent. Mann-Whitney U test and spearman correlation were applied for statistical test by using graph pad prism.

Results: A total of 80 study participants comprising of 40 newly diagnosed smear positive TB patients and 40 healthy blood donors were included in the study. From our study participants 19 (23.75%) were females and 61(76.25%) were males with mean age of 30. Results from the study showed that the median level of IFN- γ (72.8) and IL-6 (13.45) in TB patients was higher when compared with controls IFN- γ (44.8) & IL-6 (4.08) ($P > 0.001$) while on the other hand, the median level of CD4 (565.5) and BMI (18.00) were lower compared to controls CD4 (857) & BMI (20.9) ($P > 0.001$).

Conclusion: Our study showed high level of IFN- γ , IL-6 and low level of monocyte, CD4 and BMI in TB patients than the control groups.

Keywords: Tuberculosis; CD4T cell; IFN- γ and IL-6; gondar; Ethiopia.

ABBREVIATIONS

BMI : body mass index
CD : cluster of differentiation
EDTA : Ethylenediaminetetraacetic acid
ELISA : Enzyme linked immunosorbant assay
IL : Interlukine
IFN- γ : Interferon gamma
TB : Tuberculosis
TNF- α : Tumor necrosis factor alpha
SD : Standard deviation
WHO : world health organization

1. INTRODUCTION

Tuberculosis (TB) continues to be major public health problem, causing global morbidity and death yearly. Based on the recent WHO report, there were 9.6 million new TB cases per year and close to 2 million deaths [1]. Factors related to the environment, host and pathogen could play significant roles for TB susceptibility [2]. Immune status of the patient play a great role in the progression of TB and for the control of TB disease [3]. The ability of the host to elicit potent cell mediated immune response determines disease development and outcomes associated with TB. Macrophage activation, cytokine production and other immune cells play roles in TB immunity and determine the outcome of the disease [4,5].

T-helper 1(Th1) and T-helper 2 (Th2) immune responses are the major arms of immune responses for humoral and cell mediated immune system, respectively. The induction of the Th1 type immune response is very crucial to

combat TB infection and characterized by secretion of interferon-gamma (IFN- γ). Cytokine such as IFN- γ , Tumor necrosis factor – alpha (TNF- α), IL-12 and Interleukin-6 (IL-6) are responsible chemicals to build the proactive immunity against TB pathogens during early stage of the infection. IFN- γ has versatile roles including facilitating the intracellular killing of the TB bacilli by macrophages [6,7]. It has a great role for the function and maturation of immune cells, and it stimulates production of other cytokines and activation factors to augment the function of innate immune system during infections [8,9]. In addition to IFN- γ , the inflammatory cytokines like IL-6 also play pivotal roles in TB immunity. The secretion of IFN- γ for the activation of macrophage by CD4 cells during TB infection is stimulated by the production of IL-6 [10]. IL-6 is a cytokine known as a kind of pleiotrophic nature and produced by many cells including T and B-lymphocytes. It plays a great role in the synthesis of C-reactive protein in various stimuli [11,12] and also has been revealed that it has a relevant role in the pathogenesis of tuberculosis like the secretion of IFN- γ during infection [10,13].

Therefore, the present study aims to assess the levels of these two essential cytokines IL-6 and IFN- γ , which are greatly needed to combat TB infection and as well as assessing the levels of two immune cells which are T cells and monocyte. Thus, the level of these two cytokines and cells among active TB patients will be compared to the level from apparently healthy control.

2. MATERIALS AND METHODS

2.1 Study Design and Area

A cross-sectional study was conducted in newly diagnosed TB patients at the Gondar of University referral hospital from February to May 2014. Gondar University hospital is a tertiary level teaching and referral hospital with 450 beds for inpatients as well as rendering referral health services for over 5 million inhabitants in North-West Ethiopia. The hospital provides inpatient and outpatient services to the population in the surrounding area of Gondar town and the adjacent regions. The hospital has TB clinic where TB patients are getting their medication and further assessment during follow up period. For comparison purpose, apparently healthy controls were enrolled at the Gondar University Blood Bank.

2.2 Study Participant

A total 80 study participants comprising of 40 newly diagnosed smear positive TB patients and 40 healthy blood donors were selected for the study; the mean age for the TB patients and the controls were 34 years (range 18 to 57) and 26 Years (range: 20 to 42), respectively. Number of females among TB patients is (11/40) 27.5% and (9/40) 22.5% among the control group. Among the TB cases, 70% (28/40) were diagnosed in the hospital as pulmonary TB and the remaining were extra pulmonary TB. The body mass index (BMI) of the study participants were, 17.98 for Tb patients and 20.64 for controls. From all study participants' informed consent were taken. Participants with known disease (HIV, diabetic, hypertensive and cancer) and pregnant women were excluded in the study and we include with the age group of 18 up to 65 years old. Socio-demographic data were collected using a structured and pretested questionnaire.

2.3 Blood Collection

Six (6) ml of whole blood was collected from the study participant using serum separator [14] tubes and EDTA for measurement of cytokines using ELISA technique and the remaining one was used for blood cell count determination, respectively.

2.4 Serum Level IFN- γ and IL-6 Measurement

The levels of IL-6, and IFN- γ were measured using ELISA Ready-SET-Go! Kit. Anti-human

cytokines (IL-6 and IFN- γ) capture antibodies (eBioscience) was coated in ELISA (Coat corning costar 9018) plates with 100 μ l/per well in 1x coating buffer and the plates were sealed and incubated overnight at 2-8°C. On the following day, the plates were aspirated and washed three times using 250 μ l wash buffer (1x PBS, 0.05% Tween 20). The standards (human cytokines lyophilized standard) were reconstituted and diluted according to the given instruction. A twofold serial dilution of the top standards with 100 μ l were performed in duplicates to make a standard curve for 8 points and also 100 μ l supernatant of samples were added to incubate at room temperature for 2 hours. After 3-5 washes, 100 μ l/well of detection antibody (anti human cytokine biotin) was added and incubated at room temperature for 1 hour. After 3-5 washes, 100 μ l/well of Avidin-HRP was added and incubated at room temperature for 30 minutes. Finally, 1M H₃PO₄ or 2N H₂SO₄ stop solution was added to stop the reaction, and the plate was read at 450 nm length using ELISA reader. The analytical sensitivity and other technical protocols were followed as per the manufacturer guideline.

2.5 Cell Counting Using Flow Cytometry

T-lymphocyte sub-populations were characterized using Multi TEST™ fluorescent labelled monoclonal antibodies reagents against surface CD markers (CD₃ fluorescein isothiocyanate (FITC), CD₈ phycoerythrin (PE) and CD₄allophycocyanin (APC) after erythrocyte has been lysed. FACS Calibur flow cytometer (Becton Dickinson Biosciences, San Jose, CA, USA) was used for acquisition and analysis of T-lymphocyte sub-population by using Multit TEST software (Becton Dickinson), and a minimum of 15, 000 events were captured during acquisition. Monocyte cell count done by coulter counter machine.

2.6 Statistical Analysis

Data were analyzed using graph pad prism. Data were reported as frequency and percentage for categorical variables and mean and standard deviation (SD). Mann-Whitney U test was used to compare the cytokine levels between TB patients and healthy controls. Spearman correlation was used to correlate serum IFN- γ and IL-6 cytokines with peripheral CD4+, CD8+ and Monocyte cells. Statistical significance was considered at 95% level of confidence and P value less than 0.05.

3. RESULTS

3.1 IFN-gamma, IL-6, and Peripheral T-cell Subpopulation and Monocyte

The level IFN- γ and IL-6 were significantly higher among active TB patients compared to healthy controls (median of 72.8 Pg/ml versus 44.0 pg/ml, $p < 0.0001$ for IFN-gamma; and 27.4 pg/ml versus 6.0 pg/ml, $p < 0.0001$ for IL-6) (Table 2). Similarly, significant difference in CD4⁺ T cell subsets ($p < 0.0001$) and monocyte ($p=0.018$)

was obtained between active TB patients and healthy controls (Table 1) & (Fig. 1).

3.2 Correlation of Serum IFN- γ and IL-6 Cytokines with Peripheral CD4⁺, CD8⁺ and Monocyte Cells

By using spearman correlation among cytokine and cells, IL-6 and CD4⁺ showed weak negative correlation ($r = - 0.4929$; $p = 0.001$) but monocyte and CD8⁺ cell showed weak positive correlation ($r = 0.3962$; $p = 0.0167$) (Fig. 2).




Fig. 1. Serum level of IFN-gamma, IL-6, and measurement of peripheral blood CD4⁺ cells, CD8⁺ cells and Monocytes among newly diagnosed TB patients (n=40) and apparently healthy controls (n=40)




Fig. 2. Correlation of serum level cytokine with peripheral cell in newly diagnosed TB patients (A) IFN- γ with CD4⁺ cells, (B) IFN- γ with IL-6, (C) IFN- γ with monocyte , (D) IFN- γ with CD8⁺, (E) IL-6 with CD4⁺ cells, (F) IL-6 with CD8⁺ cells (G) IL-6 with monocyte cells and (H) CD8+ cell with monocyte cells

Table 1. Level of IFN- γ , IL- 6, CD4T cell, CD8 T cell and Monocyte cell in TB patients and controls

Parameters	TB patients	Control	p-value
IFN- γ (pg/ μ l) (median \pm SD)	72.8	44.0	< 0.0001
IL-6 (pg/ μ l) (median \pm SD)	13.45	4.05	< 0.0001
CD4 cells($\times 10^3$ / μ l)	565.5	857.0	< 0.0001
CD8 cells($\times 10^3$ / μ l)	539.5	551.5	< 0.4162
Monocyte ($\times 10^3$ / μ l)	400.0	300.0	< 0.0193

4. DISCUSSION

In our study, the level of CD4 cell count is significantly higher in the controls when compared with that of newly diagnosed TB-patients. This is in agreement with previously literature which showed that patients with TB case had low level of CD4⁺T cell subsets when compared with that of the apparently health controls [14]. Due to the protective effect of the immune response against this pathogen mediated by Th1 cells, lower levels of the cells leads to the progression of tuberculosis in the patients [3]. Moreover, most of TB cases show low BMI in our study, which may play a great role for the decreasing of CD4 counts and their functionality by reducing the transportation of different co-factors and elements than controls with normal BMI value. This Low level of CD4 cells play a great role for the pathogenesis of TB disease, because it may be due to the low respond in delayed type immune response and due to low level of cell frequency may lead to decrease in granuloma formation, which is fundamental for proper immune response.

IFN- γ is one of the Th1 cytokine identified as the most important agents of anti-mycobacterial cytokine by activating macrophage to enhance intracellular killing [15,16] as well as enhance the production of other cytokine [17]. In our study the level of IFN- γ was significantly higher in TB patients compared with the control groups. The finding is consistent with other similar studies on TB [18-20] and the level of this cytokine is decrease through and after treatment [20] The probable reason why INF- γ is higher in TB patients while CD4 cell are low is perhaps due to the fact that IFN- γ comes out from both local production and spill over of it from the activated lymphocytes sequestered at the site of MTB infection. This condition can make the cytokine not being able to elicit downstream events involving ineffective activation of macrophages and intracellular killing of MTB. All these conditions play a great role for the pathogenesis of MTB infections.

The level of IL- 6 in TB patients is also higher than that of the controls, similar studies had the result which is in line with our findings [21,22]. On the other hand also study showed that there is increased IL- 6 in active TB disease condition when compared with that of the latent condition of TB infection [23]. In addition study showed that the concentration of IL-6 is very high in TB patients with pulmonary cavities than without cavities. This is a great indication that IL-6 plays its own role for the TB disease outcome [24]. High level of IL-6 during TB infection can inhibit the type 1 interferon signalling on macrophage and consequently lead to the disease progression [25]. Moreover , increased level of IL-6 is correlated with disease severity and decrease in the nutritional status of the patients [24], our findings showed this based on the level of BMI in TB patients in the study that were significantly lower than that of health controls.

The level of monocyte cell in the blood of TB patients is higher than that of the controls and show significant association, one of the probable reason that the increased frequency of the monocyte during TB infection is due to the production of pro-inflammatory cytokines and chemo attractant factors that direct the monocyte from the bone marrow to lymph node and other infected tissue where it act like macrophages. In this condition there may be an increase of monocyte in the vascular. On the other hand cytokines that is produced by infected macrophages may have an impact on monocyte by not changing it to macrophages thereby preventing it from going to the infected area. This high frequency of monocyte in TB patients may also contribute to the increased level of IL- 6 in TB patients since they have high level than the controls.

5. CONCLUSION

In our study patients with tuberculosis showed higher level of IFN-Y, IL-6 and monocyte with lower level of CD4 and BMI than the controls.

This may indicate the progression of the disease. Future studies are required to describing the mechanisms of this value alteration during TB disease as well as for better clinical application of these parameters.

CONSENT AND ETHICS APPROVAL

The study was approved by the ethical committee of the School of Biomedical and Laboratory Sciences, University of Gondar. In addition, written informed consent was obtained from each participant.

AVAILABILITY OF DATA AND MATERIALS

The datasets supporting the conclusions of this article are included within the article.

ACKNOWLEDGEMENT

We would like to thank Yohans Sink, Eden W/gerima, Debaka Belet and Samirawit Sisay in the form university of Gondar hospital HIV laboratory. And also Esay Derebe Asnkew Belet Tigist, Koku Sisay, Nebyu Mekonn from University of Gondar hospital TB clinic and laboratory for their indispensable cooperation. Finally we would like to acknowledge University of department of Immunology and molecular biology for great cooperation to finalize this work.

COMPETING INTERESTS

The authors declare that there is no competing interest with regard to the present study.

REFERENCES

1. Organization WH. Global tuberculosis report 2015: World Health Organization; 2015.
2. McShane H. Susceptibility to tuberculosis—the importance of the pathogen as well as the host. *Clinical & Experimental Immunology*. 2003;133(1):20-1.
3. Kinjo Y, Kawakami K, Uezu K, Yara S, Miyagi K, Koguchi Y, et al. Contribution of IL-18 to Th1 response and host defense against infection by Mycobacterium tuberculosis: A comparative study with IL-12p40. *The Journal of Immunology*. 2002; 169(1):323-9.
4. Munk M, Emoto M. Functions of T-cell subsets and cytokines in mycobacterial infections. *The European Respiratory Journal Supplement*. 1995;20:668s-75s.
5. Jo EK, Park JK, Dockrell HM. Dynamics of cytokine generation in patients with active pulmonary tuberculosis. *Current opinion in infectious diseases*. 2003;16(3):205-10.
6. Kaufmann SH. Is the development of a new tuberculosis vaccine possible? *Nature medicine*. 2000;6(9):955-60.
7. Mosmann T, Coffman R. TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties. *Annual review of immunology*. 1989;7(1):145-73.
8. Miller CH, Maher SG, Young HA. Clinical use of interferon- γ . *Annals of the New York Academy of Sciences*. 2009;1182(1):69-79.
9. Mootoo A, Stylianou E, Arias MA, Reljic R. TNF- α in tuberculosis: A cytokine with a split personality. *Inflammation & allergy-drug targets (Formerly Current Drug Targets-Inflammation & Allergy)*. 2009; 8(1):53-62.
10. Saunders BM, Frank AA, Orme IM, Cooper AM. Interleukin-6 induces early gamma interferon production in the infected lung but is not required for generation of specific immunity to mycobacterium tuberculosis Infection. *Infection and immunity*. 2000;68(6):3322-6.
11. Papanicolaou DA, Wilder RL, Manolagas SC, Chrousos GP. The pathophysiologic roles of interleukin-6 in human disease. *Annals of Internal Medicine*. 1998;128(2): 127-37.
12. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *New England Journal of Medicine*. 1999;340(6):448-54.
13. Ladel CH, Blum C, Dreher A, Reifenberg K, Kopf M, Kaufmann S. Lethal tuberculosis in interleukin-6-deficient mutant mice. *Infection and immunity*. 1997;65(11): 4843-9.
14. Skogmar S, Schön T, Balcha TT, Jemal ZH, Tibesso G, Björk J, et al. CD4 cell levels during treatment for tuberculosis (TB) in Ethiopian adults and clinical markers associated with CD4 lymphocytopenia. *PLoS One*. 2013;8(12):e83270.
15. Lin PL, Flynn JL. Understanding latent tuberculosis: a moving target. *The Journal of Immunology*. 2010;185(1):15-22.
16. Cooper AM, Khader SA. The role of cytokines in the initiation, expansion, and control of cellular immunity to tuberculosis.

- Immunological Reviews. 2008;226(1):191-204.
17. Fulton SA, Johnsen JM, Wolf SF, Sieburth DS, Boom WH. Interleukin-12 production by human monocytes infected with Mycobacterium tuberculosis: Role of phagocytosis. Infection and Immunity. 1996;64(7):2523-31.
 18. Liu Y, Janeway CA. Interferon gamma plays a critical role in induced cell death of effector T cell: A possible third mechanism of self-tolerance. Journal of Experimental Medicine. 1990;172(6):1735-9.
 19. Katti MK. Assessment of serum IL-1, IL-2 and IFN- γ levels in untreated pulmonary tuberculosis patients: role in pathogenesis. Archives of Medical Research. 2011; 42(3):199-201.
 20. Lee SW, Lee CT, Yim JJ. Serial interferon-gamma release assays during treatment of active tuberculosis in young adults. BMC Infectious Diseases. 2010; 10(1):300.
 21. Lopes FHA, Assis LCd, Pires Neto RdJ, Botelho KP, Sá KM, Frota CC, et al. Serum levels of interleukin-6 in contacts of active pulmonary tuberculosis. Jornal Brasileiro de Patologia e Medicina Laboratorial. 2013;49(6):410-4.
 22. Correia J, Freitas M, Queiroz J, Pereira Perrin M, Cavadas B. Interleukin-6 blood levels in sensitive and multiresistant tuberculosis. Infection. 2009;37(2):138-41.
 23. Djoba Siawaya J, Beyers N, Van Helden P, Walzl G. Differential cytokine secretion and early treatment response in patients with pulmonary tuberculosis. Clinical & Experimental Immunology. 2009;156(1): 69-77.
 24. Karyadi E, Dolmans WM, West CE, Van Crevel R, Nelwan RH, Amin Z, et al. Cytokines related to nutritional status in patients with untreated pulmonary tuberculosis in Indonesia. Asia Pacific journal of clinical nutrition. 2007;16(2):218-26.
 25. Martinez AN, Mehra S, Kaushal D. Role of interleukin 6 in innate immunity to Mycobacterium tuberculosis infection. The Journal of Infectious Diseases. 2013; 207(8):1253-61.

© 2019 Lemma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/44844>*