



A Review on Ethnobotanical Study of Traditional Medicinal Plants Used for Treatment of Liver Problems in Ethiopia

Belachew Garedew^{1*} and Behailu Bizuayehu¹

¹*Department of Biology, College of Natural and Computational Science, Wolkite University, Wolkite, Ethiopia.*

Authors' contributions

The authors conceptualized the studies, wrote and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2018/38153

Editor(s):

- (1) Dr. Paola Angelini, Department of Applied Biology, University of Perugia, Italy.
(2) Dr. Marcello Iriti, Professor, Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

- (1) M. Sasikala, Karpagam College of Pharmacy, India.
(2) Emmanuel Nachamada Solomon, Ahmadu Bello University, Nigeria.
Complete Peer review History: <http://www.science domain.org/review-history/27534>

Review Article

Received 03 September 2017
Accepted 06 December 2017
Published 01 December 2018

ABSTRACT

Traditional medicinal plants are used for human ailments treatment throughout the world primary health care need practiced typically liver problems by indigenous medical practitioners. This review was employed with the objective of compiling and documenting ethnobotanical studies on the use of medicinal plants, associated with the treatment of liver problems from different ethnic groups in Ethiopia. By using published and unpublished research articles, a total of 46 journals meeting inclusion criteria were used for analyzing and compiling this review literature. The ethno-medicinal use of 114 plant species belonging to 90 genera and 50 families were reported and documented from eight regions of Ethiopia. The highest family in terms of species number was Euphorbiaceae accounts 13 (11.4%) species and followed by Asteraceae 12 (10.53%) species and Fabaceae 10 (8.77%) species. Herbs were dominant habit 48(42.11%) followed by shrubs 33 (28.95%). Most of the medicinal plant species (52.7%) were collected from the wild. The most frequently cited plant species were *Justicia schimperiana* 9 (12.5%) followed by *Phytolacca dodecandra* 8 (11.11%), *Croton macrostachyus* 7 (9.72%). The most frequently utilized plant parts for treatment of liver disease was root (24.78%) followed by leaf (23.89 %). The medicinal plant preparations were administered via oral gavage or Oral route of administration was employed for the medicinal plant

*Corresponding author: E-mail: belachewgar@gmail.com, begaredew@gmail.com;

preparations most commonly used route of application 99 (88.39%) followed by dermal and oral 4 (3.57%) and dermal application 3 (2.68%). The healers used different method of preparation for application such as crushing, concoction, decoction, chewing, powdering, etc. This review indicated that the liver problem is common disease in Ethiopia. Adapting a recommended diagnostic and treatment using physical diagnosis by indigenous healers/ practitioners attempted curing liver problem implementing prevention and control policies in the general population needs an urgent attention in the country.

Keywords: Ethnobotany; indigenous knowledge; liver problems; medicinal plants.

1. INTRODUCTION

Traditional medicine is used throughout the world as it is dependent on locally available plants, which are easily accessible, and capitalizes on traditional wisdom-repository of knowledge, simple to use and affordable in cost compared to modern medication of western [54,56]. These medical systems are heavily dependent on various plant species and plant based products. Traditional systems of medicine are popular in developing countries and up to 80% of the population relies on traditional medicines or folk remedies for their primary health care need practiced by indigenous practitioners [47]. In Ethiopia, most of the human population is dependent on traditional medicine plants for primary healthcare services or human ailments treatment [41].

Most studies included surveys of utilization of medicinal plants for the treatment different diseases in different parts of Ethiopia, as recommended by traditional healers /practitioners [52]. Various types of plants identified in Ethiopia claimed to be used for treatment of liver problems. Liver is the largest organ in the body. Liver diseases remain serious health problems and are caused, among others, by drugs, chemicals, and alcohol. Although liver disease(s) is stereotypically linked to alcohol or drugs or chemicals, the truth is that there are over 100 known forms of liver disease caused by a variety of factors and affecting everyone from infants to older adults worldwide [49,50-52]. Chronic liver disease is a major cause of morbidity and mortality throughout the world. Conventional medical therapy for many common liver disorders, including non-alcoholic fatty liver disease and viral hepatitis, has limited efficacy and potentially life-threatening side effects. This has increased dependence on complementary and alternative medicine (CAM), especially herbal therapy. Various medicinal plants and their formulations and dosage are used in traditional medicine for their effects on liver disorder and a number of herbal preparations are

available on the market traditionally since ancient time [52,53,55].

The diagnosis of major liver problems can be made by herbalists or medical practitioners from prior medical history of patient and physical examination and laboratory tests among other. As [52,53] stated comparative studies on medicinal plants in different cultures or ethnic groups of a country or among different countries may contribute to the identification of the most usable species of plant for treating different ailments, typically liver cases and its related disorders. A considerable amount of research has been conducted worldwide and in Ethiopia too on medicinal plants and ethnobotany with an emphasis on field surveys and documentation of people's indigenous knowledge on the usage of traditional medicinal plants that used for curing of liver complication. Hence, we found it worthwhile to go through ethnobotanical information on traditional medicinal plants used for treatment of human liver problem in Ethiopia from different ethnic groups and perspective and compile existing information as a basis lead for further investigations into these plants. Hence, the main objective of this review was ethnobotanical study on the medicinal plants with emphasis on the compilation and documentation of indigenous knowledge associated with the treatment of liver problems by these plants reported from different ethnic groups in Ethiopia.

2. METHODS

The present reviewed literatures on the major herbal medicine that has contributed the most in the protection of liver diseases/problems in Ethiopia was assessed from different sources. Relevant information were searched and studied on traditional medicinal plants pertaining to the treatment of liver problems in human. On this regard, a systematic analysis and review of research literature associated to medicinal plants used for traditional medication of liver diseases practiced by indigenous people which were conducted from different ethnic groups from the

past to this present time. These profoundly help to indigenous practice aiding in medicinal plant selection for treatment of liver problems along the ethnicity based on the availability of plants in the districts or locality.

2.1 Approaches of Article Selection

A comprehensive research literature search strategy was carried out from webs (Google search): Ethno-botanical/ethno-medicinal studies reporting on medicinal plants used for traditional liver problem treatment in Ethiopia were congregated by the following concrete search approaches: Unpublished MSc thesis or/ and PhD dissertation research reports using Google search engine and local university websites; Published articles of journal using international scientific databases including PubMed, Science direct, Researchgate, Web of Science, Google scholar, AJOL, Hinari, etc.

All possible literature search was made to address the expected objectives using the following important initial key terms: Ethiopian medicinal plants, anti-liver problems plants, Traditional knowledge medicinal plants, Medicinal Plants/Medicinal herbs, Indigenous knowledge, Plants/Herbal/Medicine/Remedies, Folk Medicine/Folk remedies/ Home remedies/ Herbal remedies, Ethnobotany/Ethnobotanical, Ethnopharmacology/ Ethnopharmacological, Ethnomedicine/ Ethnomedicinal, Ethnopharmaceutical, Medico-cultural.

2.2 Inclusion and Exclusion Criteria

The author selected the articles based on the relevancy and clearness of the data related to the traditional medication. The reliable and related articles/theses/ dissertation were downloaded from the sources and critically inspected for inclusion in the review. The comprehensive literature or article screening was made on the following inclusion and exclusion criteria.

2.2.1 Inclusion criteria

Published and unpublished ethno-botanical and ethnomedicinal surveys reporting on anti-liver problems medicinal plant(s), conducted at any time period in Ethiopia. All liver related problems were included from original literature reported as liver disease, liver disorder, hepatitis, jaundice, etc. The inclusion was restricted to original research articles published in English language studied in all parts of the Ethiopia. If the

information is not clearly stated or missed for analyzing, then rephrasing/correction was made, particularly local name and habit of the plants, and misspelled scientific names were retrieved from Natural Database for Africa (NDA), Version 2.0 and Google searching.

2.2.2 Exclusion criteria

The following types of research data were excluded by looking the tangible data that used for analysis: Published and unpublished ethno-botanical and ethno-medicinal surveys lacking information on anyone of the following: study areas/localities, scientific plant names, informant's involvement and not reporting information about anti- liver problems medicinal plants; Non-open access journal articles or partially accessed (abstract only) articles; Review articles, historical documents or experimental studies or article from other countries.

2.3 Data Analysis

The collected Ethnobotanical data reported medicinal plants used for treatment of liver problems and associated indigenous knowledge was entered in to Excel spreadsheet and summarized using descriptive statistics. The spreadsheet data filter facility was employed to determine frequencies and percentage of citations so as to identify the most common use, parts used and route of administration and habit of medicinal plants preferred for treatment of liver problem throughout the country. The results were presented using pie charts, bar chart and tables.

3. RESULTS AND DISCUSSION

3.1 Overview of Ethnobotanical Studies on Medicinal Plants

Ethno-medicinal and ethnobotany studies on plants demand standard procedures for botanical identification and reliable documentation of indigenous knowledge pertaining to plant distribution, management and traditional medicinal use in Ethiopia. By searching different webs (Google search), a total of 46 original ethno-medicinal studies representing from eight different regions in Ethiopia were included in this review. Both published and unpublished (M.Sc. and Ph.D. thesis) research reports were reviewed. Study quality inconsistencies were noted with regard to sampling and number of knowledgeable informants, as well as completeness of herbal remedy recipe, prescription and dosage and antidote information

reported. Current literature survey reflects potentially important information gaps and need for standardization of ethno-medicinal studies on indigenous medicinal plants in Ethiopia. This regional distribution of anti-liver problem medicinal plants indicated that prevalence of the disease in the locality and practiced for treatment using their indigenous knowledge. From these regions of Ethiopia, majority of the medicinal plants species used for treatment of liver problems reported from Oromia region accounts (36.59%) and followed by Amhara (25%), SNNPR (18.90%), Tigray (6.71%) and least in Dire Dawa (1.22%) (Fig. 1 and Table 2). The remaining three regions of Ethiopia: i.e. Gambella, Somali, Benshangul Gumuz and Addis Ababa could not report compiled information on medicinal plants used for curing of liver diseases traditionally.

On this review, medicinal plants used for treatment of liver problems in different parts of the country reported by different authors selected for analysis with respect to number of citation and geographical distribution in the regional state of Ethiopia. On this regard, the most commonly cited medicinal plant species by different authors were *Justicia schimperiana* 9 (12.5%) [1,2,11, 16,29,31,34,43,45] followed by *Phytolacca dodecandra* 8 (11.11%) [11, 22, 23, 32, 34, 35, 37,44], *Croton macrostachyus* 7 (9.72%) [1,3,13, 18, 32, 34, 46], *Rumex abyssinicus* [13,18,29, 46] and *Cucumis ficifolius* [1,21,34,41] each represented by 4 (5.56%) and it indicated that these medicinal plants are the most preferable one for treatment of liver problems by the traditional healers in different locality (Table 2). The biodiversity of Ethiopian flora offers great possibilities in the search for natural medicinal plants that used for treatment of various liver disorders. This diversity of plants helps for preparing medicine to treat liver and its related problems. There are various reports of plants being used in treatment of jaundice and hepatitis worldwide. The use of medicinal plants selected as a source of drugs to cure various diseases in local community is as old as humankind itself. Even to the present day, medicinal plants are available as cheap and accessible source of drug for most of developing and also in developed countries [48].

3.2 Medicinal Plants and Associated Indigenous Knowledge

In this reviewing, a total of 114 medicinal plants used for treatment of liver problems were

reported from 46 articles (40 published and 6 unpublished journals) by different authors conducted from different ethnic groups of Ethiopia. These information were recorded primarily from the informants the study area by researchers as being used for treatment of liver problems in the country. These plant species are belongs to 50 families and 90 genera. From the particular mentioned above, family Euphorbiaceae was represented by 13 (11.4%) species followed by Asteraceae which was represented by 12 (10.53%) species, Fabaceae was represented by 10 (8.77%) species and Lamiaceae 7 (6.14%) species, Amaranthaceae 5(4.39%) species, Cucurbitaceae and Moraceae represented by 4 species each, 4 families represented by 3 species each, 8 families represented by 2 species each and 31 families represented 1 species each (Fig. 2). The detail of all identified medicinal plants with their respective family, vernacular name, habitat and habit was presented in (Table 2).

This review of literature is a good indicator for the presence of a considerable diversity of plant species in Ethiopia. The existence and utilization of such a large number of medicinal plants by indigenous people in the specified area indicates that majority of the people used and continue to use indigenous medicinal plant practices to cater medication liver problems. The herbalists collected the medicinal plant from natural environment, forest, farm land and even there was a relaxed practice to cultivation medicinal plants in the areas. There were various reports of plants being used in treatment of liver problems in Tribal Communities of Paschim Medinipur District, West Bengal, India *Mentha spicata* dry leaves powdered and eaten with chilli and *Achryanthes aspera* root was crushed to powder and boiled in water [48]. In similar report, leaves of *Ricinus Communis* in India community used for treatment of liver diseases [50].

3.3 Habits of the Medicinal Plants

The assessment on the habits of the medicinal plants depicted that herbs constituted the highest fraction 48 species (42.11%) followed by shrubs 33 species (28.95%), trees 23 species (20.18%) and climbers 10 species (8.77%) (Fig. 3). Disagreement with research reported by (6) indicated mostly shrub was frequently used for remedy preparation for different human ailments treatment. This could be due to the fact that naturally there are more herbs than woody plants species. Herbs are overused because of their

diverse value for humans. It might also indicate that the threats that exist on other growth form particularly shrubs and trees. This might show that there was abundance of herbs because the area was rich with average rain fall for most of the regions of Ethiopia. This made conducive for growth of herbs and collected easily. Therefore, the trend of using more of herbaceous plants could be advantageous as it is easier to cultivate them when they are in short supply and they are annual. Relatively high number of herbs and shrubs for medicinal purpose treating different human ailments were also previously reported in Ethiopia [13].

3.4 Plant Parts Used for the Treatment of Liver Problems

With regard to the plant parts used for medicinal purposes, different parts of the plants were reported to be used for medicines. The most frequently utilized plant parts was root (24.78%) followed by leaf (23.89 %), combination parts (two parts combined) (21.24%), whole part (8.85%) and bark (5.31%) whereas flower, fiber, stem and Aerial apex of plant parts were the least used (0.88 %) each (Fig. 4). This review is in agreement with the work of [2,6,35] which reported roots as one of the most extensively




Fig. 1. Distribution of medicinal plants used for treatment of liver problem in Ethiopia




Fig. 2. The top nineteen representative families of medicinal plants used for treatment of liver problems reported in Ethiopia




Fig. 3. Habit of medicinal plants used for treatment of liver problem in Ethiopia




Fig. 4. Parts of medicinal plants used for treatment of liver problem in Ethiopia




Fig. 5. Route of administration of medicinal plants used for treatment of liver problem in Ethiopia

used part of the plant in remedy preparations Ethiopia. However it disagreed with the report of used for treatment of multiple diseases in [1,4,5] which reported the leaf as mostly

preferred for remedy preparations. The report also shows the flower parts as been highly utilized for treatment of jaundice and Hepatitis among Indian people [54].

Table 1. Mode of preparation reported traditional medicinal plants for treatment of liver problem

Mode of preparation	Frequency (F)	Percentage (%)
Crushing	23	20.18
Squeezing	3	2.63
Decoction	5	4.39
Concoction	1	0.88
Steaming	2	1.75
Smoking	1	0.88
Cooking	2	1.75
Chewing	2	1.75
Fumigating	1	0.88
Powdering	8	7.02
Chopping	1	0.878
Grinding	7	6.14
Pounding	3	2.63
Creaming	1	0.88
Boiling with Water	5	4.39
Washing	2	1.75
Baking	1	0.88
Heat	1	0.88
Four or more methods	9	7.89
Combination of three methods	15	13.16
Combination of two methods	21	18.42

3.5 Preparation, Dosage and Route of Administration of Medicinal Plants

Plant parts were prepared as medicine using fresh and dried plant materials. The dependency of the inhabitants on fresh materials in the study area including the removal of fresh leaves and roots put the plants under serious threat than the dried form, as fresh materials are harvested directly and used soon with its extra deterioration with no chance of preservation i.e. not stored for later use. However, during this review local healers were argued that fresh materials are effective in treatment as the contents are not lost before use compared to the dried forms. This review revealed that different forms of preparations were investigated. These include: homogenizing in water, crushing, decoction, squeezed, chewing, smoking, extract with cold

water, chopping and concoction. Medicinal plant parts were reported mostly to be crushing (20.18%), combination of two methods (18.42%), combination of three methods (13.16%), powdering (7.02%), grinding (6.14%), boiling with water (4.39%), decoction (4.39%), squeezing and pounding during preparation of remedies. Research conducted on medicinal plants used for the treatment of jaundice and hepatitis based on socio-economic documentation in India by [54] decoction was the major method for preparation of remedy practiced by indigenous healers.

A single herbal preparation was taken by mixing with different ingredients recommended antidotes includes drinking milk, eating honey, bread of black *tef* and occasionally drinking water slowly accordingly. In similar manner different preparations and application methods of medicinal plants were mentioned for internal and external use in which water was mostly used to dilute plant preparations while some remedies were prepared from dry and fresh plant parts.

This review of literature indicated that herbal remedies were prepared by healers and used medicinal plants for treatment of liver problem in the form of fresh 84 species (73.68 %) while 17 species (14.91%) were prepared and used either fresh or dry form of condition (Fig. 6).

Regarding the route of application medicinal plants, the eminent method was mainly 99 (88.39%) through oral followed by the combination of dermal and oral 4 (3.57%), Dermal 3 (2.68%), and dermal and nasal 2 (1.79%), smoking (0.89%) account the least number (Fig. 5). After preparing the remedy properly, the required dosage of medicine taken mostly through oral because the healers assumed that it assuring the effectiveness for curing human diseases [53]. According to healers, preparations were prescribed to patients differently for different age groups. The dosage prescription for children was mostly lower than for adults. Dosages were estimated using lids, spoons, cups, glasses, pinches or handfuls. The amounts of remedy and prescription rates were generally dependent on the degree and duration of the degree and severity of the disease. Treatment durations varied between a minute to some days [1-39]. The diagnosis of major liver problems can be made by herbalists or medical practitioners from prior medical history of patient and physical examination and laboratory tests among other.

Table 2. List of medicinal plants used for treatment of liver problem in Ethiopia

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Abrus precatorius</i> L.	Fabaceae	Cl	Badaluwa(Dw)	S	Crushed and concocted	Fresh	Oral	SNNPR	[39]	1
<i>Acacia abyssinica</i> Benth.	Fabaceae	T	Girar (Am)	L	Crushed	Fresh	Oral	Amhara	[24]	1
<i>Acacia seyal</i> Del.	Fabaceae	T		L	Crushed	Fresh	Oral	Afar	(33)	1
<i>Acacia tortilis</i> (Forssk.) Hayne	Fabaceae	T	Assel	R	Crushed and Decoction	Fresh	Oral	Dire Dawa	[28]	1
<i>Acalypha indica</i> L.	Euphorbiaceae	H	-----	L	Decoction and extract	Fresh	Oral	Afar	[33]	1
<i>Acalypha villicaulis</i> A. Rich.	Euphorbiaceae	H	Wak'ak'uwa (Dw)	R	Crushed	Fresh	Oral	SNNPR	[39]	1
<i>Achyranthes aspera</i> L.	Amaranthaceae	H	Telenj (Am)	L	Ground and powdered	Fresh	Oral	Amhara	[32]	1
<i>Acmella caulirhiza</i> Del.	Asteraceae	H		WP	Concocted, crushed, powdered	Fresh	Oral	Oromia	[45]	1
<i>Acokanthera schimperi</i> (A. DC.) Schweinf.	Apocynaceae	T	Merz/Mirez(Am)	AA	Crushed and fumigate	Fresh/dry		Amhara	[11]	1
<i>Adhatoda schimperiana</i> Hochst.ex.	Acanthaceae	Sh	Simiza (Am)	L	Boiled; crushed	Fresh	Oral	Amhara	[37, 38]	2
<i>Afrocarpus falcatus</i> (Thunb.) C.N.Page	Podocarpaceae	T	Dagucho	L	Crushed	Fresh	Oral	SNNPR	[15]	1
<i>Ajuga integrifolia</i> Ham.Buch.	Lamiaceae	H	Armagusa (Oro)	WP	Heat	Fresh	Oral	Oromia	[25]	1
<i>Aloe gilbertii</i> Sebsebe & Brandham	Aloaceae	H	Hargessa(Oro)	La	Crush and extract	Fresh	Oral	Oromia	[6]	1
<i>Aloe macrocarpa</i> Tod.	Aloaceae	H	Hargessaa (Oro)	R	Concocted, crushed	Fresh	Oral	Oromia	[45]	1
<i>Aloe monticola</i> Reynolds	Aloaceae	H	Genenno (Had.)	La	Squeezing	Fresh	Oral	SNNPR	[31]	1
<i>Amaranthus caudatus</i> L.	Amaranthaceae	H	Bertefi (Oro)	S	Crushed and homogenizing with water, powdering	Fresh/dry	Oral	Oromia	[41]	1
<i>Amaranthus hybridus</i>	Amaranthaceae	H	-----	Fr	Bake	Fresh	Oral	Oromia	[40]	1

Spp. Linn.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Arisaema schimperianum</i> Schott	Araceae	H	Amoch (Am)	L, Fr	Crushed and Concoction; homogenizing with water	Fresh	Oral	Oromia	[41]	1
<i>Aristolochia bracteolata</i> Lam.	Aristolochiaceae	Cl	----	WP	Crushed	Fresh	Oral	Afar	[33]	1
<i>Asparagus africanus</i> Lam.	Asparagaceae	Cl	Shuko (Oro)	L,Fr	Concoction , chewing	Fresh	Oral	Oromia	[41]	1
<i>Asparagus leptocladodius</i> Chiov.	Asparagaceae	Cl	Keleme sere (Oro)	L,R	Crushed, Concoction and infusion	Fresh	Oral	Afar, Oromia & Harar	[4]	2
<i>Aspilia gillettii</i> Wild.	Asteraceae	H	----	L,R	Crushed	Fresh	Oral	Oromia	[6]	1
<i>Balanites rotundifolia</i> (van Tiegn.) Blatter	Balanitaceae	T	----	L,B	Chewed	Fresh	Oral	Afar	[33]	1
<i>Bersama abyssinica</i> Fresen.	Melanthaceae	T	Xewerako (Sd)	B	Crushed	Fresh	Oral	SNNPR	[15]	1
<i>Bidens macroptera</i> (Sch. Bip. ex Chiov.) Mesfin	Asteraceae	H	Kello (Oro)	R	Crush and Concoction and homogenizing with water	Fresh	Oral	Oromia	[41]	1
<i>Boerhaavia plumbaginea</i> Cav.	Myrtaceae	H	-----	L	Decoction	Fresh	Dermal	Ethiopia	[3]	1
<i>Boswellia microphylla</i> Chiov.	Burseraceae	T	-----	B	Crush and squeezing	Fresh	Oral	Oromia	[10]	1
<i>Brucea antidysenterica</i> J. F. Mill.	Simaroubaceae	T	Aballo (Am) Kilisa adi (Oro)	L, Fr	Boiling and steaming	Fresh	Smoke, Oral	Amhara Oromia	[35] [40]	2
<i>Cadaba glandulosa</i> Forssk.	Capparidaceae	Sh		L	Crushed		Or, Der	Afar	[33]	1
<i>Calpurnea aurea</i> (Alt.) Benth	Fabaceae	Sh	Cekkatta (Sd), Digta (Am)	S, L	Crushed; boil and inhale	Fresh	Oral, Smok	SNNPR. Amhara	[26] [37]	2
<i>Carica papaya</i> L.	Caricaceae	T	Papaya	S,L,R	Crushed; pouring and squeezing	Fresh/Dry	Oral	Oromia Tigray	[41, 43] [14]	3
<i>Catha edulis</i>	Celastraceae	Sh	Jimaa (Dw)	L	Decoction	Fresh	Oral	SNNPR	[39]	1
<i>Celosia polystachia</i>	Amaranthaceae	H	----	L	Crushed	Fresh	Oral	Afar	[33]	1

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
(Forssk.) C.C. Towns.										
<i>Cineraria deltoidea</i> Sond.	Asteraceae	H	---	WP	Crushed and concoction, homogenizing with water	Fresh	Oral	Oromia	[41]	1
<i>Cirsium englerianum</i> O. Hoffm.	Asteraceae	H	Umbahoo(Oro)		Squeezed	Fresh	Oral	Oromia	[46]	1
<i>Cissampelos mucronata</i> A.Rich.	Menispermaceae	H	Kawuro (M)	L	Ground	Dry	Oral	SNNPR	[17]	1
<i>Citrus aurantifolia</i>	Rutaceae	T	Lomi	Fr	Ground	Dry	Oral	Tigray	[14]	1
<i>Clerodendrum myricoides</i> (Hochst.) R.Br. ex Vatke	Lamiaceae	Sh	Marasisa (Oro)	Fr	Concocted, crushed, powdered	Fresh	Oral	Oromia	[41]	1
<i>Clutia lanceolata</i> Forssk.	Euphorbiaceae	H	Fiyele feji (Am)	L, R	Ground , powdered	Dry	Oral	Amhara SNNPR	[32] [31]	2
<i>Coccinia abyssinica</i>	Cucurbitaceae	Cl	Ushushiya (Dw)	R	Crushed	Fresh/dry	Oral	SNNPR	[39]	1
<i>Combretum sp.</i>	Combretaceae	T	Fongera	R,B	Ground	Fresh/dry	Oral	Amhara	[32]	1
<i>Cordia africana</i> Lam.	Boraginaceae	T	Wanza (Am), Awhi (Tg)	L,R,B	Crushed, boiled; decoction and chewed	Fresh	Oral	Amhara Tigray	[19, 32] [1]	3
<i>Crepis ruepellii</i> Sch. Bip.	Asteraceae	H	Kartasa (Oro)	R	Concoction, Crushed, Powdered	Fresh/dry	Oral	Oromia	[41]	
<i>Croton macrostachyus</i> Del.	Euphorbiaceae	T	Makanisaa (Oro), Missana (Am), Tambok (Tg)	Wp	cooked; boiled and pounded; decoction; paste; boiling; powdering	Fresh/dry	Oral	Oromia Amhara Tigray Ethiopia	[46] [18, 32,34] [1, 13] [3]	7
<i>Cucumis dipsaceus</i> Ehrenb. ex Spach	Cucurbitaceae	Cl	Hare goge (Oro)	R, Fr	Crushed, Concoction	Fresh	Oral	Afar, Oromia & Harar Harar	[4] [36]	2
<i>Cucumis ficifolius</i> A. Rich.	Curcurbitaceae	Cl	Yemidir Embuay (Am), Ramboambo (Tg), Anchote (Oro)	WP, R	Crushed; Powdered	Fresh/dry	Oral	Tigray Amhara Oromia	[1] [21, 34] [41]	4
<i>Cyathula polycephala</i> Bak.	Amaranthaceae	H	Hacho (Oro)	R	Concoction , Crushed, chewed	Fresh	Oral	Oromia	[41]	1

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Cymbopogon commutatus</i> (Steud.) Stapf	Poaceae	H		Ap	Fumigated	Fresh	Oral, Der	Afar	[33]	1
<i>Cynoglossum coeruleum</i> Hochst.	Boraginaceae	H	Qarchaba (Oro)	R	Concoction, Decoction, Crushed	Fresh	Oral	Oromia	[41]	1
<i>Dioscorea alata</i> L.	Dioscoriaceae	Cl	Boyna (Sd)	St	Cooked	Fresh	Oral	SNNPR	[26]	1
<i>Dodonaea angustifolia</i> L. f.	Sapindaceae	Sh	Etancha (Sd), Iticha (Oro)	L	Decoction, crushed, powdered	Fresh	Oral	SNNPR Oromia	[26] [29]	2
<i>Dorstenia barnimiana</i> Schwienf.	Moraceae	H	Work Bemeda (Am)	R	Powdered	Dry	Oral	Amhara Tigray Amhara	[34] [13] [35]	3
<i>Dorstenia foetida</i> (Forssk.) Schweinf.	Moraceae	H	Worq-bemeda	R	Crushed and pounded	Fresh	Oral	Amhara	[19]	1
<i>Euphorbia abyssinica</i> J. F. Gmel.	Euphorbiaceae	T	Kulkual (Am)	R, La	Crushed ; squeezing	Fresh	Oral	Amhara	[11,32, 34]	3
<i>Euphorbia dumalis</i> S. Carter	Euphorbiaceae	H	Dargu adi (Oro)	R	Crushed	Fresh	Oral	Oromia	[6]	1
<i>Euphorbia lathyris</i> L.	Euphorbiaceae	H	Ambuluk(Oro)	Fr	Concocted, crushed	Fresh	Oral	Oromia	[41]	1
<i>Euphorbia schimperiana</i> Scheele	Euphorbiaceae	Sh	Gurii (Oro)	L	Concoction, Crush, chew	Fresh	Oral	Oromia	[41]	1
<i>Euphorbia triaculeata</i> Forssk.	Euphorbiaceae	Sh		L	Crushed	fresh	Na, De	Afar	[33]	1
<i>Fagonia schweinfurthii</i>	Zygophyllaceae	Sh		St,R, WP	Crushed, boiled & fumigating	fresh	Oral, Der	Oromia Afar	[10] [33]	2
<i>Ferula communis</i> L.	Apiaceae	H	Gnida (Oro)	R	Decoction	fresh	Oral	Oromia	[41]	1
<i>Ficus carica</i> L.	Moraceae	Sh	Beles (Am)	L, St	Ground	fresh	Oral	Amhara	[32]	1
<i>Ficus sycomorus</i> L.	Moraceae	T	Odaa (Oro.)	La	Creaming	Fresh	Dermal	Oromia	[5]	1
<i>Flueggea virosa</i> Guill. & Perr.	Euphorbiaceae	Sh	Shasha (Am)	L	Powdered	Dry	Nasal/Oral	Amhara	[32]	1
<i>Galinsoga parviflora</i>	Asteraceae	T	Ematiya/bizdiya (Dw)	L,R	Crushed	Fresh	Oral	SNNPR	[39]	1
<i>Galinsoga parviflora</i> Cav.	Asteraceae	H	Midirberbere, (Am)	Fl	Chopped	Fresh	Oral	SNNPR	[17]	1

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Gnidia glauca</i>	Thymelaeaceae	Sh	Migra (Dw)	R	Crushed	Fresh	Oral	SNNPR	[39]	1
<i>Grewia villosa</i> Will.	Tiliaceae	Sh		WP	Crushed	Oral	Oral, Nasal	Afar	[33]	1
<i>Hypericum quartinianum</i> A. Rich.	Hypericaceae	Sh	Muke fonii (Oro)	L	Pounding and homogenized in water	Fresh	Oral	Oromia	[2]	1
<i>Indigofera arrecta</i> A.Rich.	Fabaceae	Sh	Wareami (A)	L	Smoked	Dry	Smoke	SNNPR	[17]	1
<i>Indigofera articulata</i> Gouan	Fabaceae	Sh	----	R	Powdered	Oral	Oral	Afar	[33]	1
<i>Indigofera spicata</i> Forsk.	Fabaceae	Sh	----	R	Powdered	Oral	Na, Der	Afar	[33]	1
<i>Jatropha curcas</i> L.	Euphorbiaceae	T	Beeroo faranji (Oro)	S	crushed, powdered	Dry	Oral	Oromia	[46]	1
<i>Justicia ladanoides</i>	Acantaceae	H	Mulu muk'uwa (Dw)	L,R	Crushing and rubbing; powdering	Fresh/dry	Derma & Ora	SNNPR	[39]	1
<i>Justicia schimperiana</i> (Hochst. ex Nees) T.Anders.	Acanthaceae	Sh	Shimieya (Tg), Dhumuga(Oro), Tumunigga(Hd), Sensel (Smiza) (Am)	L,R	Crushed; pounded; Concocted, powdered; decocted/squeezed; chopped, crushed, and boiled	Fresh/dry	Oral, Der, nasal	Tigray SNNPR Oromia Amhara	[1] [31] [2, 29, 43, 45] [11,16, 34]	9
<i>Kalanchoe petitana</i> A. Rich.	Crassulaceae	H	Endehula (Am)	L	Powdered	Dry	Oral	Amhara	[32]	1
<i>Kanahala laniflora</i> (Forssk.) R. Br.	Asclepidaceae	Sh	Wundiffo (Oro)	R	Powdered	Fresh/dry	Oral	Oromia SNNPR	[9] [8]	2
<i>Leucas stachydiformis</i> (Benth.) Hochst. Ex Briq.	Lamiaceae	H	Businae (M)	L,B	chopped and drench (soak)	Fresh		SNNPR	[17]	1
<i>Maytenus arbutifolia</i> (A. Rich.) Wilczek	Celastraceae	Sh	Qartame (Oro)	L,Fr	Concoction , crushed	Fresh	Oral	Oromia	[41]	1
<i>Mentha spicata</i> L.	Lamiaceae	H	Nana	L	Boiling or pounding	Fresh/dry	Oral	Dire Dawa	[28]	1
<i>Microglossa pyrifolia</i> (Lam.) Kuntze,	Asteraceae	Sh	Y/m- meqenet (Sk.)	L	Crushed	Fresh	Oral	SNNPR	[20]	1
<i>Nicandra physaloides</i>	Solanaceae	H	Puqaqiya (Dw) Hawwixii(Oro)	L,B	Crushed	Fresh	Oral	SNNPR Oromia	[39] [44]	2

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Nigella sativa</i> L.	Ranunculaceae	H	Tikur Azmud (Sd)	S	Ground	Fresh	Oral	SNNPR	[42]	1
<i>Oncocalyx schimperi</i> (A. Rich.) M. G. Gilbert	Loranthaceae	Sh		L	Crushed	Fresh	Oral	Afar	[33]	1
<i>Phyllanthus reticulatus</i>	Euphorbiaceae	Sh	Wusiwisiya mala (Dw)	L,R	Crushed	Fresh	Oral	SNNPR	[39]	1
<i>Phytolacca dodecandra</i> L.Heri	Phytolacaceae	Cl	Endod (Am), Handoode (Oro)	R, L	Crushed and pounded; Ground	Fresh	Oral	Oromia Amhara	[23,22, 44] [11,32, 34, 35, 37]	8
<i>Portulaca quadrifida</i> L.	Portulacaceae	H	Akalkaraha (Oro)	WP	chopped, powdered	Fresh	Oral	Oromia	[29]	1
<i>Rhamnus prinoides</i>	Rhamnaceae	Sh	Gesho (Am)	R	Ground	Fresh	Oral	Amhara	[37]	1
<i>Rhus retinorrhoea</i> Steud, ex Olive	Ancardaceae	Sh	Tilem (Am)	L,R	Powdered; Crushed	Fresh/dry	Oral	Amhara Oromia	[12] [27]	2
<i>Ricinus communis</i> L.	Euphorbiaceae	Sh	Qobboo (Oro)	L,R	Crushed	Fresh	Oral	Oromia	[22, 30]	2
<i>Rumex abyssinicus</i> Jacq.	Polygonaceae	H	Dhangago (Oro), Mekemeko (Am)	AP, R	pounded and then decocted; powdered; boiled; crushed,	Fresh/dry	Oral	Oromia Tigray Amhara	[29, 46] [13] [18]	4
<i>Rumex nepalensis</i> Spreng.	Polygonaceae	H	Germach (Bn), Shabbee (Oro)	R	Decoction, Crushed	Fresh	Oral	SNNPR Oromia	[13] [41]	2
<i>Salvia merjamie</i> Forssk.	Lamiaceae	H	Okota (Oro)	R	Concoction, Decoction, Crush	Fresh	Oral	Oromia	[41]	1
<i>Satureja punctata</i> Benth. Briq	Lamiaceae	H	Lomishet (Am)	L	Cooked	Fresh	Oral	Tigray	[13]	1
<i>Schinus molle</i> L.	Anacardiaceae	T	Tikur berbere (Tg)	L	Crushed and filter	Fresh	Oral	Tigray	[1]	1
<i>Securidaca longepedunculata</i> Fresen.	Polygalaceae	T	Xamanaayii (Oro)	B	Powdering	Dry	Oral	Oromia	[22]	1
<i>Senna petersiana</i> (Bolle) Lock	Fabaceae	Sh	Ramso (Oro)	L	Washed body	Fresh	Dermal	Oromia	[43]	1
<i>Senna alexandrina</i> Mill.	Fabaceae	Sh		B	Crushed and extract	Fresh	Oral	Afar	[33]	1
<i>Sida schimperiana</i> Hochst. ex A.Rich.	Malvaceae	S		R	Boiled and extract	Fresh	Oral	SNNPR	[7]	1

Spp.	Family	Hb	Local	PU	Mode preparation	Condition	RA	Region	References	Fr.
<i>Silene macrosolen</i> A. Rich.	Caryophyllaceae	H	Wagartii (Oro)	R	Concoction, crush, chew	Fresh	Oral	Oromia	[41]	1
<i>Solanum nigrum</i>	Solanaceae	H	Tut'naye (Sd)	L	Boiled	Fresh	Oral	SNNPR	[42]	1
<i>Solanum sp.</i>	Solanaceae	Sh	Puk'ek'iya(Dw)	R	washed/crushed		Oral	SNNPR	[39]	1
<i>Sonchus bipontini</i> Asch.	Asteraceae	H	Kartasa (Oro)	R	Concoction, Crushed or chew	Fresh	Oral	Oromia	[41]	1
<i>Stellaria sennii</i> Chiov. H	Caryophyllaceae	H		R	Decoction	Fresh	Oral	SNNPR	[8]	1
<i>Syzygium guineense</i> (Willd.) DC.	Myrtaceae	T	Duwancho (Sd)	B	Concoction	Fresh	Oral	SNNPR	[15]	1
<i>Terminalia brownie</i> Pers.	Combretaceae	T	Weyba (Am)	B	Boiled and extraction	Fresh	Oral	Amhara	[32]	1
<i>Thymus serrulatus</i> (hoechst ex. Benth)	Lamiaceae	Sh	Tosegn (Am)	L	Decoction	Fresh	Oral	Oromia	[29]	1
<i>Triumfetta heterocarpa</i> Sprague and Hutch.	Tiliaceae	Sh	Yelam tut (Am)	R	Crushed	Fresh	Oral	Amhara	[35]	1
<i>Verbena officinalis</i>	Verbenaceae	H	Atuch (Am)	R	Squeezed	Fresh	Oral	Amhara	[37]	1
<i>Vernonia myriantha</i> Hook.f.	Asteraceae	Sh	Regi (Oro)	L	Crushed , Powdered, chewed	Fresh/Dry	Oral	Oromia	[41]	1
<i>Vernonia sp.</i>	Asteraceae	H	Yesheshuwa (Dw)	R	Powdered	Dry	Oral	SNNPR	[39]	1
<i>Vitis vinifera</i> L.	Vitaceae	H	Weyne (Am)	L,R	Ground	Fresh	Oral	Amhara	[32]	1
<i>Woodfordia uniiflora</i> (A. Rich.) Koehne	Lythraceae	H	Itecha (Oro)	R	Crushed	Fresh	Oral	Oromia	[6]	1
<i>Ximenia Americana</i> L.	Olacaceae	T		Fb	Crushed and soaking	Fresh	Oral	Oromia	[10]	1
<i>Zehneria scabra</i> (Linn.f.) Sond.	Cucurbitaceae	Cl	Areg Resa (Am), Harola (Oro)	L, R	pounded and squeezed; Decoction, Crushed	Fresh	Oral	Amhara, Oromia	[21,41]	2

Hb=Habit (T=Tree, Sh=Shrub, H=Herb, Cl=Climber); PU=part use (L=Leaves, R=Roots, B=bark, Fr=Fruit, S=Seed, Fl=flower, La=latex, Fb=fiber, AP=Aerial part, AA=aerial apex, WP=Whole plant); RA=Route of administration (O=Oral, D=Dermal, N=Nasal); Language (Oro=Oromia, Am=Amharic, Dw=Dawuro, Sd= Sidama, Gd= Gedeo, Tg=Tigray, Bn=Bench, Sk=Sheko, Hd=Hadiya) , Fr.=frequency




Fig. 6. Condition of medicinal plant preparation

4. CONCLUSION

This research review revealed that there were considerable numbers of medicinal plants in Ethiopia from different ethnic group which were used to treat liver problems. Hence, medicinal plants still play significant role in the health care of system of the rural community as first choice for curing different diseases. Generally, in Ethiopia 114 medicinal plants used for treatment of liver problems were recorded from different authors (46 articles). The most common habit of medicinal plants used by different ethnicity was herbs and shrubs respectively. The most utilizable parts of medicinal plants for treating liver cases were root and leaf. Major medicinal plants were cited by different authors in different areas were *Justicia schimperiana*, *Phytolacca dodecandra* and *Croton macrostachyus*. This review indicates the prevalence of liver problems in Ethiopia which necessitates further investigations of herbs used in light of modern, clinical and scientific research approach. Most of the medicinal plants were harvested from the wild that the natural habitats need to be managed properly to minimize threats of medicinal plants in the near future.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

We would like thank you all researchers and contributors for generating these research data's compiled as a review document.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Teklay A, Abera B, Giday M. An ethnobotanical study of medicinal plants used in Kilte Awulaelo District, Tigray Region of Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2013;9:65.
2. Kefalew A, Asfaw Z, Kelbessa E. Ethnobotany of medicinal plants in Ada'a District, East Shewa Zone of Oromia Regional State, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2015;11:25. DOI:10.1186/s13002-015-0014-6.
3. Getahun A. Some common medicinal and poisonous plants used in Ethiopian Folk Medicine. Faculty of Science, Addis Abeba University, Addis Abeba, Ethiopia; 1976.
4. Belayneh A, Asfaw Z, Demissew S, Bussa NF. Medicinal plants potential and use by pastoral and agro-pastoral communities in Erer Valley of Babile Wereda, Eastern

- Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2012;8:42.
5. Abera B. Medicinal plants used in traditional medicine by Oromo people, Ghimbi District, Southwest Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2014;10:40.
 6. Lulekal E, Kelbessa E, Bekele T, Yineger H. An ethnobotanical study of medicinal plants in Mana Angetu District, southeastern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2008; 4:10.
DOI:10.1186/1746-4269-4-10.
 7. Mesfin F, Seta T, Assefa A. An ethnobotanical study of medicinal plants in Amaro Woreda, Ethiopia. *Ethnobotany Research & Applications*. 2014;12:341-354.
 8. Mesfin F, Demissew S, Teklehaymanot T. An ethnobotanical study of medicinal plants in Wonago Woreda, SNNPR, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2009;5:28.
DOI:10.1186/1746-4269-5-28.
 9. Bekele G, Reddy PR. Ethnobotanical study of medicinal plants used to treat human ailments by Guji Oromo Tribes in Abaya District, Borana, Oromia, Ethiopia. *Universal Journal of Plant Science*. 2015;3(1):1-8.
DOI:10.13189/ujps.2015.030101.
 10. Gebeyehu G, Beche D, Feyisa K. Indigenous utilization and management of useful plants in and around Awash National Park, Ethiopia. *Journal of Plant Biology & Soil Health*. 2016;3(1):12.
 11. Chekole G, Asfaw Z, Kelbessa E. Ethnobotanical study of medicinal plants in the environs of Tara-gedam and Amba remnant forests of Libo Kemkem District, northwest Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2015; 11(4):1-38.
 12. Alemayehu G, Asfaw Z, Kelbessa E: Ethnobotanical study of medicinal plants used by local communities of Minjar-Shenkora District, North Shewa Zone of Amhara Region, Ethiopia. *Journal of Medicinal Plants Studies*. 2015;3(6):01-11.
 13. Giday M, Asfaw Z, Woldu Z, Teklehaymanot T. Medicinal plant knowledge of the Bench ethnic group of Ethiopia: An ethnobotanical investigation. *Journal of Ethnobiology and Ethnomedicine*. 2009;5:34.
 14. Yirga G. Assessment of traditional medicinal plants in Endrta District, South-eastern Tigray, Northern Ethiopia. *African Journal of Plant Science*. 2010;4(7):255-260.
 15. Kewessa G, Abebe T, Demessie A. Indigenous knowledge on the use and management of medicinal trees and shrubs in Dale District, Sidama Zone, Southern Ethiopia. *Ethnobotany Research & Applications*. 2015;14:171-182.
 16. Reta H. An ethnobotanical study of useful plants of the farming site in Gozamen Wereda, East Gojjam Zone of Amhara Region, Ethiopia. Addis Ababa University, MSc. Thesis; 2010.
 17. Tolossa K, Debela E, Athanasiadou S, Tolera A, Ganga G, Houdijk J. Ethnomedicinal study of plants used for treatment of human and livestock ailments by traditional healers in South Omo, Southern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2013;9:32.
 18. Wolde-Mariam M, Limenih Y, Umer S. Ethnobotanical study on traditional medicinal plants in Dega Damot Woreda, Amhara Region, North Ethiopia. *International Journal of Research in Pharmacy and Chemistry*. 2015;5(2):258-273.
 19. Giday M, Teklehaymanot T, Animut A, Mekonnen Y. Medicinal plants of the Shinasha, Agew-awi and Amhara peoples in northwest Ethiopia. *Journal of Ethnopharmacology*. 2007;110:516–525.
 20. Giday M, Asfaw Z, Woldu Z. Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *Journal of Ethnopharmacology*. 2010;132:75–85.
 21. Meragiaw M, Asfaw Z, Argaw M. The status of ethnobotanical knowledge of medicinal plants and the impacts of resettlement in Delanta, Northwestern Wello, Northern Ethiopia. *Evidence-Based Complementary and Alternative Medicine*. 2016;24.
 22. Megersa M, Asfaw Z, Kelbessa E, Beyene A, Woldeab B. An ethnobotanical study of medicinal plants in Wayu Tuka District, East Welega Zone of Oromia Regional State, West Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2013;9:68.

23. Alito MK. Use and management of medicinal plants by indigenous people of Jima rare District in Oromia Region, Ethiopia. Haramaya University, M.Sc. Thesis Research; 2014.
24. Rangunathan M, Abay SM. Ethnomedicinal survey of folk drugs used in bahirdar zuria district in northwestern Ethiopia. *Indian Journal of Traditional Knowledge*. 2009;8(2):281-284.
25. Parvez N, Yadav S. Ethnopharmacology of single herbal preparation of medicinal plants in Asendabo district, Jimma, Ethiopia. *Indian Journal of Traditional Knowledge*. 2010;9(4):724-729.
26. Regassa R. Assessment of indigenous knowledge of medicinal plant practice and mode of service delivery in Hawassa city, southern Ethiopia. *Journal of Medicinal Plants Research*. 2013;7(9):517-535.
27. Getaneh S, Girma Z. An ethnobotanical study of medicinal plants in Debre libanos wereda, central Ethiopia. *African Journal of Plant Science*. 2014;8(7):366-379.
28. Ayalew S, Kebede A, Mesfin A, Mulualem G. Ethnobotanical study of medicinal plants used by agro pastoralist Somali people for the management of human ailments in Jeldesa Cluster, Dire Dawa Administration, Eastern Ethiopia. *Journal of Medicinal Plants Research*. 2017;11(9): 171-187.
29. Suleman S, Alemu T. A survey on utilization of ethnomedicinal plants in Nekemte Town, East Wellega (Oromia), Ethiopia. *Journal of Herbs, Spices & Medicinal Plants*. 2012;18:34–57.
30. Birhanu T, Abera D, Ejeta E. Ethnobotanical study of medicinal plants in selected Horro Gudurru Woredas, Western Ethiopia. *Journal of Biology, Agriculture and Healthcare*. 2015;5(1):83-93.
31. Temam T, Dillo A. Ethnobotanical study of medicinal plants of Mirab-Badwacho district, Ethiopia. *Journal of BioScience and Biotechnology*. 2016;5(2):151-158.
32. Mekuanent T, Zebene A, Solomon Z. Ethnobotanical Study of Medicinal Plants in Chilga District, Northwestern Ethiopia. *Journal of Natural Remedies*. 2015;15(2): 88-112.
DOI:10.18311/jnr/2015/476.
33. Teklehaymanot T. An ethnobotanical survey of medicinal and edible plants of Yalo Woreda in Afar regional state, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2017;13:40.
DOI:10.1186/s13002-017-0166-7.
34. Teklehaymanot T, Giday M. Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2007;3:12.
DOI:10.1186/1746-4269-3-12.
35. Teklehaymanot T. Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. *Journal of Ethnopharmacology*. 2009;124:69–78.
36. Fenetahun Y, Eshetu G, Worku A, Abdella T. A survey on medicinal plants used by traditional healers in Harari regional State, East Ethiopia. *Journal of Medicinal Plants Studies*. 2017;5(1):85-90.
37. Getnet Z, Chandrodyam S, Masresha G. Studies on traditional medicinal plants in Ambagiorgis area of Wogera District, Amhara Regional State, Ethiopia. *International Journal of Pure and Applied Bioscience*. 2016;4(2):38-45.
38. Birhanu Z, Endale A, Shewamene Z. An ethnomedicinal investigation of plants used by traditional healers of Gondar town, North-Western Ethiopia. *Journal of Medicinal Plants Studies*. 2015;3(2):36-43.
39. Agize M, Andarge E, Shonga A, Tora A. Utilization and conservation of medicinal plants and their associated Indigenous Knowledge (IK) in Dawuro Zone: An ethnobotanical approach. *International Journal of Medicinal Plant Research*. 2015;4(3):330-337.
40. Yirga G, Zerabruk S Traditional knowledge of medicinal plants in Gindeberet district, Western Ethiopia. *South African Journal of Botany*. 2012;78:165–169.
41. Yineger H. A study on the ethnobotany of medicinal plants and floristic composition of the dry Afromontane Forest at Bale Mountains National Park, Ethiopia. Addis Ababa University, M.Sc Thesis; 2005.
42. Busse H, Tefera G. Handbook of Sidama Traditional Medicinal Plants. University of Wisconsin-Madison; 2013.
43. Kumbi ET. Use and conservation of traditional medicinal plants by indigenous people in Gimbi Woreda, Western Wellega, Ethiopia. Msc Thesis; 2007.
44. Amenu E. Use and management of medicinal plants by indigenous people of

- Ejaji Area (Chelya Woreda) West Shoa, Ethiopia: An Ethnobotanical Approach. MSc Thesis; 2007.
45. Yineger H, Kelbessa E, Bekele T, Lulekal E: Plants used in traditional management of human ailments at Bale Mountains National Park, Southeastern Ethiopia. *Journal of Medicinal Plants Research*. 2008;2(6):132-153.
 46. Etana B. Ethnobotanical study of traditional medicinal plants of Goma Wereda, Jima Zone of Oromia Region, Ethiopia. MSc Thesis; 2010.
 47. World Health Organization: WHO monographs on selected medicinal plants. Geneva, WHO Publications; 2007.
 48. Sarkhel S. Ethnomedicinal uses of some plants in treatment of jaundice by tribal communities of Paschim Medinipur District, West Bengal, India. *Medicinal & Aromatic Plants*. 2015;4:205. DOI:10.4172/2167-0412.1000205.
 49. Rajaratnam M, Prystupa A, Lachowska-Kotowska P, Zaluska W, Filip R. Herbal medicine for treatment and prevention of liver diseases. *Journal of Pre-Clinical and Clinical Research*. 2014;8(2):55–60. DOI:10.5604/18982395.1135650.
 50. Vishal R. Protective role of Indian medicinal plants against liver damage. *The Journal of Phytopharmacology*. 2013;2(3):1-3.
 51. Féher J, Lengyel G. Silymarin in the prevention and treatment of liver diseases and primary liver cancer. *Curr Pharm Biotechnol*. 2012;13(1):210–217.
 52. Umer S, Asres K, Veeresham C. Hepatoprotective activities of two Ethiopian medicinal plants. *Pharmaceutical Biology*. 2010;48(4):461–468.
 53. Mukazayire MJ, Minani V, Christopher K, Ruffo E, Bizuru C, Stévigny P. Duez. Traditional phytotherapy remedies used in Southern Rwanda for the treatment of liver diseases. *Journal of Ethnopharmacology*. 2011;138:415–431.
 54. Abbasi AM, Khan MA, Ahmad M, Zafar M, Khan H, Muhammad N, Sultana S. Medicinal plants used for the treatment of jaundice and hepatitis based on socio-economic documentation. *African Journal of Biotechnology*. 2009;8(8):1643-1650.
 55. Anand K, Lal UR. Hepatitis and medicinal plants: An overview. *Journal of Pharmacognosy and Phytochemistry*. 2016;5(6):408-415.
 56. Dhiman RK, Chawla YK. Herbal medicines for liver diseases: Review article. *Digestive Diseases and Sciences*. 2005;50(10): 1807–1812.

© 2018 Garedew and Bizuayehu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/27534>