



Assessment of Production Constraints of Large Cardamom in the Eastern Hills of Nepal

Keshav Prasad Shrestha^{1*}, K. C. Gopal², Raju Chaudhary³, Amar Bahadur Pun⁴,
Jiban Shrestha⁵, Surendra Yadav⁶ and Dhanik Lal Mandal⁴

¹Socioeconomic Agriculture Research and Policy Division (SARPOD), NARC, Nepal.

²Institute of Agriculture and Animal Science, Tribhuvan University, Nepal.

³Commercial Crop Research Division, Khumaltar, Lalitpur, Nepal.

⁴Agricultural Research Station, Pakhribas, Dhankuta, Nepal.

⁵National Commercial Agriculture Research Program, Pakhribas, Dhankuta, Nepal.

⁶Agricultural Research Station, Jaubari, Ilam, Nepal.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAHR/2018/45905

Editor(s):

(1) Dr. Ahmed Medhat Mohamed Al-Naggar, Professor of Plant Breeding, Department of Agronomy, Faculty of Agriculture, Cairo University, Egypt.

Reviewers:

(1) Martín Maria Silva Rossi, Argentina.

(2) Ferenc Bagi, University of Novi Sad, Serbia.

Complete Peer review History: <http://prh.sdiarticle3.com/review-history/27736>

Original Research Article

Received 19 September 2018

Accepted 04 December 2018

Published 14 December 2018

ABSTRACT

Large Cardamom, one of the high value spice crop has gained renewed interest among farmers since last decade in Nepal. Higher net return and suitability in wider range of mid hills in Nepal is a major factor attracting this crop for the replacement of other crops. Despite profitability, there are major challenges in production due to biotic factors. Serious yield losses in this crop has reported during past 10 years and as this is perennial, farmers are disappointed with the current production and profitability for long run. In this study, an attempt has been made to explore potential causes of the decline in in eastern hills of Nepal. For this, household survey was conducted in five districts of eastern Nepal during 2016. Lack of disease resistant/tolerant varietal option and inadequate management practices are reported to be the major problems for promoting disease spread in the eastern region. The public sectors have been unable to fulfil the high demand of new saplings.

*Corresponding author: Email: kpshresthasocio@gmail.com;

Rhizome rot remains the most prevalent disease in studied districts followed by wilting. The study suggest plan of actions to implement for the good orchard management to address the problem of biotic factors in short run; technology development and adoption to mitigate biotic problems in large cardamom in long run.

Keywords: *Chhirkey; compound annual growth rate; Foorkey; Rhizome rot; large cardamom decline.*

1. INTRODUCTION

Large Cardamom (*Amomum subulatum* Roxb.) is a high potential crop of Nepal for the export markets in the world. The annual export value of Large Cardamom (LC) was estimated at NPR 4.85 billion in 2017 [1]. Particularly, it has been the major farm income of majority farmers in the eastern hill region of Nepal. Moreover, Nepal is the largest producer of LC in the world, supplying over 50% of the world's market demand [2,3]. Currently, it is grown in 17,002 ha, producing of 6,521 t, with 522 kg ha⁻¹ productivity [4,5]. Currently, LC is grown in 51 hill districts [6]. The LC has been known to be originated in Nepal as many wild races are reported in the eastern Himalayan region of Nepal [7]. The area under LC plantation is growing slowly and steadily. It is a low volume, high value crop with medicinal properties. It was introduced in Nepal from Sikkim long time back [8].

However, in the recent years, its' production has been severely affected by the cardamom decline. There are multiple factors associated with the decline; however the major factors are comprised of Rhizome/clump rot; and viral disease like Foorkey and Chhirkey. This problem has become widespread across the LC growing districts. As a result, the problem has received national priority that carries important for the assessment of associated factors of this problem in the respective districts.

In this context, the study was conducted to identify the status of different factors associated with the Large Cardamom decline in the major growing districts so as to recommend the control measures for the management of this problem.

2. MATERIALS AND METHODS

The study was conducted in Taplejung, Panchthar, Ilam, Dhankuta, and Tehrathum during January to July, 2016. The following approaches were adapted to conduct the study:

- i. Focus Group Discussions (FGD),

- ii. Key Informants Interviews,
- iii. Stakeholders meeting,
- iv. Literature review, and
- v. Household survey.

Literatures published by the public institutions: District Agriculture Development Offices (DADOs), Cardamom Development Office (CDC), Nepal Agriculture Research Council (NARC) and other related stakeholders were reviewed to get the secondary information on various aspects of LC decline. Household survey was carried out with an objectively designed household survey questionnaire. Altogether 30 households from the representing VDC of each five districts: were selected as the sample size and sampling for the household survey. Stakeholder meeting was organized among the stakeholders of LC to identify the associated factors of LC decline in Ilam during 2016 The information was edited by following the procedure of random checking, triangulation, and thorough discussion with the key informants. The data was analyzed with Statistical Package for Social Science (SPSS) and excel, while the qualitative information was analyzed and presented in paragraphs, charts and diagrams. Information on perception, behavior, and attitude and other variables for above indicators was processed by Content Analysis. Such information was systematically organized and presented in a logical sequence and smooth flow so as to answer to indicators already defined. The preference data for disease were categorized into rank based on the percentage of ranking for all the diseases and the disease ranking coefficient value for the districts was calculated with the help of following formula:

$$Rd_1 = \frac{Dr_1 + Dr_2 + Dr_3 \dots \dots \dots Dr_n}{N}$$

Where,

Rd₁ = disease ranking coefficient value of district d1,

N = total number of disease taken into consideration,

Dr = rank occupied by the district d1 for disease r₁, r₂, r₃..... r_n

This rank coefficient value were delineated into disease intensity and categorized into three groups: high, moderate and low disease intensity. Lower the value of rank coefficient, higher the disease intensity.

Farmers' perception was ranked based on likert scale based on 1-5 point scale for disease based on symptom asked with respondents with formula:

$$I = \sum \frac{s_i f_i}{N}$$

Where,

- I = Index of importance
- S_i = scale value
- F_i = frequency of perception on satisfaction given
- N = Number of observation

3. RESULTS AND DISCUSSION

3.1 Overview of Large Cardamom Decline

In Nepal, currently the LC is cultivated in 51 districts under 17,002 ha area producing of 6,526 t in 2016/17 with productivity 522 kg ha⁻¹ [4]. The LC decline has become havoc across the country. There are multiple factors associated with the LC decline. However, particularly two viral diseases: Chhirkey, Foorkey, and Rhizome rot have been identified as the root causal factors of LC decline in Nepal. Besides these, declining soil fertility and moisture, poor orchard management, invasion of rhizome rot disease and various insects and low manuring and fertilization are among the other major factors of the decline.

The current production of LC is reported to be two folds higher than that of 1994s. However, the productivity has been declining. The major factors of the productivity shrinkage as reported in the various areas are the invasion of Chhirkey and Foorkey, rhizome rot and blight, and caterpillar.

3.1.1 Area of Large Cardamom in study districts

The area of LC in the study districts is decreasing over 10 year's period in almost all

districts. In total, the area in 2005/06 was 8646 ha, which decreases each year thereafter and reach to 6786 ha in 2013/14. However, it again slightly increased in the last year and reach to 7645 ha in 2014/15. It is due to increase in awareness in the farmers and government intervention (Fig. 1).

While calculating Compound Annual Growth Rate (CAGR) of area with regression analysis on total area of the study districts from 2005/06 to 2014/15, the CAGR significantly decreased with (minus) -1.743 percent annually with p value 0.0003. It also depict that the area of LC in the study districts are decreasing significantly.

3.1.2 Production of Large Cardamom in study districts

Production trend of LC in the study districts is also decreasing from year 2005/06 to 2014/15. In total, the production was 5262 tons in year 2005/06 which decreases each year and reach down to 2956 tons in year 2014/15. The main reason of decline is the very old plantation orchard in the district affected by diseases like foorkey, chhirkey and rhizome/clump rot which causes destroying of several plantation orchards in the districts (Fig. 2).

We also calculated CAGR with regression analysis of total production of LC of the study districts from 2005/06 to 2014/15, the CAGR of production also significantly decreasing with (minus) -0.284 percent annually with p value 0.0002. It also reveals that the area of LC in the study districts is also decreasing significantly.

3.2 Assessment of Farmers Practices for the management of LC decline

3.2.1 Cultivars and Propagation

There are many cultivars with varied yield potentials and morphological characteristics which are specific to different climates and altitudes. The popular cultivars commonly adopted in the region are Ramsay, Golsay, Ramla, Chibesay, Sawane, Varlange, Damarsay, Jirmale, Salakpure, and Pakhe [9]. The study found that cultivars: Golsay and Chibesay had lower productivity compared to Ramsay in Panchthar, while it was found similar among these cultivars in Taplejung and Terhrathum.

Most of the farmers (96.32%) prefer rhizome propagation (Fig. 3). Among them, 70% farmers buy suckers from neighbors and 30% farmers use own orchards suckers for multiplication. In Tehrathum, all the farmers use suckers for




Figure 1. Trend of area (ha) of LC in the study districts




Figure 2. Production (tons) trend of LC in the study districts




Figure 3. Method of propagation of LC adopted by respondent in percentage
(Source: Field Survey, 2016)

Overall, farmer buy suckers for propagation from other's orchard (58.82%) and some transfer suckers from own field (41.18%) (Fig. 4). This may be due to ease of sucker transplantation and lack of knowledge about healthy plant establishment. Less than 1% farmers adopt other means of propagation such as tissue culture nursery plants.

Every year 20,000 saplings were supplied through Haspokhari LC Production Cooperative, Kolbung, Ilam. CDC, Ilam produces around 4.7 lakh saplings annually [10]. Similarly, NARC, Dhankuta also produces 50 thousand seedlings through seed and tissue culture but this is not sufficient to fulfill the high demand of farmers.

LC seedling demand is very high throughout Nepal. The farmers from Province 1, whose orchards have been collapsed for last few years are willing to replant with healthy seedlings. Due to high suckering nature of some cultivars; and high demand and price, farmers from Kolbung, Ilam are more interested towards sapling production than capsules.

The study revealed that twenty seven nurseries are in Ilam, mostly concentrated in Jirmale, Kolbung, Fikkal and Mabu. There is nearly 10 lakh seedlings demand per annum, but the supply is only 1.5 lakh. Likewise seedling production internally in Taplejung is also insufficient.




Figure 4. Source of clone for propagation adopted by respondents in percentage
(Source: Field Survey, 2016)

3.2.2 Manure and fertilizer management

The study revealed that almost all LC growers do not apply manure and fertilizer. The general perception is that natural source of plant nutrient derived from the decaying leaf litter of the shading trees is sufficient for the crop. The LC orchards without additional manuring and fertilizing look poor growth and yield. The soil fertility in most of the areas has been declining over time. The possible reason for declining the soil fertility is because of exhausting the plant nutrients from the continuous cultivation for long time.

3.2.3 Irrigation

Ninety percent of respondents were irrigating orchards by different techniques such as canal/surface irrigation (39.9%), sprinkler (15%) and both canal and sprinkler (32.7%). The canal irrigation system could be the potential source of transmission of various fungal diseases as these crops grown mainly in terrain. The knowledge of disease transmission process through water found to be known by 29.33% of the respondents but they did not adopt any precaution measures to check the disease spread.

3.2.4 Current status of LC decline

LC production is declining in most of the growing districts in Nepal. There are many reasons of this problem. In the study areas, rhizome/clump rot, chhirkey, and foorkey diseases are reported as the major factors of LC decline. The fruits remaining unripe due to unknown reason and declining in subsequent year has been reported as the new problem in Taplejung. The inadequate irrigation, use of disease infected seed materials, lack of knowledge about the mode of disease spread and continuous deforestation of shade trees from the LC orchards are observed as the major causal factors of the decline in the study areas. The diseases appeared most severe in older orchards in Ilam. Seed abortion and unripening has also become a threat to LC growing areas in Ilam. However, newer plantations are also poor due to the plantation of unhealthy planting materials. The plantations have almost collapsed in Ilam. However, some plantations of those farmers, who have given good care of management and sanitation, are still performing well for good yield.

Inappropriate plantation of cultivar as per the altitude and climate are also being major problem

among the farmers, for instance *Ramsay* cultivar performs better in high altitude but it was also cultivated in lower belts. Although number of diseases has been diagnosed, the remedial pace for the control of these diseases is very slow. Farmers reported that over 40% of loss is associated with diseased orchards and will be collapsed completely by next 2-3 years. Comparatively lower incidence of diseases as reported in Taplejung could be the reason for higher production, but there were many reports that showed a trend of gradual increase in the diseases incidence. All these curtailed the income of growers, forcing them to look to alternate farming enterprises.

3.2.5 harvesting techniques as a source of disease spread

Use of single knives to harvest spikes of all plants in an orchard was found as main causal factor of spreading viral as well as fungal diseases in LC the plantation.

3.3 Pattern of Disease Distribution in studied districts

Distributions of diseases were mapped into three categories viz. places with low incidence (01-15%), moderate incidence (15-30%) and high incidence (>30%) based on disease incidence data as a perception of farmers and FGD with the concerned stakeholder.

As a whole, there was decline in LC production due to disease problems in Ilam, Panchthar, Tehrathum and Dhankuta districts. There was no evidence of decline in Taplejung district in spite of increasing infection of diseases. The decline was highest in Ilam because of higher incidence of chhirkey, wilt, rhizome rot, and leaf blight, whereas severity of foorkey and others were low to moderate (Table 1).

Similarly, Panchthar district reported greater decline due to higher incidence of diseases like wilt, leaf blight and medium incidence of chhirkey, foorkey and low incidence of rhizome rot. Dhankuta and Tehrathum also reported some decline due to low/moderate incidence of disease and pests. Thus, Ilam and Panchthar can be grouped in one category with higher decline due to severe incidence. Taplejung, Tehrathum and Dhankuta can be classified into second group with moderate decline with medium incidence of diseases.

Table 1. Severity of different diseases in the study districts

District	Disease severity				
	Rhizome rot	Wilting	Chhirkey	Foorkey	Immature capsule
1. Ilam	High	high	Moderate	Low	Low
2. Panchthar	High	Moderate	High	Low	Moderate
3. Tehrathum	High	Low	Low	Moderate	Moderate
4. Dhankuta	Moderate	High	Low	Low	Low
5. Taplejung	High	Low	Moderate	Low	Moderate

Source: Field survey, 2016

3.3.1 Distributions of rhizome rot disease

Rhizome rot is a fungal disease in LC caused by *Rhizoctonia solani* Kuhn. Teleomorph stadium *Thanatephorus cucumeris* Donk. It is characterized by foul smell and red pink color of rhizome subsequently wilting symptoms and death of plants. It is transmitted by fungal movement from infected plant to healthy plants. It is most damaging disease of LC wherever this is grown but the intensity is high in higher altitude (1550 m) and during heavy raining months and under deep shade [11]. This disease ranked top priority by the farmers of eastern hill districts [12,13].

This disease is prevalent in most of the LC growing region throughout Eastern hills during November to February. Dhankuta and Ilam are more prone to this disease. Other districts are less affected. Good orchard management with adequate irrigation system was reported to be very effective in controlling this disease.

3.3.2 Distribution of wilting

Wilting is not serious in case if it is due to abiotic factors and there is chance of recovery. It has ranked second most important disease in the study districts. It was also supported by the Yadav et al. [13]. Lack of appropriate drainage was reported to be major cause of wilting. This disease was seen during February and March. In Godak and Panchkanya of Ilam and Subang, Bharappa and Ranitar of Panchthar were reported to have this disease. Wilt disease caused by *Fusarium oxysporum* Schlecht shows symptoms of chlorosis of the older leaves at the junction of petiole with pseudo stem or their collapse while still green.

Wilting due to stem borer and rhizome borer are also reported in Dokhu, and Hangdeva of Taplejung. But this disease is not much serious in comparison to others.

3.3.3 Distribution of leaf blight disease

Leaf blight is complex diseases seen in LC. It has got third rank of the disease in the study district. It was also supported by Yadav et al. [13]. It mainly occurs due to bacteria and fungi (*Pestalotiopsis versicolor* (Sepp.). It is characterized by minute grey spots with chlorotic holes on the leaves mostly from the tip and margin. This disease is seen during January to May. It is often confusing with the wilting symptoms produced by fungal from wilting due to borer. Leaf blight was reported in all the districts, especially in Taplejung, Panchthar and Ilam districts. Farmers reported that leaf blight is also a major cause of decline of LC production.

3.3.4 Distribution of foorkey disease

Foorkey is also a viral disease belonging to the genus Nanovirus and family Nanoviridae [14] and causal agent is Cardamom Bushy Dwarf Virus (CBDV). This disease is characterized by dwarf tillers with small slightly curled pale green leaves; and mainly reported in lower altitude. The tillers do not grow beyond a few inches in height and appear bushy. The virus induces reduction in size of leafy and leaf shoots of the infected plants. The diseased plants remain unproductive and gradually degenerated. Foorkey symptoms appear both on seedlings and grown up plants [15]. It is also called as stunt mosaic virus of LC. Locally, this disease is called *Jurjure*. This disease is observed mainly during rainy season. It is also sap transmissible disease and is mainly transmitted by aphids (*Mollitrichosiphum* sp.). Clonal propagation from diseased plants is main reason of spread of this disease. So, regular rouging, uprooting and burying the infected samples in an isolated region were mentioned to minimize the spread of these two viral diseases in the affected plantations [16].

Its' severity is found to be low in high hills. Earlier Ilam was found to be affected by this disease but

now-a-days its severity is low. A total of 10% clumps were infected by this disease [11]. In Kolbung, Panchkanya and Godak of Ilam district losses due to this disease was found to be more than 50%. Low incidence of this disease was found in Taplejung district but in some part of Khejunum showed more incidence of this disease. Medium severity exists in Dhankuta and Panchthar. Similarly, low severity is found in Tehrathum district.

3.3.5 Distribution of chhirkey disease

Chhirkey is an important viral disease of LC and is also called as Streak Mosaic Virus. The causal agent of chhirkey disease is Large Cardamom Chhirkey Virus (LCCV), genus *Macluravirus*, family *Potyviridae*. It was reported since last 50 years in most of the growing areas with significant losses in production. This disease is characterized by light and dark green streaks on leaves. This disease is transmitted by aphids (*Rhopalosiphum maidis* and *Brachycaudus helichrysi*) in a non-persistent manner [17]. The yield loss due to this disease was reported at ranging from 33.97% to 52.84% in the affected plants but Raychoudhuri experiment showed up to 85.2% loss [18].

Chhirkey was initially seen in Ilam district with high severity and later transmitted to other districts. It is the second priority disease of Ilam after rhizome rot [12]. Low severity of this disease is seen in Taplejung and Tehrathum district. This disease is not transmitted through seed so propagation from seed is recommended. Treatment of disease by 0.075% of hydroquinone and soil drenching with 0.1% of thiouracil showed virus inhibition [18,19].

3.3.6 Distribution of Immature capsule problem

Immature capsule refers to unripening of capsule that means at the time of harvesting seeds are still white in color. This problem is new in LC orchard. Since last few years, this disease is affecting quality parameter as well as weight of capsule. Etiology of this problem is unknown. During this study it was reported that the severity of this disease in more than 50% in all the districts and nearly 80% in Panchthar district.

Most problematic disease was reported to be foorkey followed by rhizome rot, leaf blight and chhirkey in descending order during 1998 [20]. Whereas at present severity of foorkey disease

has found low. Chhirkey affected leaves shows streak mosaic which after drying causes degeneration of clumps resulting in reduction in yield and production. Foorkey causes dwarf tillers and small leaves and stunted growth of suckers which results in whole plant drying. Rhizome rot shows wilting symptoms and death of plants. All these effects of diseases result in decline in production and productivity. It is general perception that LC cultivation is being severely suffered by the disease problems. Majority of farmers estimated the loss to be more than 30%. Foorkey was the burning problem with rhizome rot in second position as problematic disease.

4. CONCLUSION AND RECOMMENDATIONS

The study identified the LC decline due to disease complex and its' spread to new areas are main problem of decreasing production and productivity of LC. The complete collapse of the LC plants from old orchards in Ilam is havoc among farmers. The farmers are hesitating to replant due to fear of disease complex. In this context, following plan of actions has been suggested for the management of LC decline:

1. The NARC should focus on developing technologies in the following areas:
 - a. Management of disease complex
 - b. Location specific varietal development
 - c. Nutrient and water management
 - d. Insect management
 - e. Orchard husbandry including shade tree management, mulching, inter-cultural operation, sanitation
 - f. Effect of climate change on cardamom.
2. Capacity building of all actors of cardamom value chain
3. Production and supply of virus-free nursery plants
4. Nurserymen/owners, firms and commercial farmers should be trained on appropriate nursery management techniques for the production and multiplication of saplings.
5. Develop and enforce effective quarantine system within the country to control disease spread within cardamom growing district
6. Carry out awareness campaign and empower the stakeholders with technical

and technology knowledge and skill for the overall decline management

7. Destroy diseased orchards and replant with assured healthy planting materials.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shrestha KP. Marketing of Large Cardamom in Mechi hills, Nepal. Paper accepted and published in November in International Journal of Graduate Research and review.). Socioeconomics Agriculture Research Policy Division (SARPOD), Nepal Agricultural Research Council (NARC); 2018a.
2. Shrestha KP. Growth trends analysis of large cardamom in Nepal. Paper submitted to Nepalese Horticulture Society. Socio-economics Agriculture Research Policy Division (SARPOD), Nepal Agriculture Research Council (NARC), Nepal; 2018b.
3. Kaini BR. A policy paper for the overall development of Large Cardamom sectors in an economically viable and environmentally sustainable manner in Nepal. Paper presented in National dialogue on setting up an Autonomous entity for Large Cardamom Development in Nepal. Hotel Mall Lainchaur, Kathmandu; 2018.
4. Mo ALMC. Statistical information on nepalese agriculture, ministry of agriculture, land management and cooperatives, monitoring, evaluation and statistical division, agriculture statistics section. Singha Durbar, Kathmandu, Nepal; 2017.
5. NSCDP. Annual Report 2015/16. National Spice Crop Development Program, Government of Nepal, Ministry of Agriculture Development, Khumaltar, Lalitpur, Nepal; 2016.
6. Shrestha J, Prasai HK, Timsina KP, Shrestha KP, Pokhrel D, Poudel K, Yadav M. Large cardamom in Nepal: Production practice and economics, processing and marketing. Nepal Agriculture Research Council, National Commercial Agriculture Research Program, Pakhribas, Dhankuta, Nepal; 2018. [ISBN: 978-9937-0-5098-2]
7. Chaudhary R, Ghimire SK, Joshi BK, Ojha BR, Niroula RK, Yadav PK. Genetic diversity in large cardamom (*Amomum subulatum* Roxb.) dissected using RAPD markers. Int. J. of Adv. Res. 2016;4(6): 1443-1451.
8. Shrestha KP, Karn PL, Shrestha CB. A study report on large cardamom, marketing in Nepal and India. Nepal Agriculture Research Council, Agriculture Research Station, Pakhribas, Dhankuta; 2001.
9. Chaudhary R, Subedi M. Distribution of large cardamom cultivars in eastern hills of Nepal. Proceedings of 11th Outreach Workshop. RARS, Lumle; 2014.
10. Adhikari PP, Sigdel K. Activities of cardamom development center and DoA in large cardamom development. In: Chaudhary R, Vista SP (eds). Proceedings of the Stakeholders consultation workshop on large cardamom development in Nepal held in April 20, 2015. Commercial Crop Division, NARC, Khumaltar, Nepal; 2015.
11. Srivastava LS. Control of pest and diseases and revitalization of Large cardamom in the project area. Commercial Agriculture Development Project, MoAD, Kathmandu; 2012.
12. Dhakal DP, Shrestha SK. Status of diseases of large cardamom in nepal. agricultural research for enhancing livelihood of Nepalese people. In: (Joshi BK, Joshi SL, Paudel KP (Eds), Proceedings of 2nd SAS Convention, 30 July-1 August 2003, Kathmandu; 2004.
13. Yadav PK, Chaudhary R, Shrestha S, Shrestha KP. Farmers perception on disease and insect incidences in Large Cardamom: A case study of Mechi Zone, Nepal. Proceedings of 11th Outreach Workshop, RARS, Lumle; 2014.
14. Mandal B, Mandal S, Pun KB, Varma A. First report of association of nanovirus with foorkey disease of large cardamom in India. PI Dis. 2004;88:428.
15. Varma PM, Capoor SP. 'Foorkey' disease of large cardamom. Indian J. Agri. Sci. 1964;34:56-62.
16. Chattopadhyay SB, Bhowmik TP. Control of 'Foorkey' disease of large cardamom in West Bengal. Indian J. Agri. Sci. 1965;35:272-275.
17. Mandal B, Vijayanandraj S, Shilpi S, Pun KB, Singh V. Pant, Varma A. Disease distribution and characterization of a new maclura virus associated with Chhirkey disease of large cardamom. Annals of Applied Biology. 2012;160(3):225-236.
18. Paudel J, Belbase S, Gautam S, Bhusa R. Viral disease of large cardamom their

- effect on production and their management. International Journal of Research studies in Agriculture Sciences (IJRSAS). 2018;4(2):21-26.
19. Raychowdhury SP, Gangly B. Transmission of chhirkey disease of large cardamom by aphid species. Indian J. Ent. 1965;27:272-276.
20. CCD. Annual Report 2063/64 (2007/08). Commercial Crop Division, Nepal Agricultural Research Council, Khumaltar, Lalitpur, Nepal; 2009.

© 2018 Shrestha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://prh.sdiarticle3.com/review-history/27736>