



**International Journal of Plant & Soil Science**  
3(6): 751-762, 2014; Article no. IJPSS.2014.6.017



SCIENCEDOMAIN international  
[www.sciencedomain.org](http://www.sciencedomain.org)

---

## Effects of Macro- and Nano-cobalt Oxide Particles on Barley Seedlings and Remediation of Cobalt Chloride Toxicity Using Sodium Hypochlorite

Sonia<sup>1</sup> and A. K. Thukral<sup>1\*</sup>

<sup>1</sup>Department of Botanical and Environmental Sciences, Guru Nanak Dev University, Amritsar, Punjab- 143005, India.

### Author's contribution

This work was carried out by the first author Sonia. The second author AKT supervised the work and developed software for statistical analysis.

Original Research Article

Received 4<sup>th</sup> December 2013  
Accepted 17<sup>th</sup> March 2014  
Published 28<sup>th</sup> March 2014

---

### ABSTRACT

This study was undertaken to determine the comparative effects of cobalt (II, III) oxide (Co<sub>3</sub>O<sub>4</sub>) macro- and nano-particles and cobaltous chloride hexahydrate (CoCl<sub>2</sub>.6H<sub>2</sub>O) on seed germination, growth and some biochemical parameters of *Hordeum vulgare* L. seedlings. Macro- and nano-Co was added to the sand medium at four levels (50 to 200 mg kg<sup>-1</sup> sand). Macro-Co was found to increase the growth of both shoots and roots at concentrations up to 200 mg Co kg<sup>-1</sup> sand. Increase in concentration of nano-Co decreased the root length. Lipid peroxidation was maximum at 200 mg Co kg<sup>-1</sup> sand for macro-Co in roots. Increase in the lipid peroxidation was found in nano-Co treated roots and shoots. Nano- and macro-Co<sub>3</sub>O<sub>4</sub> behaved differently with respect to effects on barley seedlings. The present study also demonstrated the ameliorative effect of NaOCl against CoCl<sub>2</sub>.6H<sub>2</sub>O toxicity in barley seedlings. NaOCl also decreased the lipid peroxidation induced by CoCl<sub>2</sub>.6H<sub>2</sub>O and increased chlorophyll content in seedlings.

**Keywords:** Detoxification; heavy metals; nanotoxicology; sodium hypochlorite; *Hordeum vulgare* L.

---

\*Corresponding author: E-mail: [akthukral.gndu@gmail.com](mailto:akthukral.gndu@gmail.com);

## 1. INTRODUCTION

Fast pace of industrialization and irrational uses of natural resources have led to metal accumulation in the environment. Metal accumulation in soil is of great concern in agriculture due to its adverse effects on food safety and marketability, plant growth and soil microflora and fauna [1]. Metal toxicity has high impact on the plants which consequently affect the whole ecosystem due to interdependence of living organisms. Cobalt (Co) is a transition metal with atomic number 27 and atomic weight  $58.9 \text{ g mol}^{-1}$ . The role of Co in nutrition of leguminous plants is well known, but its importance to the rest of the plant species is still ambiguous [2]. It is an essential element for the synthesis of various enzymes and coenzymes like vitamin B<sub>12</sub> (cyanocobalamin), which are required for human and animal nutrition. Co is safer for consumption up to 2.5 - 3.0 mg daily, without any adverse health effects [3]. It acts as a coenzyme in a number of cellular processes including the oxidation of fatty acids and the synthesis of DNA. Toxic concentrations of Co inhibit active transport in plants. Relatively higher concentrations of Co have toxic effects, including leaf fall, inhibition of greening, discolored veins, premature leaf closure and reduced shoot weight [4].

Two salts of Co are used in industry on a large scale, Co (II, III) oxide, also known as  $\text{CoO} \cdot \text{Co}_2\text{O}_3$  or  $\text{Co}_3\text{O}_4$ , macro- and nano-scale particles which are insoluble in water and cobalt chloride ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ , macroscale particles, water soluble). Nano- $\text{Co}_3\text{O}_4$  is a recent discovery and needs to be investigated in detail.  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  is toxic at higher concentrations.

Nanotechnology is the engineered convergence of biology, chemistry and informatics at nanoscale. The products of these exertions are called nanomaterials, consisting of nanoparticles (NPs), having a size smaller than 100 nm in at least one dimension. Among the latest technological innovations, nanotechnology possesses the top position [5]. The properties of nanomaterials raise concern about their potential adverse effects on biological systems at cellular level. Because of their small size, NPs get incursion into the living cell membrane. In contrast to the classical macroscale particles, due to their smaller size and huge surface area, NPs may interact more expeditiously with biological systems. Metal oxide-based NPs are increasingly used in applications such as opacifiers, fillers, catalysts, semiconductors, cosmetics, microelectronics etc. [6]. Therefore, interaction between inorganic nanoparticles and biological systems is one of the most promising areas of research in modern nanoscience and technology.

The present work is aimed at studying the differential effects of macro- and nano-particles of  $\text{Co}_3\text{O}_4$  and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  in combination with sodium hypochlorite (NaOCl) on barley seedlings in sand medium.  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  helps in color change in glass industry, organic synthesis and electroplating objects, production of pigments in ceramics and as a mordant in dry cleaners.  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  is a catalyst used for metal surface treatment also. The waste from these industries contains Co more than the prescribed limit. Such industrial effluents when reaching the crop fields cause toxicity to plants [7]. So, to remediate Co rich soil we have tried to use sodium hypochlorite (NaOCl) for detoxification. NaOCl converts transition metal complexes into their oxides [8]. NaOCl is used in the pesticide and textile industries and is a disinfectant, cleaner and bleach.

## 2. MATERIALS AND METHODS

### 2.1 Study Material

Certified and disease-free seeds of barley (*Hordeum vulgare* L.) variety PL-426 were purchased from Punjab Agricultural University, Ludhiana (India). Barley is generally grown as a summer crop in temperate areas and winter crop in tropical areas (including India). It is an important cereal of India, ranking next to wheat, maize and rice in the world.

### 2.2 Macro- and Nano-Co<sub>3</sub>O<sub>4</sub> Treatments

Salts of Co and other chemicals used in the study were purchased from Sigma-Aldrich, Bangalore, India; HIMEDIA Laboratory Pvt Ltd; Loba Chemie Pvt Ltd and BTL Research Lab. Suspensions of both macro- and nano-Co<sub>3</sub>O<sub>4</sub> were made in distilled water. Different concentrations of both macro- and nano-Co<sub>3</sub>O<sub>4</sub> containing 0, 50, 100, 150 and 200 mg Co kg<sup>-1</sup> sand were prepared respectively.

### 2.3 CoCl<sub>2</sub>.6H<sub>2</sub>O and NaOCl treatments

Seeds of barley were grown in sand containing various binary combinations of CoCl<sub>2</sub>.6H<sub>2</sub>O and NaOCl (Table 1). Growth and biochemical parameters were studied for any modulation in CoCl<sub>2</sub>.6H<sub>2</sub>O toxicity to seedlings.

**Table 1. CoCl<sub>2</sub>.6H<sub>2</sub>O treatments (given in numerator) in binary combinations with NaOCl treatments (given in denominator)**

| NaOCl conc. in sand medium (mg kg <sup>-1</sup> ) | CoCl <sub>2</sub> .6H <sub>2</sub> O conc. (mg kg <sup>-1</sup> ) in sand medium |          |          |          |           |
|---------------------------------------------------|----------------------------------------------------------------------------------|----------|----------|----------|-----------|
|                                                   | 0                                                                                | 250      | 500      | 750      | 1000      |
| 0                                                 | 0/0                                                                              | 250/0    | 500/0    | 750/0    | 1000/0    |
| 250                                               | 0/250                                                                            | 250/250  | 500/250  | 750/250  | 1000/250  |
| 500                                               | 0/500                                                                            | 250/500  | 500/500  | 750/500  | 1000/500  |
| 750                                               | 0/750                                                                            | 250/750  | 500/750  | 750/750  | 1000/750  |
| 1000                                              | 0/1000                                                                           | 250/1000 | 500/1000 | 750/1000 | 1000/1000 |

### 2.4 Sand Cultures and Plant Material Raising

Seeds of *H. vulgare* were surface sterilized with 0.01% HgCl<sub>2</sub> and then washed under running tap water for 10 min. After that, the seeds were soaked in distilled water for 1 h for imbibition. Sand was filtered through sieve of 300 nm size, washed with 0.1 N HCl and thrice with deionised water and was dried on filter paper in the oven at 80 - 85°C for 3 days. The imbibed seeds were then sown in polypropylene plastic jars of diameter 11 cm containing 0.5 kg sand supplemented with different concentrations of Co. In each jar, 30 seeds of nearly the same size were sown. These sand cultures were maintained at a temperature of 25±0.5°C, 70 – 80% relative humidity and 16:8 hour dark: light photoperiod (1700 lux). Then, different plant parts (shoots, roots) were harvested after 7 days of growth for the estimation of root length (RL) and shoot length (SL), fresh weight (fw) and dry weight (dw). Biochemical parameters were studied in terms of oxidative stress caused by metal salts. These included lipid peroxidation and estimation of chlorophyll content. Malondialdehyde (MDA) was estimated according to Heath and Packer [9] and chlorophyll content was measured by the method described by Arnon [10].

## 2.5 Statistical Analysis

The experimental data were expressed as mean  $\pm$  SE. One-way and two-way analysis of variance (ANOVA) were done to check the significance of differences within and between treatments and interactions if any. Significance levels of F-ratios were checked at  $P = 0.05$ . Honestly significant differences (HSD) were calculated using Tukey's multiple comparison test at  $P = 0.05$ . Difference between any two means in ANOVA if larger than the HSD value reveals a statistically significant difference. Linear regression and multiple linear regression with interaction analyses were carried out in MS-Excel using self-coded software. Pearsonian correlation and multiple correlation analyses were done to determine the significance of correlativity among the variables. Unitless beta ( $\beta$ ) regression coefficients in multiple regression analysis were calculated in order to measure the relative effects of independent variables (Co, NaOCl and Co $\times$ NaOCl interaction) on the dependent variable [11].

## 3. RESULTS

### 3.1 Growth Characteristics

#### 3.1.1. Co<sub>3</sub>O<sub>4</sub> macro- and nano-particles treatment

Seedlings cultured in sand medium containing Co<sub>3</sub>O<sub>4</sub> (macro) showed increase in root and shoot length with increase in Co concentration (50, 100, 150 and 200 mg kg<sup>-1</sup>). Further it was observed that treatment of Co<sub>3</sub>O<sub>4</sub> nano-particles significantly increased shoot length but decreased root length (Table 2).

**Table 2. Effect of Co<sub>3</sub>O<sub>4</sub> macro- and nano-particles on root length (RL) and shoot length (SL) (mean  $\pm$  S.E.) of *H. vulgare* seedlings**

| Co <sub>3</sub> O <sub>4</sub> (mg kg <sup>-1</sup> ) | Macro-particles |                 | Nano-particles  |                 |
|-------------------------------------------------------|-----------------|-----------------|-----------------|-----------------|
|                                                       | RL (cm)         | SL (cm)         | RL (cm)         | SL (cm)         |
| 0                                                     | 08.8 $\pm$ 0.60 | 13.8 $\pm$ 0.30 | 12.2 $\pm$ 1.40 | 15.0 $\pm$ 1.70 |
| 50                                                    | 09.1 $\pm$ 0.80 | 14.3 $\pm$ 0.30 | 11.6 $\pm$ 2.10 | 16.0 $\pm$ 2.40 |
| 100                                                   | 09.9 $\pm$ 0.30 | 14.6 $\pm$ 0.10 | 11.4 $\pm$ 1.70 | 16.4 $\pm$ 1.80 |
| 150                                                   | 10.2 $\pm$ 0.40 | 15.0 $\pm$ 0.10 | 10.5 $\pm$ 0.67 | 17.7 $\pm$ 1.80 |
| 200                                                   | 11.0 $\pm$ 0.10 | 15.5 $\pm$ 0.50 | 10.1 $\pm$ 0.85 | 18.2 $\pm$ 1.90 |
| F- ratio (* $P = 0.5$ )                               | 6.84*           | 12.07*          | 4.74*           | 5.17*           |
| HSD                                                   | 1.53            | 0.84            | 2.01            | 2.72            |

#### 3.1.2 CoCl<sub>2</sub>.6H<sub>2</sub>O treatments in binary combinations with NaOCl

A significant decrease in shoot, root length and fresh weight (fw) dry weight (dw) of *H. vulgare* was observed upon addition of various concentrations (250, 500, 750 and 1000 mg kg<sup>-1</sup>) of Co as CoCl<sub>2</sub>.6H<sub>2</sub>O. Further the role of NaOCl as a potent inhibitor of CoCl<sub>2</sub>.6H<sub>2</sub>O is elucidated in Tables 3, 4 and 5. 750 mg kg<sup>-1</sup> of NaOCl concentration increased shoot length of seedlings grown in 1000 mg kg<sup>-1</sup> Co amended sand by 58.57 % and root length by 86.67%. 500 mg kg<sup>-1</sup> of NaOCl increased shoot fresh weight of 1000 mg kg<sup>-1</sup> Co treated seedlings by 91.5%. Two-way ANOVA variance ratio (F) describes the statistically significant difference among shoot and root lengths on CoCl<sub>2</sub>.6H<sub>2</sub>O and the NaOCl treatments. Multiple regression models showed that Co has negative effect on shoot and

root length, while NaOCl has a positive effect. Interaction between Co and NaOCl was found to be statistically significant. Fresh and dry weight of shoots also showed significant differences (Table 6).

### 3.2 Lipid Peroxidation

Variations in shoot and root MDA content of *H. vulgare* grown in sand media containing  $\text{Co}_3\text{O}_4$  macro- and nano-particles are presented in Table 7. The MDA content of *H. vulgare* treated with macro- $\text{Co}_3\text{O}_4$  was increased significantly for shoots, while a decreasing trend was found in roots. The MDA content for both shoots and roots showed an increasing trend with increase in concentration (50, 100, 150 and 200  $\text{mg kg}^{-1}$ ) of  $\text{Co}_3\text{O}_4$  nano-particles in a dose dependent manner. The lowest value for MDA (shoots and roots) was found at concentration of 50  $\text{mg Co kg}^{-1}$  sand, while other concentrations showed increased amount of lipid peroxidation. 750  $\text{mg kg}^{-1}$  of NaOCl decreased lipid peroxidation in 1000  $\text{mg kg}^{-1}$  Co-treated shoots and roots up to 10.65% and 14.63% respectively. One-way ANOVA showed significant increase in MDA content in both roots and shoots treated with macro- and nano-Co. Two-way ANOVA revealed that there are significant differences among MDA contents of both shoots and roots in binary treatments (Table 8). The interaction between Co and NaOCl was found to be negative for both shoots and roots (Table 9).

### 3.3 Chlorophyll Estimation

The effects of  $\text{Co}_3\text{O}_4$  macro- and nano-particles and binary combinations of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  with NaOCl on chl content (chl 'a', chl 'b' and total chl) are presented in Tables 10, 11. ANOVA depicted statistical significant differences among different treatments on, chl 'b' and total chl. Multiple regression analysis showed positive effect of NaOCl on chl 'a', which as a result compensated the negative effect of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ . Co and NaOCl significantly increased the chl 'b' content, whereas in the case of total chl, Co showed negative, while NaOCl showed positive  $\beta$ - regression coefficient. It was found that chl 'a', chl 'b' and total chl showed maximum values at 200  $\text{mg kg}^{-1}$ . Significant increase was found in the chl 'a', chl 'b' and total chl contents with increase in concentration of  $\text{Co}_3\text{O}_4$  nano-particles in sand medium. Such results depicted that nano-Co modulated chlorophyll synthesis. 500  $\text{mg kg}^{-1}$  of NaOCl concentration increased chl 'a', chl 'b' and total chl contents of 1000  $\text{mg kg}^{-1}$  Co-treated leaves by 76.06 %, 79.35 % and 77.81 % respectively.

Table 3. Effect of binary treatments of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  and  $\text{NaOCl}$  on shoot length and root length (mean  $\pm$  S.E.) of *H. vulgare* seedlings

| $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$<br>( $\text{mg kg}^{-1}$ ) | $\text{NaOCl}$ ( $\text{mg kg}^{-1}$ ) |                |                 |                |                 |                |                 |                |                 |                 |
|----------------------------------------------------------------------|----------------------------------------|----------------|-----------------|----------------|-----------------|----------------|-----------------|----------------|-----------------|-----------------|
|                                                                      | 0                                      |                | 250             |                | 500             |                | 750             |                | 1000            |                 |
|                                                                      | SL (cm)                                | RL (cm)        | SL (cm)         | RL (cm)        | SL (cm)         | RL (cm)        | SL (cm)         | RL (cm)        | SL (cm)         | RL (cm)         |
| 0                                                                    | 11.2 $\pm$ 0.50                        | 8.7 $\pm$ 0.79 | 12.2 $\pm$ 0.40 | 8.9 $\pm$ 0.39 | 09.9 $\pm$ 0.40 | 9.0 $\pm$ 0.60 | 10.7 $\pm$ 0.60 | 8.8 $\pm$ 0.40 | 11.0 $\pm$ 0.90 | 09.9 $\pm$ 0.52 |
| 250                                                                  | 10.9 $\pm$ 0.50                        | 8.6 $\pm$ 0.43 | 11.1 $\pm$ 0.60 | 8.3 $\pm$ 0.42 | 10.6 $\pm$ 0.70 | 8.1 $\pm$ 0.46 | 12.2 $\pm$ 0.80 | 9.1 $\pm$ 0.45 | 11.2 $\pm$ 0.70 | 10.7 $\pm$ 0.48 |
| 500                                                                  | 09.4 $\pm$ 1.10                        | 8.4 $\pm$ 0.38 | 12.3 $\pm$ 0.60 | 9.0 $\pm$ 0.37 | 10.1 $\pm$ 0.70 | 8.7 $\pm$ 0.26 | 10.8 $\pm$ 0.50 | 9.6 $\pm$ 0.26 | 10.9 $\pm$ 0.80 | 09.6 $\pm$ 0.51 |
| 750                                                                  | 07.2 $\pm$ 1.10                        | 5.7 $\pm$ 0.79 | 11.3 $\pm$ 0.60 | 9.5 $\pm$ 0.47 | 10.7 $\pm$ 0.30 | 9.6 $\pm$ 0.25 | 09.4 $\pm$ 0.90 | 8.4 $\pm$ 0.28 | 12.1 $\pm$ 0.40 | 09.7 $\pm$ 0.42 |
| 1000                                                                 | 07.0 $\pm$ 1.10                        | 5.3 $\pm$ 0.62 | 09.8 $\pm$ 0.30 | 8.9 $\pm$ 0.30 | 09.6 $\pm$ 0.40 | 8.7 $\pm$ 0.66 | 11.1 $\pm$ 0.60 | 9.8 $\pm$ 0.48 | 11.1 $\pm$ 0.40 | 08.2 $\pm$ 0.43 |

F- ratios for shoots; 4.08\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 9.21\* ( $\text{NaOCl}$ ), 2.39 ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 2.74

F- ratios for roots; 3.47 ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 15.97\* ( $\text{NaOCl}$ ), 4.17\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 1.93

Table 4. Effect of binary treatments of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  and  $\text{NaOCl}$  on fresh weight (fw) and dry weight (dw) of shoots (mean  $\pm$  S.E) of *H. vulgare* seedlings

| $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ( $\text{mg kg}^{-1}$ ) | $\text{NaOCl}$ ( $\text{mg kg}^{-1}$ ) |                  |                 |                  |                 |                  |                 |                  |                 |                  |
|-------------------------------------------------------------------|----------------------------------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
|                                                                   | 0                                      |                  | 250             |                  | 500             |                  | 750             |                  | 1000            |                  |
|                                                                   | fw (g)                                 | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           |
| 0                                                                 | 1.08 $\pm$ 0.05                        | 0.12 $\pm$ 0.02  | 1.73 $\pm$ 0.03 | 0.15 $\pm$ 0.007 | 1.24 $\pm$ 0.02 | 0.10 $\pm$ 0.007 | 1.26 $\pm$ 0.01 | 0.11 $\pm$ 0.008 | 1.43 $\pm$ 0.03 | 0.12 $\pm$ 0.004 |
| 250                                                               | 0.84 $\pm$ 0.02                        | 0.09 $\pm$ 0.004 | 1.66 $\pm$ 0.01 | 0.15 $\pm$ 0.005 | 1.50 $\pm$ 0.01 | 0.12 $\pm$ 0.022 | 1.32 $\pm$ 0.02 | 0.11 $\pm$ 0.009 | 1.55 $\pm$ 0.06 | 0.14 $\pm$ 0.006 |
| 500                                                               | 0.81 $\pm$ 0.01                        | 0.08 $\pm$ 0.003 | 0.85 $\pm$ 0.01 | 0.08 $\pm$ 0.003 | 0.72 $\pm$ 0.03 | 0.07 $\pm$ 0.005 | 0.67 $\pm$ 0.06 | 0.05 $\pm$ 0.009 | 1.26 $\pm$ 0.02 | 0.11 $\pm$ 0.008 |
| 750                                                               | 0.75 $\pm$ 0.03                        | 0.08 $\pm$ 0.004 | 0.87 $\pm$ 0.01 | 0.07 $\pm$ 0.002 | 1.14 $\pm$ 0.01 | 0.11 $\pm$ 0.006 | 1.05 $\pm$ 0.01 | 0.09 $\pm$ 0.003 | 1.16 $\pm$ 0.02 | 0.11 $\pm$ 0.006 |
| 1000                                                              | 0.71 $\pm$ 0.01                        | 0.07 $\pm$ 0.003 | 1.11 $\pm$ 0.01 | 0.10 $\pm$ 0.008 | 1.36 $\pm$ 0.03 | 0.14 $\pm$ 0.007 | 1.15 $\pm$ 0.05 | 0.11 $\pm$ 0.007 | 1.72 $\pm$ 0.02 | 0.16 $\pm$ 0.010 |

F- ratios for shoots (fw) ; 990.59\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 915.10\* ( $\text{NaOCl}$ ), 153.83\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 0.07

F- ratios for shoots (dw) ; 81.48\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 48.05\* ( $\text{NaOCl}$ ), 16.36\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 0.02

Table 5. Effect of binary treatments of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  and NaOCl on fresh weight (fw) and dry weight (dw) of roots (mean  $\pm$  S.E) of *H. vulgare* seedlings

| $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (mg kg <sup>-1</sup> ) | NaOCl (mg kg <sup>-1</sup> ) |                  |                 |                  |                 |                  |                 |                  |                 |                  |
|------------------------------------------------------------------|------------------------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|
|                                                                  | 0                            |                  | 250             |                  | 500             |                  | 750             |                  | 1000            |                  |
|                                                                  | fw (g)                       | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           | fw (g)          | dw (g)           |
| 0                                                                | 1.21 $\pm$ 0.07              | 0.07 $\pm$ 0.003 | 1.68 $\pm$ 0.05 | 0.15 $\pm$ 0.005 | 1.02 $\pm$ 0.01 | 0.09 $\pm$ 0.005 | 1.02 $\pm$ 0.01 | 0.08 $\pm$ 0.007 | 1.13 $\pm$ 0.11 | 0.08 $\pm$ 0.004 |
| 250                                                              | 1.04 $\pm$ 0.06              | 0.07 $\pm$ 0.007 | 1.35 $\pm$ 0.13 | 0.14 $\pm$ 0.007 | 1.12 $\pm$ 0.04 | 0.10 $\pm$ 0.001 | 0.91 $\pm$ 0.01 | 0.08 $\pm$ 0.002 | 1.34 $\pm$ 0.06 | 0.09 $\pm$ 0.003 |
| 500                                                              | 0.94 $\pm$ 0.03              | 0.06 $\pm$ 0.008 | 0.67 $\pm$ 0.04 | 0.05 $\pm$ 0.006 | 0.87 $\pm$ 0.04 | 0.05 $\pm$ 0.009 | 0.54 $\pm$ 0.02 | 0.09 $\pm$ 0.004 | 1.2 $\pm$ 0.03  | 0.11 $\pm$ 0.004 |
| 750                                                              | 0.85 $\pm$ 0.05              | 0.06 $\pm$ 0.007 | 0.73 $\pm$ 0.04 | 0.06 $\pm$ 0.009 | 0.95 $\pm$ 0.04 | 0.07 $\pm$ 0.005 | 0.83 $\pm$ 0.03 | 0.09 $\pm$ 0.004 | 1.01 $\pm$ 0.01 | 0.11 $\pm$ 0.006 |
| 1000                                                             | 0.74 $\pm$ 0.04              | 0.05 $\pm$ 0.007 | 0.96 $\pm$ 0.01 | 0.13 $\pm$ 0.007 | 1.03 $\pm$ 0.02 | 0.09 $\pm$ 0.011 | 1.02 $\pm$ 0.01 | 0.09 $\pm$ 0.007 | 1.31 $\pm$ 0.07 | 0.11 $\pm$ 0.003 |

F- ratios for roots (fw) ; 162.88\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 97.04\* (NaOCl), 44.21\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 0.13

F- ratios for roots (dw) ; 71.07\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 31.17\* (NaOCl), 64.99\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 0.02

Table 6. Multiple regression models for shoot length (SL) and root length (RL), fresh weight (fw) and dry weight (dw) of *H. vulgare* seedlings in binary combination of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (mg kg<sup>-1</sup>) and NaOCl (mg kg<sup>-1</sup>)

| Dependent variable (Y) | Multiple regression equation                                                        | r                 | $\beta$ -regression coefficients |       |                   |
|------------------------|-------------------------------------------------------------------------------------|-------------------|----------------------------------|-------|-------------------|
|                        |                                                                                     |                   | Co                               | NaOCl | Co $\times$ NaOCl |
| SL (cm)                | Y= 11.69 – 0.0038 Co – 0.0008 NaOCl + 5 $\times$ 10 <sup>-6</sup> Co $\times$ NaOCl | 0.720*            | - 1.02                           | -0.22 | 0.99              |
| RL (cm)                | Y= 8.66 – 0.0017 Co + 0.0011 NaOCl + 2 $\times$ 10 <sup>-6</sup> Co $\times$ NaOCl  | 0.673*            | - 0.53                           | 0.33  | 0.41              |
| Shoot fw (g)           | Y= 1.23 – 0.0005 Co + 0.0001 NaOCl + 6 $\times$ 10 <sup>-7</sup> Co $\times$ NaOCl  | 0.58*             | - 0.61                           | 0.13  | 0.50              |
| Shoot dw (g)           | Y= 8.66 – 0.0017 Co + 0.0011 NaOCl + 2 $\times$ 10 <sup>-6</sup> Co $\times$ NaOCl  | 0.673*            | - 0.53                           | 0.33  | 0.41              |
| Root fw (g)            | Y= 1.27 – 0.0002 Co – 0.0006 NaOCl + 7 $\times$ 10 <sup>-7</sup> Co $\times$ NaOCl  | 0.56*             | -0.35                            | -0.90 | 0.79              |
| Root dw (g)            | Y= 0.095 – 0.00 Co – 0.00 NaOCl + 6 $\times$ 10 <sup>-8</sup> Co $\times$ NaOCl     | 0.47 <sup>#</sup> | - 0.12                           | -0.50 | 0.65              |

\*P= .05, <sup>#</sup>P= .10

Table 7. Lipid peroxidation ( $\mu$  mole MDA; mean  $\pm$  S.E) of *H. vulgare* seedlings after treatment with  $\text{Co}_3\text{O}_4$  macro- and nano-particles

| $\text{Co}_3\text{O}_4$ (mg $\text{kg}^{-1}$ ) | Macro-particles |                 | Nano-particles  |                 |
|------------------------------------------------|-----------------|-----------------|-----------------|-----------------|
|                                                | MDA shoots      | MDA roots       | MDA shoots      | MDA roots       |
| 0                                              | 2.72 $\pm$ 0.04 | 1.98 $\pm$ 0.04 | 1.71 $\pm$ 0.12 | 1.18 $\pm$ 0.02 |
| 50                                             | 2.43 $\pm$ 0.18 | 1.74 $\pm$ 0.01 | 1.26 $\pm$ 0.04 | 1.26 $\pm$ 0.04 |
| 100                                            | 2.24 $\pm$ 0.18 | 1.54 $\pm$ 0.01 | 1.65 $\pm$ 0.12 | 1.28 $\pm$ 0.01 |
| 150                                            | 2.48 $\pm$ 0.03 | 1.50 $\pm$ 0.03 | 1.78 $\pm$ 0.06 | 1.64 $\pm$ 0.11 |
| 200                                            | 2.99 $\pm$ 0.03 | 0.91 $\pm$ 0.05 | 1.97 $\pm$ 0.06 | 1.71 $\pm$ 0.12 |
| <b>F-ratio</b> (*P= .05)                       | 17.77*          | 466.81*         | 63.05*          | 31.99*          |
| HSD                                            | 0.31            | 0.086           | 0.15            | 0.19            |

Table 8. Lipid peroxidation ( $\mu$  mole MDA ; mean  $\pm$  S.E) of *H. vulgare* seedlings after binary treatments with  $\text{CoCl}_2.6\text{H}_2\text{O}$  and NaOCl

| $\text{CoCl}_2.6\text{H}_2\text{O}$ (mg $\text{kg}^{-1}$ ) | NaOCl (mg $\text{kg}^{-1}$ ) |                 |                  |                 |                  |                 |                  |                 |                 |                 |
|------------------------------------------------------------|------------------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|-----------------|-----------------|
|                                                            | 0                            |                 | 250              |                 | 500              |                 | 750              |                 | 1000            |                 |
|                                                            | MDA shoots                   | MDA roots       | MDA shoots       | MDA roots       | MDA shoots       | MDA roots       | MDA shoots       | MDA roots       | MDA shoots      | MDA roots       |
| 0                                                          | 2.76 $\pm$ 0.03              | 0.39 $\pm$ 0.05 | 2.96 $\pm$ 0.032 | 0.45 $\pm$ 0.05 | 2.54 $\pm$ 0.012 | 0.42 $\pm$ 0.08 | 2.15 $\pm$ 0.006 | 0.37 $\pm$ 0.07 | 3.17 $\pm$ 0.01 | 0.52 $\pm$ 0.12 |
| 250                                                        | 2.87 $\pm$ 0.02              | 0.50 $\pm$ 0.06 | 2.93 $\pm$ 0.006 | 0.68 $\pm$ 0.15 | 2.65 $\pm$ 0.006 | 0.48 $\pm$ 0.03 | 2.28 $\pm$ 0.006 | 0.44 $\pm$ 0.06 | 2.99 $\pm$ 0.01 | 0.56 $\pm$ 0.06 |
| 500                                                        | 2.96 $\pm$ 0.03              | 0.59 $\pm$ 0.06 | 3.11 $\pm$ 0.006 | 0.52 $\pm$ 0.03 | 2.36 $\pm$ 0.006 | 0.63 $\pm$ 0.07 | 3.21 $\pm$ 0.05  | 0.40 $\pm$ 0.06 | 2.89 $\pm$ 0.01 | 0.53 $\pm$ 0.03 |
| 750                                                        | 3.12 $\pm$ 0.08              | 0.69 $\pm$ 0.11 | 3.49 $\pm$ 0.005 | 0.51 $\pm$ 0.16 | 2.88 $\pm$ 0.006 | 0.53 $\pm$ 0.07 | 3.03 $\pm$ 0.02  | 0.49 $\pm$ 0.02 | 3.08 $\pm$ 0.03 | 0.63 $\pm$ 0.07 |
| 1000                                                       | 3.66 $\pm$ 0.04              | 0.82 $\pm$ 0.08 | 2.96 $\pm$ 0.017 | 0.39 $\pm$ 0.04 | 2.09 $\pm$ 0.006 | 0.54 $\pm$ 0.01 | 3.27 $\pm$ 0.01  | 0.70 $\pm$ 0.08 | 3.2 $\pm$ 0.006 | 0.44 $\pm$ 0.07 |

F- ratios for MDA (Shoots ) ; 399.79\* ( $\text{CoCl}_2.6\text{H}_2\text{O}$ ), 850.19\* (NaOCl), 262.63\* ( $\text{CoCl}_2.6\text{H}_2\text{O}\times\text{NaOCl}$ ) , \*P= .05, HSD= 0.09

F- ratios for MDA (Roots ) ; 8.37\* ( $\text{CoCl}_2.6\text{H}_2\text{O}$ ), 4.79\* (NaOCl) , 6.22\* ( $\text{CoCl}_2.6\text{H}_2\text{O}\times\text{NaOCl}$ ) , \*P= .05, HSD= 0.21

Table 9. Multiple regression models for lipid peroxidation ( $\mu$  mole MDA  $\text{g}^{-1}$  tissue) in shoots and roots, and chl content (mg  $\text{g}^{-1}$  fw) of *H. vulgare* in binary combinations of  $\text{CoCl}_2.6\text{H}_2\text{O}$  and NaOCl

| Dependent variable (Y) | Multiple regression equation                                                                        | r                 | $\beta$ -regression coefficients |        |          |
|------------------------|-----------------------------------------------------------------------------------------------------|-------------------|----------------------------------|--------|----------|
|                        |                                                                                                     |                   | Co                               | NaOCl  | CoxNaOCl |
| MDA shoot              | $Y = 2.71 - 0.0005 \text{ Co} - 2 \times 10^{-5} \text{ NaOCl} - 2 \times 10^{-7} \text{ CoxNaOCl}$ | 0.40 <sup>#</sup> | 0.48                             | -0.016 | -0.16    |
| MDA root               | $Y = 0.44 - 0.0002 \text{ Co} + 4 \times 10^{-5} \text{ NaOCl} - 2 \times 10^{-7} \text{ CoxNaOCl}$ | 0.52*             | 0.76                             | 0.14   | 0.53     |
| Chl 'a'                | $Y = 5.35 - 0.0013 \text{ Co} - 0.0009 \text{ NaOCl} + 2 \times 10^{-6} \text{ CoxNaOCl}$           | 0.27              | -0.44                            | -0.28  | 0.56     |
| Chl 'b'                | $Y = 2.48 - 0.0008 \text{ Co} - 0.0002 \text{ NaOCl} + 2 \times 10^{-6} \text{ CoxNaOCl}$           | 0.37 <sup>#</sup> | -0.35                            | -0.11  | 0.58     |
| Total Chl              | $Y = 7.83 - 0.0021 \text{ Co} - 0.0011 \text{ NaOCl} + 4 \times 10^{-6} \text{ CoxNaOCl}$           | 0.31              | -0.40                            | -0.21  | 0.59     |

\*P= .05, #P= .10



Table 10. Chlorophyll content (mean  $\pm$  S.E) of *H. vulgare* after treatment with  $\text{Co}_3\text{O}_4$  macro- and nano-particles

| $\text{Co}_3\text{O}_4$ (mg $\text{kg}^{-1}$ ) | Chl Content                     |                  |                                 |                  |                                   |                 |
|------------------------------------------------|---------------------------------|------------------|---------------------------------|------------------|-----------------------------------|-----------------|
|                                                | Chl 'a' (mg $\text{g}^{-1}$ fw) |                  | Chl 'b' (mg $\text{g}^{-1}$ fw) |                  | Total Chl (mg $\text{g}^{-1}$ fw) |                 |
|                                                | Macro-particles                 | Nano-particles   | Macro-particles                 | Nano-particles   | Macro-particles                   | Nano-particles  |
| 0                                              | 0.60 $\pm$ 0.004                | 0.61 $\pm$ 0.01  | 0.13 $\pm$ 0.004                | 0.12 $\pm$ 0.006 | 0.73 $\pm$ 0.003                  | 0.73 $\pm$ 0.02 |
| 50                                             | 0.37 $\pm$ 0.02                 | 0.49 $\pm$ 0.02  | 0.19 $\pm$ 0.003                | 0.18 $\pm$ 0.003 | 0.54 $\pm$ 0.003                  | 0.68 $\pm$ 0.03 |
| 100                                            | 0.45 $\pm$ 0.04                 | 0.52 $\pm$ 0.021 | 0.21 $\pm$ 0.004                | 0.19 $\pm$ 0.003 | 0.65 $\pm$ 0.005                  | 0.73 $\pm$ 0.07 |
| 150                                            | 0.52 $\pm$ 0.01                 | 0.54 $\pm$ 0.05  | 0.23 $\pm$ 0.03                 | 0.26 $\pm$ 0.03  | 0.76 $\pm$ 0.01                   | 0.79 $\pm$ 0.02 |
| 200                                            | 0.62 $\pm$ 0.003                | 0.68 $\pm$ 0.03  | 0.28 $\pm$ 0.04                 | 0.27 $\pm$ 0.02  | 0.91 $\pm$ 0.003                  | 0.94 $\pm$ 0.02 |
| <b>F- ratios (*P= .05)</b>                     | 78.25*                          | 22.72*           | 21.72*                          | 44.11*           | 1805.92*                          | 26.54*          |
| HSD                                            | 0.45                            | 0.57             | 0.44                            | 0.33             | 0.12                              | 0.75            |

Table 11. Chlorophyll content (mg  $\text{g}^{-1}$  fw) of *H. vulgare* seedlings after binary treatments with  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  and NaOCl

| $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$<br>(mg $\text{kg}^{-1}$ ) | NaOCl (mg $\text{kg}^{-1}$ ) |                 |                 |                  |                  |                 |                  |                 |                 |                  |                 |                 |                   |                 |                 |
|---------------------------------------------------------------------|------------------------------|-----------------|-----------------|------------------|------------------|-----------------|------------------|-----------------|-----------------|------------------|-----------------|-----------------|-------------------|-----------------|-----------------|
|                                                                     | 0                            |                 |                 | 250              |                  |                 | 500              |                 |                 | 750              |                 |                 | 1000              |                 |                 |
|                                                                     | Chl 'a'                      | Chl 'b'         | Total Chl       | Chl 'a'          | Chl 'b'          | Total Chl       | Chl 'a'          | Chl 'b'         | Total Chl       | Chl 'a'          | Chl 'b'         | Total Chl       | Chl 'a'           | Chl 'b'         | Total Chl       |
| 0                                                                   | 0.52 $\pm$ 0.06              | 0.25 $\pm$ 0.05 | 0.77 $\pm$ 0.05 | 0.49 $\pm$ 0.07  | 0.25 $\pm$ 0.07  | 0.75 $\pm$ 0.03 | 0.41 $\pm$ 0.004 | 0.18 $\pm$ 0.02 | 0.59 $\pm$ 0.02 | 0.42 $\pm$ 0.004 | 0.21 $\pm$ 0.02 | 0.62 $\pm$ 0.02 | 0.40 $\pm$ 0.09   | 0.18 $\pm$ 0.03 | 0.58 $\pm$ 0.03 |
| 250                                                                 | 0.51 $\pm$ 0.04              | 0.24 $\pm$ 0.04 | 0.75 $\pm$ 0.05 | 0.58 $\pm$ 0.02  | 0.26 $\pm$ 0.04  | 0.85 $\pm$ 0.07 | 0.40 $\pm$ 0.004 | 0.19 $\pm$ 0.05 | 0.59 $\pm$ 0.01 | 0.55 $\pm$ 0.04  | 0.27 $\pm$ 0.04 | 0.82 $\pm$ 0.03 | 0.47 $\pm$ 0.05   | 0.22 $\pm$ 0.07 | 0.68 $\pm$ 0.01 |
| 500                                                                 | 0.44 $\pm$ 0.02              | 0.21 $\pm$ 0.07 | 0.66 $\pm$ 0.05 | 0.41 $\pm$ 0.004 | 0.20 $\pm$ 0.002 | 0.62 $\pm$ 0.03 | 0.72 $\pm$ 0.06  | 0.34 $\pm$ 0.03 | 1.06 $\pm$ 0.08 | 0.62 $\pm$ 0.07  | 0.28 $\pm$ 0.03 | 0.90 $\pm$ 0.05 | 0.352 $\pm$ 0.041 | 0.18 $\pm$ 0.03 | 0.53 $\pm$ 0.03 |
| 750                                                                 | 0.43 $\pm$ 0.03              | 0.20 $\pm$ 0.02 | 0.64 $\pm$ 0.05 | 0.36 $\pm$ 0.041 | 0.19 $\pm$ 0.010 | 0.55 $\pm$ 0.06 | 0.48 $\pm$ 0.02  | 0.25 $\pm$ 0.05 | 0.73 $\pm$ 0.04 | 0.59 $\pm$ 0.08  | 0.29 $\pm$ 0.02 | 0.88 $\pm$ 0.07 | 0.16 $\pm$ 0.04   | 0.06 $\pm$ 0.01 | 0.22 $\pm$ 0.02 |
| 1000                                                                | 0.38 $\pm$ 0.01              | 0.18 $\pm$ 0.04 | 0.56 $\pm$ 0.05 | 0.45 $\pm$ 0.01  | 0.23 $\pm$ 0.01  | 0.68 $\pm$ 0.03 | 0.66 $\pm$ 0.03  | 0.33 $\pm$ 0.01 | 0.99 $\pm$ 0.06 | 0.61 $\pm$ 0.02  | 0.53 $\pm$ 0.01 | 1.14 $\pm$ 0.03 | 0.55 $\pm$ 0.04   | 0.25 $\pm$ 0.05 | 0.79 $\pm$ 0.1  |

F- ratios for Chl 'a', for binary treatments; 13.88\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 25.84\* (NaOCl), 13.24\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 1.41


F- ratios for Chl 'b' for binary treatments; 20.82\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 32.89\* (NaOCl), 11.52\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 0.94

F- ratios for Total Chl for binary treatments; 56.05\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ), 106.98\* (NaOCl), 42.43\* ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O} \times \text{NaOCl}$ ), \*P= .05, HSD= 1.20

#### 4. DISCUSSION

Heavy metals may cause major occupational and environmental hazards due to their non-biodegradable nature and long biological half life period [12]. Exposure to heavy metals is mainly due to the anthropogenic actions such as use of fertilizers, agrochemical compounds, sewage sludge and other activities like mining [13]. Such activities result in the transportation of metal ions via air and water, which ultimately bind to soil and sediments. Co is a relatively rare magnetic element with properties similar to those of iron and nickel and occurs in nature primarily as arsenides, oxides and sulphides. Most of the production of Co involves the metallic form used in the formation of Co superalloys [14]. The distribution of Co in plants is entirely species specific.

A significant increase in both root and shoot length was observed in 7 days old seedlings treated with  $\text{Co}_3\text{O}_4$  macro-particles, while treatment of  $\text{Co}_3\text{O}_4$  nano-particles increased only shoot length. These observations are in accordance with earlier studies [15] where cobalt is said to increase the seedling growth by alleviating the senescence of aged tissue by inhibiting the activity of 1-aminocyclopropane-1-carboxylate (ACC) oxidase and reducing ethylene production.  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  was found to be toxic at higher concentrations as was observed from decreased root and shoot length. NaOCl decreased  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  induced toxicity in both root and shoot length. NaOCl is known to transform Co into its oxide form either through exclusion, inclusion (i.e. sequestration and compartmentalization of metal ions in organelles) or chelation binding [16]. The reaction of  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  with NaOCl is given below:



The reason for such an observation might be attributed to the fact that NaOCl oxidises the more toxic  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  to the less toxic  $\text{Co}_3\text{O}_4$  [8]. At treatments where NaOCl was absent altogether, metal-caused toxicity resulted in reduction of shoot length. Lowest shoot length was observed at concentrations where Co is in maximum and NaOCl is in minimum amounts. The amount of NaOCl required for counteracting toxicity caused by Co is more in the case of roots as compared to shoots. This may be attributed to the fact that roots are accumulative organs of heavy metals [17].

Lipid peroxidation was found to be maximum for roots at a concentration of  $200 \text{ mg kg}^{-1}$  of  $\text{Co}_3\text{O}_4$ . The reason for such a trend can be attributed to increased production of ROS which induce membrane destabilization resulting in the formation of peroxides, as was reported by Mead et al. [18]. On the other hand,  $\text{Co}_3\text{O}_4$  inhibited lipid peroxidation by decreasing the MDA content in roots and the differences obtained were statistically significant.

A significant reduction in chlorophyll content (chl 'a', chl 'b' and total chl) induced by  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  as compared to the untreated control might be due to overproduction of reactive oxygen species, which in turn could have damaged chloroplast membrane [19] as was observed on the effect of Co on *Cajanus cajan* Mill. Application of NaOCl in combination with  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  increased levels of chl 'a', chl 'b' and total chl. Protection extended by NaOCl was evident from the fact that it significantly reduced ROS production as was observed in lipid peroxidation studies.

## 5. CONCLUSION

Our results showed that  $\text{Co}_3\text{O}_4$ , both nano- as well as macro-particles showed differential toxic effects on *H. vulgare* seedlings. Furthermore, the application of NaOCl significantly reduced the toxicity caused by  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  in *H. vulgare* seedlings. Improved Co stress mitigation by NaOCl involves biochemical ramifications. Thus, the present study presents NaOCl as effective candidate in ameliorating  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  toxicity.

## ACKNOWLEDGEMENT

The authors are thankful to the University Grants Commission, and the Department of Science and Technology, Government of India, New Delhi for financial assistance. Thanks are also due to the Head of the Department of Botanical and Environmental Sciences for providing access to research facilities.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Nagajyoti CP, Lee DK, Sreekanth MVT. Heavy metals, occurrence and toxicity for plants: a review. *Environmental Chemistry Letters*. 2010;8:199-216.
2. Collins RN, Kinsela AS. Pedogenic factors and measurements of the plant uptake of cobalt. *Plant Soil*. 2011;339:499–512.
3. Hokin B, Adams M, Ashton J, Louie H. Comparison of dietary cobalt intake in three different Australian diets. *Asia Pacific Journal of Clinical Nutrition*. 2004;13:289-291.
4. Ayeni OO, Ndakidemi PA, Snyman RG, Odendaal JP. Chemical, biological and physiological indicators of metal pollution in wetlands. *Scientific Research and Essays*. 2010;5:1938-1949.
5. Nair R, Varghese SH, Nair BG, Maekawa T, Yoshida Y, Kumar DS. Nanoparticulate material delivery to plants. *Plant Science*. 2010;179:154-163.
6. Mortimer M, Kasemets K, Heinlaan M, Kurvet I, Kahru A. High throughput kinetic *Vibrio fischeri* bioluminescence inhibition assay for study of toxic effects of nanoparticles. *Toxicology In Vitro*. 2008;221:412-417.
7. Husain A, Ashhar MM, Javed I. Analysis of industrial wastewater in Aligarh city. *Journal of Chemical and Pharmaceutical Research*. 2014;6:614-621.
8. Lister MW. Decomposition of sodium hypochlorite: the catalyzed reaction. *Canadian Journal of Chemistry*. 1956;34:479-488.
9. Heath RL, Packer L. Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics*. 1968;125:180-198.
10. Arnon DI. Copper enzymes in isolated chloroplasts polyphenoloxidase in *Beta vulgaris*. *Plant Physiology*. 1949;24:1-15.
11. Sokal RR, Rohlf FJ. *Biometry: the Principles and Practice of Statistics in Biological Research*. WH Freeman and Co. San Francisco. 1981;859.
12. Barbier O, Jacquillet G, Tauc M, Cougan M, Poujeol P. Effect of heavy metals on, and handled by, the kidney. *Nephron Physiology*. 2005;99:105-110.
13. Schützendubel A, Polle A. Plant responses to abiotic stresses: heavy metals- induced oxidative stress and protection by mycorrhization. *Journal of Experimental Botany*. 2002;53:1351-1365.

14. Barceloux DG. Cobalt. *Clinical Toxicology*. 1999;37:201-216.
15. Li CZ, Wang D, Wang GX. The protective effects of cobalt on potato seedling leaves during osmotic stress. *Botanical Bulletin of Academia Sinica*. 2005;46:119-125.
16. Jayakumar K, Jaleel CA. Uptake and accumulation of cobalt in plants: a study based on exogenous cobalt in soyabean. *Botany Research International*. 2009;2:310-314.
17. Singh R, Gautam N, Mishra A, Gupta R. Heavy metals and living systems: an overview. *Indian Journal of Pharmacology*. 2011;43:246-253.
18. Mead JF, Wu GS, Stain RA, Gelmont D, Sevanian A, Sohlbeg E, McElhaney RN. Mechanism of the protection against membrane peroxidation. In: Yagi K, editor. *Lipid Peroxides in Biology and Medicine*, London Academic Press. 1982;161-173.
19. Gopal R. Antioxidant defense mechanism in pigeon pea under cobalt stress. *Journal of Plant Nutrition*. 2014;37:136-145.

---

© 2014 Sonia and Thukral; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sciencedomain.org/review-history.php?iid=472&id=24&aid=4160>