



Postharvest Losses and Management of Horticultural Produce: A Review

Abhishek Bisht^{a*} and Sury Pratap Singh^a

^a *Department of Food Technology, School of Allied Sciences, Dev Bhoomi Uttarakhand University, Navgaon, Manduwala, Chakrata Road, Dehradun -248007 (Uttarakhand), India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JSRR/2024/v30i31881

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112719>

Review Article

Received: 14/12/2023

Accepted: 17/02/2024

Published: 21/02/2024

ABSTRACT

Horticultural postharvest losses pose a significant challenge to the supply chain and economic growth, particularly in impoverished nations. This review paper explores the multifaceted issues surrounding postharvest losses in horticultural crops and emphasizes the critical role of technological innovations in mitigating these losses. Various stakeholders within the value chain, including farmers, handlers, and consumers, face substantial risks due to insufficient postharvest handling techniques. The study underscores the urgency of adopting advanced postharvest technologies, focusing on enhanced harvesting and packaging strategies. By using cutting-edge technologies, it is possible to lower losses after harvest and raise the general standard of fresh produce. The improper use of packing materials and inadequate packaging methods significantly contribute to these losses, making it imperative to address these aspects for sustainable solutions. This comprehensive review not only identifies the reason for fruit and vegetable losses after harvesting within the countries but also provides actionable insights into mitigating measures. This article intends to support the creation of a robust and successful horticulture supply chain by addressing the issues and offering workable solutions. The integration of technological advancements in harvesting, handling, and packaging processes emerges as a key element in the overall strategy to reduce postharvest losses and promote economic growth in resource-constrained regions.

*Corresponding author: E-mail: bisht.abhishek11@gmail.com;

Keywords: Developing nation; fruits; vegetables; canning; storehouse; traceability.

1. INTRODUCTION

Items from horticulture are necessary to maintain people's diets and health. They are rich source of water soluble vitamins like (niacin, riboflavin, ascorbic acid and thiamine) as well as fat soluble vitamin like (retinol , tocopherol) etc. also greater than 90 percent of ascorbic acid in people's diet is obtained from fruits and vegetables. They are an excellent source of dietary fiber, calcium, phosphorus, iron, and magnesium along with this it increases the good micro-flora in the intestine [87]. Crops, veggies, fruits, flowers, spices, and aromatic and therapeutic plants are all considered horticultural crops, are primarily cultivated for their culinary value and appeal, with their consumption patterns influenced by factors such as availability, quality, and affordability. Unlike staple cereals or essential industrial crops, horticultural crops do not occupy a central role in agriculture, according to [4]. The Indian horticulture industry has expanded due to the joint efforts of farmers, government, and investors. The horticulture industry has shown positive growth due to the support from government agencies and educational institutions in terms of research and training of professionals [10]. Fruit and vegetable losses after harvest can amount to as much as fifty percent since fresh food has a limited shelf life and there are inadequate facilities for processing and marketing it. Inadequate pre-production and post-harvest management practices exacerbate these losses, adversely affecting prices for consumers, farmer revenue, and produce's nutritional content. Subpar quality of harvested produce stems from the use of inadequate planting materials and less-than-ideal habits of cultivation, such as methods of harvest and managing. Perishable food without proper storage on the farms or access to a pack house packing station is often sold immediately after harvest without processing or wrapping. Solid debris from urban horticulture crops can clog drains and cause water blockages and cause environmental pollution with unsanitary conditions [26].

Following the harvest, there may be qualitative as well as quantitative losses. The term "quantitative losses" describes how food loses

bulk or volume, decreasing the volume of food suitable for people to consume. These losses might be valued in terms of weight, money, or energy. Before food is thrown out, eaten, or used in another way, it experiences qualitative losses. These losses include decreases in the produce's nutritive content, edibility, safety, and eventually economic worth. One of the main issues with after-harvesting systems and growth in the economy is qualitative degradation, in particular the growing countries, as improper postharvest handling techniques put impoverished value chain participants, such as farmers, handlers, and consumers, at serious risk. While the lack of frequently linked to qualitative losses, this is less of an issue for those who can afford to upgrade their logistical systems [87]. While both developed and developing countries experience substantial postharvest losses and waste in horticulture products, the degree of these reductions varies according to the handling chain's step. Developing countries typically suffer greater losses during the post harvest and handling stages, primarily due to inadequate infrastructure and improper handling practices. In contrast, Waste at the retail and consumer levels is often higher in industrialized nations. Often attributable to stringent cosmetic standards and consumer preferences [43]. Around a third of the global population global fruit and vegetables production is believed to be lost in the postharvest period, excluding inedible waste. In the UK, postharvest losses from harvest to retailer are approximated at 9 percent, omitting goods that, because of quality and esthetic requirements, were left in the field. In 2008, projections indicated that the United States experienced losses of \$42.8 billion in fruits and vegetables at the commercial and consumption stages, equating to approximately \$141 per person [17]. From this standpoint, it is becoming progressively crucial to mitigate or ideally eliminate the waste of fresh fruits and vegetables. Consequently, it is imperative to implement advancements in postharvest technology, including suitable harvesting methods and packing techniques, to diminish postharvest losses and increase the overall quality of fresh produce. This will ultimately lead to increased consumption of fresh fruits [88,89].




Chart 1. Top worldwide fruit production shares
Source: [28]




Chart 2. Top worldwide vegetables production shares
Source: [28]

The main forms of deterioration that contribute to the disposal of fresh fruits and vegetables include moisture loss, bruising, and eventual decay [44,65]. It is not feasible to entirely halt the natural transformation of fresh food; there are measures that can be taken to mitigate these changes. These include maintaining a low temperature, regulating relative humidity during storage, ensuring proper packaging and shipping, and so forth [3].

2. REASONS FOR AGRICULTURE LOSSES AFTER HARVEST

Increasing food production serves as a viable solution to address the growing demand for substance. Instead of merely focusing on boosting output, it is more prudent to ensure the efficient delivery of food to consumers while safeguarding agricultural products against loss that occurs during several phases, including processing, retailing, fields, shipping, and storage. This approach aims to mitigate the strain on water resources, fields, and the environment. Following production, agricultural products undergo before reaching customers, there are several after harvest unit processes dealing with stages and storage procedures, all of which have specific losses. Consequently, the

cumulative effect of these steps results in a diminished availability of food. Recent research conducted in India reveals specific percentages significant losses that occur after harvest, after-harvest activities, handling and storage for various crops, including grains, pulses, oilseeds, [54].

2.1 Harvesting

Fruit and vegetable quality and shelf life are greatly affected by the date of harvest. Premature harvesting, driven by financial concerns, can result in unripe produce with poor flavor and increased susceptibility to damage. Conversely, ripe fruits have a lower shelf life. Overripe and under ripe fruits are more likely to cause physiological issues. The nutritive and financial worth of crops is diminished by early harvesting. In certain instances, entire batches may be lost due to unripe or overripe produce [7,41]. Harvesting methods may result in losses as well [46].

These losses can be increased several times when farmers not having the proper cold or containers for storing either during or after the collecting process. This resulted in deterioration of the quality of fresh horticultural products and produce.




Fig. 1. An example of the waste generated along the whole fresh fruit and vegetable distribution chain

Inadequate implementation of cultural practices” is replaced with “inadequate cultural practices”. “Including pre harvest irrigation, fertilization, pruning, infections, physiological disorders, hail, drought,excessive rainfall, and plant nutrition” is replaced with “including a variety of factors”. Furthermore,“improper harvesting techniques, and inaccurate harvest timing” is substituted for “inaccuracies in determining the appropriate harvest time, harvesting at the wrong moment, improper use of harvesting techniques, and failure to pre-cool crops like cherries during the harvest season.” result in losses that range from (4%) to (12%) [60,59].

2.2 Storehouse

Storage technologies provide opportunities to enhance marketing efforts, boost value, and overcome quarantine limitations. Additionally they play a pivotal role in advancing and globalizing the horticulture sectors. Temperature serves as the foundational technology for various storage systems, and effective temperature management remains irreplaceable. While other storage technologies can enhance its impact, controlling relative humidity, optimizing the storage environment, implementing proper packing, and utilizing chemical treatments can all be coordinated alongside cold storage [80]. The metabolism of ethylene is complexly impacted by temperature. Elevated production of ethylene during refrigerated storage might indicate damage to perishable items that are sensitive to cold conditions [22]. While volatile aroma components play a critical role in strawberry flavor, the impact of low-temperature storage on these compounds has received less attention compared to other quality factors. Following

harvest, strawberries produce a diverse range of volatile compounds. Although this emission persists throughout chilly storage, the temperature has an impact on the particular chemicals that are created [32,49].

2.3 Transportation

The movement of goods from producers to consumers is a crucial aspect of commerce, particularly for newly manufactured products. Direct delivery from producers to consumers is an efficient approach. Transportation costs have a significant impact on consumer prices, especially in complex distribution system that serves large cities and remote locations. In some cases, these transportation costs may even exceed the price of the components that are needed to produce the goods [21]. The preservation of perishable food faces challenges in less affluent countries due to insufficient transportation, poor road conditions, and ineffective management of logistics. Additionally in numerous nations, individuals lacking literacy and skills often handle loading and unloading tasks, leading to careless treatment of goods and consequent mechanical damage to agricultural products [7].

2.4 Improper Packaging

To maximize losses and prolong the freshness of fresh f&V, proper packaging is essential. Inadequate packaging techniques and the use of unsuitable packing material significantly contribute to post-harvest losses. Insufficient protection against damage can accelerate the deterioration of fresh produce when low-quality packaging materials are utilized. Unfortunately, the widespread use of inexpensive but subpar

packing material is a common practice globally, especially in developing and underdeveloped nations [44].

2.5 Consumer's Waste

Consumer preferences refer to the views that individuals hold regarding a product, whether it is a service or an item, and are shaped through the evaluation of the choices currently accessible [45]. Analyzing consumer behavior encompasses more than just examining what individuals purchase; it also delves into the reason behind their purchases, the timing, location, manner, and frequency of their acquisitions, aiming to

fulfill their needs and desires [69]. Roughly (50%) of the food is discarded by households consists of fresh fruits and veggies [29]. A report states that fruits and vegetables contribute to (39%) of total domestic waste [85]. Another study indicates a similar proportion of around (40%) in household garbage. Factors leading to financial wastages among consumers involve excessive spending, inadequate home storage practices, and insufficient preparation. The quantity of waste produced by individuals is impacted by various factors, such as material possessions and sociocultural aspects, including income, gender, lifestyles, and choices in home storage [63].



Fig. 2. Improper packaging



Fig. 3. Improper loading



Fig. 4. Not proper market



Fig. 5. Not proper storage



Fig. 6. Not proper market for sale



Fig. 7. Not proper regulation



Fig. 8. Improper transport facilities



Fig. 9. Not proper market



Fig. 10. Not proper handling



Fig. 11. Improper transportation

3. OTHERS FACTORS LEADING TO POST HARVEST LOSSES

3.1 Insufficient Structure

It is important to note that unlike other enduring agricultural goods, fruits and vegetables need to be promptly transported to markets due to their quick perishability any delay could lead to loss [18,73,76]. For this reason, efficient transportation is essential to the horticultural product supply chain. Examples of insufficient infrastructure include limited transit choices, an underdeveloped transportation network, impractical roadways, and the absence of affordable temperature-controlled vehicles [1,19,47,50,52,55,57,71].

3.2 Absence of Communication between Farmers and Facilities

Processing businesses usually acquire raw materials through intermediaries or agents instead of directly from farmers, leading to reduced prices for farmers and increased costs for processors [72]. A key issue revolves around linkage, influencing the accessibility of suitable techniques used in fruits and vegetables

processing. The ratio of raw materials to completed goods tends to increase due to this connection factor. It's noteworthy that agricultural seasonality affects processing sectors, leading to price fluctuations. Implementing measures to stabilize F&V markets not only mitigates the risk of price fluctuations but also eases the exchange of valuable pricing information between producers and processing units [67].

3.3 Ignorance of High-Quality Seeds

Optimal weather conditions, top-notch seeds, and fertile soil take into consideration while trying to improve agricultural output [2,53,58]. Several Indian farmers lack awareness regarding the benefits of using seeds of excellent quality for agricultural cultivation, including improved wellness and greater resilience to pests and illnesses [57].

3.4 Inadequate Facilities for Tracking Traceability, and Record Keeping

From the agricultural field to the retailer, traceability ensures the monitoring of specific farm products within a supply chain. Adequate documentation is essential at every point in the

distribution process to facilitate item tracking. Companies must recognize food items, gather relevant data, and share it with various participants in the distribution chain to ensure effective food tracing [23,76,83,84]. The effectiveness of link for supplies is raised by effective information exchange and coordination between the many parties [20,34]. When traceability is implemented well, commodity prices are lowered and, in the event of a recall, distributed products may be retrieved from the same vendor. Nonetheless, the technology has its limits. For example, there is significant implementation and maintenance costs associated with the traceability facilities and the tracking software may not be user-friendly for all supply chain participants [57].

3.5 Many Middleman

From the field to the store, traceability guarantees on monitoring of specific agricultural products throughout the supply chain. Thorough documentation is crucial at every phase of the distribution chain to make item tracking easier. Companies must recognize food items, gather pertinent data, and disseminate it among various participants within the system of supply to ensure efficient commodity tracing. Effective exchange of information and coordination among the various stakeholders enhances the effectiveness of the manufacturing chain. In India, the agricultural supply system is significantly affected by powerful traditional traders with political clout. The excessive involvement of intermediaries in the distribution process disrupts the seamless flow of crucial information, resulting in coordination challenges and decreased overall efficiency. These intermediaries, who profit significantly, contribute to a decline in farmers' earnings. The addition of extra commission brokers in the supply chain might boost product turnover but could also lead to financial deprivation [9,68].

3.6 Failure to Consume

Unexpected changes to meal habits can lead to food being wasted, even for customers who make mindful food selections [25]. Or time constraints could require the substitution or removal of some items [11]. F&V from the previous regular shop may be swapped out for new ones in certain circumstances. "Fresher," products and disposed of [25]. Customers sometimes lack knowledge on how to determine if fruit is ripe or how to ripen it and certain fruits

are overlooked if they take an excessively lengthy time to mature [37].

3.7 The Setting for Food

Providing consumers with dependable and trustworthy access to nutritious and safe food is an essential aspect of the food industry [51,33]. Domestic food waste appears to be a bigger problem in wealthy metropolitan areas that are dependent on international supply systems, according to trends [61,75]. F&V that were before only available during certain seasons are now offered year-round for these wealthy customers [75]. Almost every home has at least one refrigerator. Considering that the single most significant element influencing the preservation of F&V quality is temperature [82]. Keeping F&V at room temperature in the house is probably going to cause it to deteriorate quickly. When people choose to store their fruits and vegetables in refrigerators, the way they manage the inventory within the refrigerator can result in waste, particularly if products are missed or are thought to have become contaminated by coming into touch with other foods [79]. Among the primary reasons why food is wasted among US consumers is concern regarding food contamination [56]. The most important part of food disposal is frequently the habits related to emptying refrigerators of past-due produce. Some people do this all the time, throwing away goods they feel are no longer fresh or won't provide the desired sensory experience. However, other customers take a more impromptu approach, carrying out these purges when a certain component is needed for a dish [79]. Rather than the features of residential refrigerators, these disposal practices are probably going to be a bigger source of F&V waste.

4. FRUITS AND VEGETABLES LOSSES AFTER HARVEST IN DEVELOPING NATION

It is expected that the changing tastes of middle-class consumers in developing nations like China and India, who have more purchasing power, would significantly impact global food consumption. By the middle of this century, there may be a 50–70% increase in the global food consumption due to this [35,61,13]. Even while the global need for food is growing, about a billion people continue to struggle with insufficient food and nutritional deficits [13,64]. Upcoming food safety issues are heavily

impacted by environmental factors like water shortage, deforestation, and global warming. It is the maintenance of a delicate equilibrium to guarantee an enough quantity of wholesome and nutritious food for the global population [81,46]. Food wastage has been prioritized in the global combat hunger and for enhanced food safety due to its ability to increase food accessibility and environmental well-being. Reducing FLW promotes sustainable environmental practices, strengthens food security, and increases food supply [29,70].

5. METHODS TO REDUCE LOSSES AFTER HARVEST

Various post-harvest processes must be used to preserve crops nutrition, color, and texture. Despite the natural decline in the quality of horticultural produce over time, the application of specific procedures and standard storage techniques can effectively slow down degradation, thereby extending shelf or storage life.

5.1 Drying

The most ancient method of preserving food involves extracting moisture from it, known as dehydration, to prolong its durability. In this method, the moisture content of the fruit's tissue diminishes gradually, significantly reducing the likelihood of fruit decay. In tropical regions, this

technique can be achieved by utilizing the heat from direct sunlight. Despite the discoloration of fruit tissues during the drying process, this approach represents the earliest form of drying, which has undergone enhancements through various methods over time [36]. Different methods for drying are available, such as dehydrator dryers that utilize artificial heat with controlled humidity, temperature, and airflow. Oven dryers also regulate temperature through the oven's heat source. Various traditional drying techniques have been employed, including low-temperature drying within the range of 15 to 50°C and high-temperature drying at temperatures exceeding 50°C. Freezing fruit juice and other liquids at temperatures between -10 and -40°C is another approach for drying [5].

5.2 Canning

The lifespan of the produce is prolonged through preservation in a sealed container. This approach applies to perishable acidic fruits and vegetables, involving storage in a container and heat treatment. The tomato, for instance, is immersed in boiling water for about a minute and then placed within a can that contained contents halfway with acetic acid, Vitamin C, or citric acid to facilitate the peeling of the skin [62]. The act of acidification involves the addition of substances such as vinegar or lemon juice. This process is employed to halt the proliferation of the harmful bacterium *Clostridium botulinum*.




Chart 3. Post harvest losses at global level

Source:[27]

Table 1. Harvest and post harvest losses of fruits and vegetables in India

Crop	Harvesting (%)	Collecti on (%)	Sorting (%)	Packaging (%)	Trans portati on (%)	Total loss in farm operations (%)	Farm level storage (%)	Cold storage (%)	Wholesaler level storage (%)	Retailer level (%)	Processing unit level storage (%)	Total loss in storage (%)	Overall loss (%)
Apple	4.6	0.4	4.8	0.1	1.2	11.1	0.04(2.3)	0.12(1.5)	0.52(1.0)	0.23(1.1)	0.29(1.7)	1.2	12.3
Banana	1.3	0.4	0.9	0.4	1.1	4.2	0.04(1.6)	0.16(3.3)	1.83(2.4)	0.36(2.4)	0.01(0.3)	2.4	6.6
Citrus	0.9	0.5	1.8	0.3	1.3	4.8	0.03(1.9)	0.00(0.0)	0.69(1.3)	0.77(2.3)	0.01(0.2)	1.5	6.3
Grapes	0.9	0.2	3.2	0.3	1.9	6.6	0.41(5.5)	-	0.54(1.6)	0.84(2.2)	0.30(2.7)	1.7	8.3
Guava	4.4	1.2	4.6	0.9	2.8	13.9	0.41(2.1)	-	1.83(5.9)	1.80(3.8)	0.06(5.7)	4.1	18.0
Mango	4.1	0.7	2.8	0.5	2.5	10.6	0.06(1.5)	-	0.92(2.5)	0.93(2.7)	0.19(0.9)	2.1	12.7
Papaya	1.4	0.3	2.0	0.2	1.1	5.1	0.08(2.1)	0.00(0.0)	1.02(2.3)	1.20(2.4)	0.00(0.0)	2.3	7.4
Sapota	1.5	0.2	1.4	0.1	1.1	4.3	0.02(0.8)	-	0.75(1.7)	0.73(1.7)	-	1.5	5.8
Cabbage	1.1	1.1	1.6	0.3	1.3	4.6	0.14(2.1)	0.06(1.1)	0.88(2.2)	1.19(2.6)	0.03(2.3)	2.3	6.9
Cauliflower	0.8	0.3	1.7	0.2	1.9	4.8	0.06(1.5)	0.04(0.5)	1.00(2.2)	0.89(2.3)	-	2.0	6.8
Green pea	3.5	1.1	3.3	0.2	0.5	8.6	0.06(1.2)	0.01(0.3)	0.71(1.3)	0.92(2.4)	-	1.7	10.3
Mushroom	1.4	1.8	4.3	1.6	2.0	11.0	-	-	-	1.50(1.7)	-	1.5	12.5
Onion	2.7	0.2	1.6	0.1	0.4	5.2	0.54(2.7)	0.38(2.2)	0.38(2.2)	0.81(2.2)	0.01(0.1)	2.3	7.5
Potato	3.2	0.7	2.2	0.1	0.5	6.7	0.36(3.9)	0.78(1.4)	0.96(3.9)	0.19(2.4)	0.01(0.4)	2.3	9.0
Tomato	1.7	1.1	3.2	0.8	3.1	9.9	0.17(4.6)	0.01(1.6)	1.23(2.7)	0.98(2.3)	0.11(2.0)	2.5	12.4
Tapioca	3.6	0.5	1.5	0.5	1.3	7.5	1.09(4.1)	-	0.58(1.5)	0.44(1.7)	0.19(2.3)	2.3	9.8

Source: [54]

Table 2. Shrinking produce diversity in U.S.A stores

Fresh fruit and vegetables	U.S retail weight, total (millions lbs./year)	Shrink in 2011 (%)	Shrink in 2012 (%)	Uneaten whole fresh fruit (millions lbs./year)	Nonedible share (%)	Fruit loss (edible) (millions lbs./year)
Papaya	288	54.8	30.3	124	33	83
Pineapple	1917	30.5	35.8	617	49	315
Apricots	29	39	28.9	9	7	8
Watermelon	4187	23.3	27.7	1063	48	553
Honeydew	444	33.8	18.5	100	54	46
Mango	743	22.8	20.8	157	31	108
apples	4856	20	19.2	932	10	839
Avocados	1578	25	17.2	300	26	222
Grapefruit	723	25.2	14.6	136	50	68
Turnip greens	0.4	61.7	63.9	71	30	50
Mustard greens	0.4	60.4	61.6	76	7	70
Okra	0.4	35.6	53.5	49	14	42
Kale	0.3	30.7	24.2	28	39	17
Squash	4.3	24	22.9	309	17	256
Radishes	0.4	17.9	27.2	29	10	26
Eggplant	0.8	18.1	22.9	51	19	41
Pumpkin	4.7	16.5	21.9	267	30	187
Cauliflower	1.1	17.5	17.3	59	61	23
tomatoes	17.3	11.9	14.7	790	9	719

Source: [16]

5.3 Freezing

This method involves preserving fruit by placing it in a freezer. When a liquid drops below its freezing point, it undergoes a phase change referred to as freezing. By employing this method, the fruit's lifespan is extended because it hinders the growth of microorganisms, reducing degradation. A vital aspect of this preservation method is the fruit's capacity to preserve its color and sustain a relatively high nutritional quality [62]. Cold storage rooms and freezers use a variety of freezing techniques. These consist of cryogenic freezing, liquid immersion freezing, air blast freezing, and contact (plate) freezing (Johnston, 1994). When employing the freezing technique, with the predominant methods being air blast freezing and contact freezing, commonly utilized for preserving fruits and vegetables [14].

5.4 Fresh Storage

Preserving the optimal physical and physiological environments to prolong the shelf life constitutes an integral aspect of fresh storage. Fruits that are stored are kept in their original, fresh state, with efforts focused on minimizing the degradation process and metabolic mechanisms to extend shelf life without compromising freshness. To achieve the desired storage temperature, cooling

methods are employed. Various chilling techniques, such as evaporative coolers, controlled environment storage, and freezers, are applied to keep tomato fruits in storage [6].

5.5 Refrigerators

Perishables are stored around the globe today using this extremely well-known and well-established method. For a few days, it is an effective way to store fruits and vegetables [66]. Refrigeration, the method employed to lower the temperature of a space below that of its environment, involves various techniques such as the gas cycle, thermoelectric, vapor compression, and vapor adsorption. Commonly found in residential refrigerators, the vapor compression cycle plays a key role in cooling by employing a compressor, evaporator, condenser, and refrigerant. In Heat is transmitted from the items that are kept to the refrigerant during this procedure, achieving the desired refrigeration effect [12].

5.6 Regulated Atmospheric Pressure

The idea behind a controlled environment is to delay a fruit's respiratory process by raising the level of CO₂ and lowering the level of O₂ in the atmosphere. This extension of shelf life is

achieved by altering the usual air composition [24]. The duration fruits remain fresh can be prolonged by combining them with refrigeration, which alters the temperature within the storage space. The composition of standard air includes 21% O₂, 78% N₂, 0.93% Ar, 0.04% CO₂, and fluctuating levels of water vapor. The controlled atmosphere is stabilized by the introduction of a non reacting gas, specifically nitrogen [8].

5.7 Use of Ethylene Action Inhibitors

Sometimes, to increase the amount of time that product may be stored, waxing, high carbon dioxide levels, low oxygen levels, and ripening inhibitors are combined [48,74]. Finding the best remedies for endogenous ethylene (C₂H₄) and ripening inhibition, however, is still difficult. As a result, many chemical combinations have been tested in order to maintain ethylene levels below the cutoff. When employed in conjunction with regulated storage atmospheres, ethylene absorbents like potassium permanganate (KMnO₄) and calcium chloride (CaCl₂) have considerable economic potential, albeit small-scale farmers may not be able to afford those [74].

5.8 Developments in Cold Chain Management and Logistics

The cornerstones of postharvest technology have been and now are maintaining the cold chain from harvest to market and optimizing storage temperature for fruits and vegetables [40]. Although ideal storage and transit settings may be envisioned in theoretical models, the real world is frequently messier. Sophisticated logistics systems require thorough monitoring since deviations from ideal circumstances are common [38]. A new paradigm for inventory management in the retail industry is made possible by the combination of temperature data with large primary producer databases that include information on variety, yield, location, season, pre-harvest procedures, and quality evaluations. This change enables shops to prioritize the consumption of items depending on their expiration date by switching from the antiquated FIFO model to the more advanced FEFO approach [38]. While the FEFO strategy involves a thorough grasp of the produce's quality journey, the FIFO model merely requires information on the produce's arrival date. This includes evaluating the product's initial quality upon arrival, taking into account the

temperatures throughout storage and transit, and finally predicting the product's remaining shelf life. Growers' labels now only contain limited information on variety, identity, provenance, and harvest time. However, information technology is about to make cloud-based, dynamic data available that includes all pre- and post-harvest factors. With the use of this improved data, virtual labels will be created, enabling a dynamic FEFO model for efficient produce management.

5.9 Developments in Retail Packaging

Packaging does more than just hold food; it actively prevents Waste and loss of food across the whole distribution system, maintaining its freshness and quality until it gets to your table [78]. In the world of produce, readily available "retail packages" like apple bags, banana bunches, or individually wrapped zucchinis provide several enticing advantages beyond simple convenience. These pre-packaged options offer informative labels, protection from damage, improved hygiene, extended shelf life, and in some cases, even controlled atmospheres to optimize freshness [78]. Creating an efficient packaging approach for fruits and vegetables involves customizing a strategy. Each packaging unit should ensure appropriate cooling, ventilation, and temperature regulation to suit the respiration patterns and environmental preferences of individual products, with a particular focus on managing low O₂ and high CO₂ levels [42]. By modifying the air inside the package, MAP acts as a shield, protecting the delicate freshness and quality of mangoes, strawberries, and even prepared kale during their transport and shelf life [15]. While plastic sheets and containers have long been the standard for packaging, edible waxes, and coatings have become more well-known as reliable alternatives in recent years. These substitutes, which are acceptable to eat and are made of proteins, lipids, and polysaccharides often in composite forms, offer similar protective barriers between food and surroundings [77].

5.10. Efforts to raise consumer awareness

Food waste that goes unnoticed is common; polls show that people are shockingly unaware of how much food is wasted and how it affects the surroundings and the finances [56]. 1,862 customers in the UK were interviewed, and the results indicate a general underestimation of personal food waste quantities [48].




Fig. 12. Additional methods to lower postharvest losses in fruits and vegetables

Public education on food waste is still vital because it gives people knowledge about how to reduce the amount of trash they produce. One notable example of this is the United Kingdom WRAP-initiated "LOVE FOOD HATE WASTE" crusade. Remarkably, even if the total amount of households increased by 4 percent, throughout five years (2007–2012), this initiative successfully reduced household food and beverage waste by 1.3 million tons [86].

5.11 Publication of Suggested Storage Guidelines

A crucial bottleneck in the supply chain, wholesale marketplaces' uneven cooling and poor temperature control can result in reduced shelf life and substantial fruit and vegetable loss from incorrect storage. This emphasizes how important it is to share best practices at this critical juncture [40]. Providing clear, personalized storage rules for customers and shop management is crucial for effectively tackling the problem of fruit and vegetable waste. Specific temperature and humidity requirements apply to each variety of food, and by communicating this information across the supply chain, the danger of rotting may be significantly reduced [63]. "The Food Keeper" was developed in the U.S as an outcome of cooperation between the Food Marketing Institute, and the USDA. This book gives consumers with helpful guidance on how to store a range of goods in pantry, refrigerator, and freezer areas, including fresh fruit [31].

6. CONCLUSION

During the fight against poverty, ensuring that everyone has access to enough wholesome food, and maintaining the freshness and quality of products, reducing waste after fruits and vegetables are picked may be a highly effective weapon. Poor infrastructure at every stage of the process, from picking to storing to transporting and processing is the primary cause of wasted fruits and vegetables in underdeveloped nations. As a crucial tactic for attaining ecological and worldwide food safety, recently, nations and global bodies have prioritized and directed their focus toward FLW prevention. The UN has set an extraordinarily ambitious goal to reduce world food wastage by 2030 as a result of this. However, the most of the trash in wealthy countries comes from the consumers. Implementing advanced technology for real-time monitoring and management of post-harvest processes, coupled with comprehensive training programs for farmers on best practices in handling and storage, can significantly reduce future post-harvest losses. Additionally, reducing waste and increasing access to nutrient-dense food is made possible by efficient coordination and management throughout the post-harvest phases.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Accorsi Gallo, Manzini R. A climate driven decision-support model for the distribution of perishable products. *Journal of Cleaner Production*. 2017;165:917-929.
2. Afadhali. Rwanda: Farmer Ignorance Blamed for Low Usage of Quality Seeds. *The East African*; 2015.
3. Ahmad, Siddiqui. Factors affecting postharvest quality of fresh fruits Postharvest quality assurance of fruits: Springer. 2015;7-32.
4. Alazar. Horticultural Marketing in Ethiopia. M.Sc. Thesis. 735 Haramaya University, Haramaya; 2007.
5. Aravindh, Sreekumar. Solar drying—a sustainable way of food processing. In *Energy Sustainability through Green Energy*. Springer India. 2015;27-46.
6. Ayomide, Ajayi, Ajayi. Advances in the development of a tomato postharvest storage system: towards eradicating postharvest losses. In *Journal of Physics: Conference Series IOP Publishing*. 2019;1378(2):022064.
7. Azabağaoğlu MÖ. Investigating fresh fruit and vegetables losses at contemporary food retailers. *Sosyal Bilimler Araştırma Dergisi*. 2018;7(4):55-62.
8. Babarinsa, Omodara, Okanlawon. Preservation of mature green tomatoes using controlled atmosphere storage under tropical conditions. *Journal of Stored Products and Postharvest Research*. 2015;6(9):72-76.
9. Balaji M, Arshinder K. Modeling the causes of food wastage in Indian perishable food supply chain. *Resources. Conservation Recycle*. 2016;114(1):153-167.
10. Banjaw Review of post-harvest loss of horticultural crops in Ethiopia, its causes and mitigation strategies. *Journal of Plant Sciences and Agricultural Research*. 2017;2(1):1-4.
11. Bava, Jaeger, Park. Constraints upon food provisioning practices in 'busy' women's lives: Trade-offs which demand convenience. *Appetite*. 2008;50(2-3):486-498.
12. Bhatt. Domestic Refrigerators: Field Studies and Energy Efficiency Improvement. 2001;60:591-600.
13. Bond, Meacham, Bhunnoo, Benton. Food waste within global food systems. Swindon, UK: Global Food Security; 2013.
14. Bouzari, Holstege, Barrett, Mineral, fiber, and total phenolic retention in eight fruits and vegetables: A comparison of refrigerated and frozen storage. *J Agric Food Chem*. 2015;63(3):951-56.
15. Brecht, Chau, Fonseca, Oliveira FAR, Silva, Nunes MCN. Bender, Maintaining optimal atmosphere conditions for fruits and vegetables throughout the postharvest handling chain. *Postharvest Biology and Technology*. 2003;27(1):87-101.
16. Buzby, Bentley, Padera, Ammon, Campuzano. Estimated fresh produce shrink and food loss in US supermarkets. *Agriculture*. 2015;5(3):626-648.
17. Buzby JC, Hyman J, Stewart H, Wells HF. The value of retail- and consumer-level fruit and vegetable losses in the United States. *Journal Consumers Affairs*. 2011;45:492-515.
18. Caixeta-Filho JV. Losses in the transportation of fruits and vegetables: A Brazilian case study. *Int. J. Logist.: Research and Applications*. 1999;2(3):325-341.
19. Cardoen, Joshi, Diels, Sarma PM, Pant D. Agriculture biomass in India: Part 2. Post-harvest losses, cost and environmental impacts. *Resources, Conservation and Recycling*. 2015;101:143-153.
20. Chen. Information sharing and supply chain coordination. *Handbooks in Operations Research and Management Science*. 2003;11:341-421.
21. Condratchi, Movileanu, The Role of transport means used in the transportation of fresh horticultural products. *Scientific Papers*. 2012;35.
22. Cooper WC, Rasmussen GK, Waldon ES. Ethylene evolution stimulated by chilling in *Citrus* and *Persea* sp. *Plant Physiology*. 1969;44:1194-1196.
23. Dabbene Gay, Tortia, Traceability issues in food supply chain management: A review. *Biosyst. Eng*. 2013;120(1):65-80.
24. El-Ramady, Domokos-Szabolcsy, Abdalla, Taha, Fári. Postharvest management of fruits and vegetables storage. In *Sustainable agriculture reviews*. Springer International Publishing. 2015;65-152.
25. Evans. Blaming the consumer—once again: The social and material contexts of everyday food waste practices in some English households. *Critical Public Health*. 2011;21(4):429-440.
26. Expert consultation meeting on postharvest and value addition of

- horticultural produce (2010). GFAR/APARI Date: 29 November – 2 December (2010). Preliminary Report on the Review Agricultural Towards; 2000.
27. FAO. Moving forward on food loss and waste reduction. The State of Food and Agriculture; 2019.
 28. FAO. Agricultural production statistics 2000–2021. In FAO eBooks; 2022.
 29. FAO. Global Food Losses and Waste. Extent, Causes and Prevention; 2011.
 30. Food and agricultural organization of the united nations (FAO). Global food losses and food waste- extent, causes and prevention; 2011.
 31. Food Marketing Institute (FMI), The Department of Food Science at Cornell University, The Department of Agriculture, U.S. The Food Keeper; 2014.
 32. Forney CF, Kalt W, Jordan MA. The composition of strawberry aroma is influenced by cultivar, maturity, and storage. *HortScience*. 2000;35:1022-1026.
 33. Gao, Knight, Zhang, Mather, Tan. Consumer scapegoating during a systemic product-harm crisis. *Journal of Marketing Management*. 2012;28(11-12):1270-1290.
 34. Gaur, Giloni, Seshadri,. Information sharing in a supply chain under ARMA demand. *Managgement Science*. 2005;51(6):961-969.
 35. Godfray HCJ, Beddington, Crute, Haddad, Lawrence, Muir, and Toulmin. Food security: The challenge of feeding 9 billion people. *Science*. 2010;327(5967):812-818.
 36. Gupta, Nath. Drying of tomatoes. *Journal of Food Science and Technology. India*. 1984;21:372-376.
 37. Harker, Jaeger, Lau, Rossiter. Consumer perceptions and preferences for kiwifruit: A review. In VI International Symposium on Kiwifruit. 2006 February;753:81-88.
 38. Jedermann, Nicometo, Uysal, Lang. Reducing food losses by intelligent food logistics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 2014;372(2017):20130302.
 39. Johnston. FAO fisheries technical paper. Freezing and refrigerated storage in fisheries. Rome: CSL Food Science Laboratory. 1994;4-94.
 40. Kader A perspective on postharvest horticulture (1978-2003). *HortScience*. 2003;38(5):1004-1008.
 41. Kader A. Maturity, ripening, and quality relationships of fruit-vegetables. Strategies for Market Oriented Greenhouse Production. 1995;434:249-256.
 42. Kader AA, Zagory D, Kerbel EL, Wang CY. Modified atmosphere packaging of fruits and vegetables. *Critical Reviews in Food Science & Nutrition*. 1989;28(1):1-30.
 43. Kader. Increasing food availability by reducing postharvest losses of fresh produce. *Acta Horti*. 2005;682:2169-2176.
 44. Kitinoja, AlHassan. Identification of appropriate postharvest technologies for small scale horticultural farmers and marketers in sub-saharan africa and south asia-part 1. Postharvest losses and quality assessments. Paper presented at the XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010): International Symposium on. 2010;934.
 45. Kotler, Keller. *Manajemen Pemasaran, Jilid 1*. Jakarta: Erlangga. Online Purchasing; 2008.
 46. Liu. Food losses and food waste in China: A first estimate. *OECD Food, Agriculture and Fisheries Papers*, No. 66. OECD Publishing, Paris; 2014.
 47. Lorentz, Yew Wong, Hilmola Emerging distribution systems in Central and Eastern Europe: Implications from two case studies. *International Journal of Physical Distribution and Logistics Management*. 2007;37(8):670-697.
 48. Lyndhurst, Cox, Downing. Food behaviour consumer research: Quantitative phase. *Waste & Resources Action Programme (WRAP)*: Banbury, UK; 2007.
 49. Miszczak A, Forney C, Prange R. Development of aroma volatiles and color during postharvest ripening of Kent. *Journal of the American Society Horticultural Science*. 1995;120: 650-655.
 50. Modi, Mishra, Gulati, Murugesan. Uttarakhand state cooperative federation: Can it help the horticulture farmers? *Vision: Journal Business Perspective*. 2009;13(2): 53-61.
 51. Morland, Wing, Roux,. The contextual effect of the local food environment on residents' diets: The atherosclerosis risk in communities study. *American Journal of Public Health*. 2002;92(11):1761-1768.
 52. Murthy, Gajanana TM, Sudha, Dakshinamoorthy V. Marketing and post-harvest losses in fruits: Its implications on availability and economy. *Indian Journal of Agricultural Economics*. 2009;64(2).

53. Mwendwa. Lack of Knowledge on Quality Seed Standards Attributed to Low Yields; 2015.
54. Nanda, Vishwakarma, Bathla HVL, Rai A, Chandra P. Harvest and post harvest losses of major crops and livestock produce in India. AICRP,(ICAR); 2012.
55. Narula. Reinventing cold chain industry in India: Need of the hour. Interview with Mr Sanjay Aggarwal. Journal of Agribusiness in Developing and Emerging Economies. 2011;1(2):97-99.
56. Neff, Spiker, Truant. Wasted food: US consumers' reported awareness, attitudes, and behaviors. Plos One. 2015;10(6):e0127881.
57. Negi, Anand.. Supply chain of fruits and vegetables agribusiness in uttarakhand (India): Major issues and challenges. Journal of Supply Chain Management Systems. 2015;4(1).
58. Nsom. Lack of Quality Seeds Hamper Vegetable Production. Cameroon Post line; 2015.
59. ÖZCAN M. Turunçgil ve Çay Yetiştiriciliği Ders Notu. Ondukuzmayıs Üniversitesi sf. 2020;105.
60. Özdemir, DüNDAR, Ertürk, Dilbaz,. Bazı yörelerimizde yetiştirilen starking delicious elmalarında derim öncesi ve derim sırasında görülen kayıpların belirlenmesi. IV. Ulusal Bahçe Bitkileri Kongresi. 2003;8-12.
61. Parfitt, Barthel, Macnaughton. Food waste within food supply chains: Quantification and potential for change to 2050. Philosophical transactions of the royal society B: Biological sciences. 2010; 365(1554):3065-3081.
62. Parnell,Suslow, Harris. California: Tomatoes: Safe methods to store, preserve, and enjoy. UCANR Publications; 2004.
63. Porat, Lichter, Terry, Harker, Buzby, Postharvest losses of fruit and vegetables during retail and in consumers' homes: Quantifications, causes, and means of prevention. Postharvest Biology and Technology. 2018;139:135-149.
64. Prevention UNEP. Reduction of Food and Drink Waste in Businesses and Households—Guidance for Governments. Local Authorities, Businesses and Other Organisations, Version. 2014;1.
65. Ray R, Ravi V. Post harvest spoilage of sweet potato in tropics and control measures. Critical Reviews in Food Science and Nutrition. 2005;45(7-8):623-644.
66. Rodriguez del CS, López Chaves. Effect of different treatments on the evolution of polyamines during refrigerated storage of eggplants. Journal of Agricultural and Food Chemistry. 2001;49(10):4700-4705.
67. Rottger. Strengthening Farm Agribusiness Linkages in Africa. Summary Results of Five Country Studies in Ghana, Nigeria, Kenya, Uganda and South Africa. Food and Agriculture Organization. 2004;6.
68. Sachan A, Sahay BS, Sharma D. Developing Indian grain supply chain cost model: A system dynamics approach. International Journal of Productivity and Performance Management. 2005;54(3): 187-205.
69. Schiffman Leon G, Kanuk Leslie L, Wisenblit Joseph.. Consumer Behavior Tenth Edition. Prentice-Hall; 2010.
70. Shafiee-Jood, Cai. Reducing food loss and waste to enhance food security and environmental sustainability. Environmental Science & Technology. 2016;50(16):8432-8443.
71. Shukla, Jharkharia. Agri-fresh produce supply chain management: A state-of-the-art literature review. International Journal of Operations and Production Management. 2013;33(2):114-158.
72. Subrahmanyam KV. Linkages Between Farm and Non Farm Sector: Role of Processing of Horticultural Products. Mumbai, India: National Bank for Agriculture and Rural Development; 2000.
73. Sudharshan, Anand, Sudulaimuttu D. Marketing and post-harvest losses in fruits: Its implications on availability & economy-A study on pomegranate in Karnataka. International Journal of Management and Social Sciences Research. 2013;2(7):34-43.
74. Tessema. Effect of Post Harvest Treatments on Storage Behavior and Quality of Tomato Fruits. World Journal of Agricultural Sciences. 2013;9(1):29-37.
75. Thyberg, Tonjes. Drivers of food waste and their implications for sustainable policy development. Resources, Conservation and Recycling. 2016;106:110-123.
76. Tolani, Hussain. Strategic change in model of fruit and vegetables supply chain. Global Journal of Management and Business Studies. 2013;3(9):965-970.

77. Valdes, Ramos, Beltrán, Jiménez, Garrigós. State of the art of antimicrobial edible coatings for food packaging applications. *Coatings*. 2017;7(4):56.
78. Verghese, Lewis, Lockrey, Williams. Packaging's role in minimizing food loss and waste across the supply chain. *Packaging Technology and Science*. 2015;28(7):603-620.
79. Waitt, Phillips,. Food waste and domestic refrigeration: A visceral and material approach. *Social & Cultural Geography*. 2016;17(3):359-379.
80. Watkins, Ekman. Storage technologies: Temperature interactions and effects on quality of horticultural products. In *V International Postharvest Symposium*. 2004;682:1527-1534.
81. Wheeler, Von Braun. Climate change impacts on global food security. *Science*. 2013;341(6145):508-513.
82. WILLS. Postharvest. An introduction to the physiology & handling of fruit, vegetables & ornamentals. 1998;33-46.
83. Wilson, Clarke,. Food safety and traceability in the agricultural supply chain: Using the Internet to deliver traceability. *Supply Chain Management: An International Journal*. 1998;3(3):127-133.
84. Winkworth-Smith, Foster, Morgan. The Impact of Reducing Food Loss in the Global Cold Chain, the University of Nottingham, Final Report; 2015.
85. WRAP. Helping Consumers Reduce Fruit and Vegetable Waste; 2008.
86. WRAP. Household Food and Drink Waste in the United Kingdom 2012–final Report; 2013.
87. Yahia, García-Solís, Celis MEM.. Contribution of fruits and vegetables to human nutrition and health. In *Postharvest physiology and biochemistry of fruits and vegetables* . Woodhead Publishing. 2019;19-45.
88. Barau B, Adelusi SM, Waiwada AA, Sani MG. Causes and management of postharvest losses in Nigeria: An overview. *Journal of Postharvest Technology*. 2023; 11(3):86-99.
89. Dsouza A, Mishra AK, Webster S. Vertical coordination and post-harvest losses: Implications on food loss Applied Economic Perspectives and Policy. 2023 Mar;45(1):460-86.

© 2024 Bisht and Singh; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/112719>