



Phytochemical Screening of Seaweed Flour (*Eucheuma cottonii*) Using Various Organic Solvents and Its Application in Tilapia Feed (*Oreochromis niloticus*)

Salnida Yuniarti Lumbessy ^{a*}, Zaenal Abidin ^a,
Dewi Putri Lestari ^a, Wastu Ayu Diamahesa ^a
and Thoy Batun Citra Rahmadani ^a

^a Aquaculture Study Program, Department of Fisheries and Marine Sciences, Faculty of Agriculture, University of Mataram, Jalan Pendidikan Mataram 83115, Nusa Tenggara Barat, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajfar/2024/v26i11833>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125979>

Original Research Article

Received: 26/08/2024

Accepted: 30/10/2024

Published: 05/11/2024

ABSTRACT

This study aims to qualitatively analyze the bioactive compounds in *Eucheuma cottonii* seaweed flour extract and its application in commercial feed for tilapia (*Oreochromis niloticus*). The methodology employed is experimental and consists of two phases. Phase 1 involves the extraction of *E. cottonii* flour using three different organic solvents with varying polarity (ethanol, ethyl acetate, and a combination of ethanol + ethyl acetate). The testing parameters include qualitative

*Corresponding author: Email: salnidayuniarti@unram.ac.id;

Cite as: Lumbessy, Salnida Yuniarti, Zaenal Abidin, Dewi Putri Lestari, Wastu Ayu Diamahesa, and Thoy Batun Citra Rahmadani. 2024. "Phytochemical Screening of Seaweed Flour (*Eucheuma Cottonii*) Using Various Organic Solvents and Its Application in Tilapia Feed (*Oreochromis Niloticus*)". *Asian Journal of Fisheries and Aquatic Research* 26 (11):100-108. <https://doi.org/10.9734/ajfar/2024/v26i11833>.

phytochemical tests for alkaloids, steroids/triterpenoids, saponins, flavonoids, and tannins. Phase 2 consists of the application of the extract to tilapia over a 30-day rearing period with four treatments: control/commercial feed (P0), commercial feed + ethanol extract (P1), commercial feed + ethyl acetate extract (P2), and commercial feed + combined ethanol and ethyl acetate extract (P3). The parameters tested include absolute weight, specific growth rate, survival rate, and water quality. Phytochemical and water quality data are presented in table format. Meanwhile, growth and survival data are analyzed using ANOVA, followed by Duncan's test for significant differences. The results of the researches carried out, it can be concluded that the phytochemical analysis of *E. cottonii* extract in ethanol and the ethanol + ethyl acetate combination yielded similar compounds: alkaloids, steroids/triterpenoids, and flavonoids. The ethyl acetate extract contained triterpenoids and flavonoids. Both saponins and tannins returned negative results across all organic solvent treatments. All treatments with the addition of *E. cottonii* seaweed extract in various organic solvents demonstrated better growth in tilapia compared to the control treatment.

Keywords: *Eucheuma cottonii*; extract; phytochemistry; tilapia; organic solvents.

1. INTRODUCTION

Seaweed contains secondary metabolites that have the potential to produce diverse bioactive metabolites with a wide range of activities, including antibacterial (Vallinayagam et al., 2009), antiviral (Serke djieva 2004), antifungal (Aliya and Shamaeel 1999), and cytotoxic (Vinayak et al., 2010) properties. According to Hashimoto (1997), the chemical compounds resulting from the secondary metabolism of seaweed that act as antibacterial agents include fatty acids, terpenoids, bromophenols, and tannins. Darusman et al. (1995) also explained that marine-derived natural products are a result of secondary metabolites from various groups, including alkaloids, terpenoids, and flavonoids, and can also directly originate from primary metabolite compounds such as dipeptides.

The presence of bioactive compounds in seaweed can be assessed using methods that provide information about these compounds. One such method is phytochemical screening (Setyowati et al., 2014). Phytochemical analysis involves a series of processes or techniques to identify and characterize the types of bioactive compounds in seaweed, such as alkaloids, flavonoids, and tannins, qualitatively (Soamole et al., 2018). Bioactive compounds resulting from secondary metabolism can be obtained through extraction processes. The solvents used in extraction can be polar, semi-polar, or non-polar (Agusman et al., 2022).

Previous studies have shown that the ethanol fraction of *E. spinosum* contains triterpenoids, alkaloids, and flavonoids (Sari et al., 2015). Research by Hanapi et al. (2013) demonstrated that the active compounds in the methanol

extract of *E. spinosum* include flavonoids, triterpenoids, and ascorbic acid. Similar findings were reported by (Yainahu et al., 2022), indicating that phytochemical screening of both fresh and dried *E. spinosum* extracts contained flavonoids, alkaloids, terpenoids, saponins, and tannins. Meanwhile, phytochemical tests of the methanol extract from *E. cottonii* qualitatively revealed the presence of alkaloids, flavonoids, steroids/terpenoids, and saponins. The n-hexane fraction contained steroids/terpenoids and saponins, while the ethyl acetate fraction contained alkaloids, flavonoids, and steroids/terpenoids, and the butanol fraction contained alkaloids (Agusman et al., 2022). The addition of *K. alvarezii* ethanol extract 1:5 to the feed showed the best results because it could increase the growth and utilization of tilapia feed with a survival rate of tilapia of 83% (Lumbessy et al., 2024). Meanwhile, the addition of 1:5 ethyl acetate extract can increase growth and better feed utilization in tilapia with a survival rate of tilapia of 90% (Zahara et al., 2024).

Based on the description above, this study aims to qualitatively analyze the bioactive compounds in *Eucheuma cottonii* seaweed flour extract and its application in commercial feed for tilapia (*Oreochromis niloticus*).

2. MATERIALS AND METHODS

2.1 Materials and Methodology

The materials used in the extraction process include *E. cottonii* flour, ethanol, and ethyl acetate. The materials for phytochemical testing include *E. cottonii* flour extract, Mg-HCl (magnesium chloride), NaOH (sodium hydroxide), H₂SO₄ (sulfuric acid), Mayer's

reagent, Dragendorff's reagent, Wagner's reagent, Liebermann-Burchard reagent, distilled water, and FeCl_3 . All chemicals used are classified as PA (Pro-analysis). The materials for the feed test include Hi Pro Vite 783 commercial feed and tilapia.

This research was carried out in October – December at the Laboratory of Fish Production and Reproduction, University of Mataram. This study employs an experimental method consisting of two phases:

Phase 1. Preparation of crude extract and identification of phytochemical compounds: the treatments involve organic solvents with varying polarity (ethanol, ethyl acetate, and a combination of ethanol + ethyl acetate). The parameter tested is the qualitative identification of phytochemical compounds in *E. cottonii* flour extract.

Phase 2. Application testing of the crude seaweed extract in feed: This phase evaluates the growth and survival of tilapia cultivated for 60 days. The treatments tested are the effects of adding *E. cottonii* extract with different solvents to commercial tilapia feed, as follows:

- P1: Commercial feed (control)
- P2: Commercial feed + ethanol extract
- P3: Commercial feed + ethyl acetate extract
- P4: Commercial feed + combination of ethanol and ethyl acetate extracts

The experimental design used in this phase was a Completely Randomized Design (CRD) consisting of four treatments, each repeated three times, resulting in a total of 12 treatments.

2.2 Preparation of Seaweed

The seaweed used was fresh seaweed obtained from cultivators. A total of 2 kg of fresh seaweed was air-dried for 14 days, depending on weather conditions, until fully dried. The dried seaweed was then cut into small pieces using scissors. The cut seaweed was ground using a blender and sieved to obtain flour.

2.3 Preparation of Seaweed Extraction

The extraction process of *E. cottonii* flour was conducted using a maceration method. Ground seaweed was weighed at 100 g and placed into three Erlenmeyer flasks, to which 96% ethanol, ethyl acetate, and a combination of ethanol + ethyl acetate was added according to the treatment used. The ratio of extract to solvent

used was 1:5 (m/v). The maceration time followed the guidelines from Purba et al. (2019), lasting 30 hours at 30°C. During maceration, the solution was stirred and shaken every 5 hours. The macerated flour was then filtered using Whatman No. 42 filter paper to produce a filtrate. The resulting filtrate was evaporated at 40°C using a vacuum rotary evaporator (Podungge et al., 2017).

2.4 Qualitative Analysis of Bioactive Compounds (Followed the Guidelines from Harborne (1996))

2.4.1 Flavonoids

An amount of 15 mg of extract was added to 10 mL of ethanol and homogenized. It was then boiled for 5 minutes and filtered. Then, 5 mL of the filtrate was mixed with 0.05 mg of powdered Mg and 1 mL of concentrated HCl and shaken vigorously. A positive test is indicated by the formation of red, yellow, or orange color.

2.4.2 Alkaloids

An amount of 50 mg of extract was mixed with 2 mL of chloroform and 2 mL of ammonia and then filtered. The filtrate was treated with 3-5 drops of concentrated H_2SO_4 and shaken to form two layers. The acidic fraction is collected. Mayer's and Dragendorff's reagents (4-5 drops each) were added. The formation of a precipitate indicates the presence of alkaloids, with Mayer's reagent producing a white precipitate and Dragendorff's reagent yielding a reddish-orange precipitate.

2.4.3 Steroids and terpenoids

A 50 mg extract was treated with 10 drops of concentrated CH_3COOH and 2 drops of concentrated H_2SO_4 . The solution was gently mixed and allowed to stand for a few minutes. Steroids produce a blue or green color, while terpenoids yield a red or purple color.

2.4.4 Saponins

An amount of 50 mg extract was mixed with 10 mL of water and shaken for 1 minute, then 2 drops of 1N HCl were added. If stable foam forms for approximately 7 minutes, the extract is considered positive for saponins.

2.4.5 Tannins

An amount of 0.5 g of *E. spinosum* extract was infused with 10 mL of distilled water and then

filtered. The resulting filtrate was diluted with water until colorless. Then, 2 mL of this solution was taken, and 1-2 drops of 1% ferric (III) chloride reagent were added. The development of blue or dark green color indicates the presence of tannins.

2.5 Testing of the *E. cottonii* Extract in Tilapia Commercial Feed

The application of the extract to tilapia feed involved spraying the seaweed extract at a dosage of 2 g/kg of feed according to the treatment. Before spraying, the extract was diluted with distilled water, and after even application, the feed was air-dried. The feed was then provided to the tilapia in a 45 L container for 30 days, with a stocking density of 10 fish per container. Throughout the cultivation period, growth and water quality were measured every 10 days, along with monitoring fish mortality and performing siphoning. The parameters analyzed include absolute weight, specific growth rate (SGR), survival rate (SR), and water quality parameters such as temperature, pH, and dissolved oxygen (DO).

2.6 Data Analysis

The results of the phytochemical tests and water quality are presented in table format. Meanwhile, the fish growth and survival rate data were analyzed using ANOVA, followed by Duncan's test if significant differences were observed.

3. RESULTS AND DISCUSSION

3.1 Phytochemical Testing

The results of the phytochemical testing to identify secondary metabolite compounds qualitatively in *E. cottonii* seaweed using ethanol, ethyl acetate, and a combination of ethanol and ethyl acetate are presented in Table 1. A positive result (+) indicates the presence of specific compound groups, while a negative result (-) indicates their absence.

Table 1 shows that the extracts of *E. cottonii* in both ethanol and the combination of ethanol + ethyl acetate yielded the same positive results for the same compound groups, namely alkaloids, steroids/triterpenoids, and flavonoids. In contrast, the ethyl acetate extract contained triterpenoids and flavonoids. The groups of compounds saponins and tannins showed negative results for all treatments with the organic solvents.

The alkaloid test for *E. cottonii* in ethyl acetate showed no precipitate after reacting with Mayer's and Dragendorff's reagents. In contrast, both ethanol and the combination of ethanol + ethyl acetate resulted in precipitates when treated with these reagents. According to Purba (2001), alkaloids contain nitrogen in their cyclic structure and have varying functional groups, such as amines, amides, phenols, and methoxy groups, which makes alkaloids semi-polar. This semi-polar characteristic allows alkaloids to be more soluble in semi-polar solvents and dissolve in both semi-polar and polar solvents (Harborne et al., 1996).

The steroid and terpenoid tests using the Liebermann-Bouchard method showed positive results, with color changes observed in the ethanol and combination of ethanol + ethyl acetate extracts of *E. cottonii*. The ethyl acetate extract showed a positive result for steroids but negative for terpenoids due to the absence of color change. Saidi et al. (2018) state that terpenoid and steroid compounds are soluble in non-polar to semi-polar solvents. Some triterpenoid compounds may have cyclic alcohol structures, contributing to their semi-polar nature (Titis et al., 2013). Steroids can exist in glycoside forms. Glycosides, which consist of sugar and aglycone, have a polar sugar component that allows them to dissolve in polar solvents. Hence, steroids were detected in the methanol extract.



















The bioactive compound test for saponins yielded negative results across all organic solvent treatments. The saponin test involves the hydrolysis of saponins in water. Saponins are triterpenoid glycosides that tend to be polar due to their glycosidic bonds (Sangi et al., 2008). According to Robinson (Robinson et al., 1995), saponin compounds possess both polar and non-polar groups that exhibit surface-active properties. When saponins are shaken with water, hydrolysis occurs, forming micelles. The structure of these micelles causes the polar groups to face outward while the non-polar groups face inward, resulting in a foam-like appearance.







The identification of flavonoids in the *E. cottonii* flour extract, treated with magnesium powder (Mg) and 2 N hydrochloric acid (HCl), showed positive results across all organic solvent treatments due to the color change to red. This is attributed to the reduction reaction of flavonoids induced by hydrochloric acid and magnesium (Simaremare 2014). According to Sjahid (2008),

flavonoids belong to the phenolic group and are polar compounds due to the presence of several untraceable hydroxyl groups or sugars, allowing them to dissolve in polar solvents such as methanol, ethanol, butanol, acetone, and

dimethyl sulfoxide. The presence of flavonoids in all solvent treatments indicates that flavonoid compounds share similar polarity with ethanol and ethyl acetate, thus allowing the formation of flavonoids in these solutions.

Table 1. Phytochemical Test Results of *Eucheuma cottonii* Extract with Various Solvents

Phytochemical Testing	Standard Color (+)	Solvents		
		Ethanol	Ethyl Acetate	Ethanol + Ethyl Acetate
A. Alkaloids				
a. Dragendrof's Reagent	Orange-red precipitate	 (+)	 (-)	 (+)
b. Meyer's Reagent	A white or yellowish precipitate	 (+)	 (-)	 (+)
c. Wanger's Reagent	Brown precipitate	 (+)	 (-)	 (+)
B. Triterpenoids and steroid	Greenish-blue and blue color (steroid)	 (+)	 (-)	 (+)
	Brown, red, orange, or purple ring (triterpenoids)	 (+)	 (+)	 (+)
C. Saponins	Stable foam for 3-5 minutes	 (-)	 (-)	 (-)

Phytochemical Testing	Standard Color (+)	Solvents		
		Ethanol	Ethyl Acetate	Ethanol + Ethyl Acetate
D. Flavonoids	Color change to red or yellow	 (+)	 (+)	 (+)
E. Tannins	Color change to bluish-black or green	 (-)	 (-)	 (-)

3.2 Testing of the *E. cottonii* Extract in Tilapia Commercial Feed

The ANOVA results for the application of the extract in commercial feed for tilapia during the 30-day cultivation period indicated that the addition of commercial feed with extracts of the seaweed *E. cottonii* using various organic solvents significantly affected the absolute weight and specific growth rate of tilapia, but it did not have a significant effect on the survival rate of the fish. Duncan's post hoc test revealed that all treatment groups that received commercial feed supplemented with extracts of *E. cottonii* (P1, P2, and P3) resulted in higher absolute weight and specific growth rates compared to the control group (P0) (Figs. 1–3).

The better growth of tilapia in all feed treatments supplemented with *E. cottonii* extracts using

various solvents (P1, P2, and P3) is believed to be related to the presence of phytochemical compounds, specifically flavonoids and terpenoids. According to Munglue (2014), flavonoids and terpenoids have the potential to enhance growth and feed utilization in Nile tilapia. Phytochemical compounds such as flavonoids and terpenoids can stimulate specific enzymes in the digestive tract, such as trypsin, amylase, and cytochrome c oxidase, which help improve digestion and nutrient absorption in fish. Mansour et al. (2022) note that feed rich in flavonoids can significantly enhance weight gain, specific growth rate, and feed ratio in tilapia up to a level of 5 g/kg. These bioactive compounds directly increase digestive enzyme activity, positively affecting fish growth performance. Furthermore, Zhai and Liu (2013) state that flavonoid bioactive compounds can enhance animal growth.




Fig. 1. Absolute weight of tilapia (*O. niloticus*)




Fig. 2. Specific growth rate of tilapia (*O. niloticus*)




Fig. 3. Survival rate of tilapia (*O. niloticus*)

Meanwhile, all feed treatments in this study did not affect the survival rate of the tilapia. This indicates that the addition of *E. cottonii* extracts using various solvents in commercial feed is not harmful and does not have toxic effects

compared to the control treatment. This is supported by temperature, pH, and dissolved oxygen (DO) measurements in the cultivation water, which remained within the optimal range for the survival of tilapia (Table 2).

Table 2. Water quality parameters

Parameter	Treatments				Source Amalia (2018)
	P0	P1	P2	P3	
Suhu (°C)	26.9 – 28.6	26.6 - 29.4	26.9 - 28.3	26.9 - 28.5	25 - 30°C
DO (mg/L)	4.9 - 7.9	6 - 6.5	6 - 8	5.7 – 7.5	> 5 mg/L
pH	6.1 - 7.8	6.5 - 7.5	6.8 - 7.4	6.5 - 7.4	5 - 11

4. CONCLUSION

The phytochemical test results of *E. cottonii* extract using ethanol and a combination of ethanol + ethyl acetate produced the same compounds: alkaloids, steroids/triterpenoids, and flavonoids. Meanwhile, the ethyl acetate solvent contained triterpenoids and flavonoids. The groups of compounds saponins and tannins yielded negative results across all organic solvent treatments. All treatments involving the addition of commercial feed with extracts of the seaweed *E. cottonii* using various organic solvents significantly improved the growth of tilapia compared to the control treatment.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Agusman, I., Diharmi, A., & Sari, N. I. (2022). Identification of bioactive compounds in red seaweed fraction (*Eucheuma cottonii*). *Aquatic Sciences Journal*, 9(2), 60-64.
- Aliya, R., & Shamaeel, M. (1999). Phytochemical evaluation of four coenocytic green seaweed from the coast of Karachi. *Pakistan Journal of Marine Biology*, 5, 65-76.
- Amalia, R., Amrullah, & Suriati. (2018). Management feeding in tilapia (*Oreochromis niloticus*) farming. In *Proceedings of the 2018 National Seminar on Multidisciplinary Synergy of Science and Technology* (pp. 252-257).
- Darusman, L. K., Sayuti, D., Komar, & Pamungkas, J. (1995). Extraction of bioactive components as drug ingredients from corals, coral flowers and algae in the waters of Pari Island, Seribu Islands. Phase II: Fractionation and Bioassay. *Chemical Bulletin*, 10, 18-27.
- Hanapi, A., Fasya, G., Mardiyah, U., & Mittahurrahmah. (2013). Activity test antioxidant and antibacterial methanol extract of red algae *Eucheuma spinosum* from Wongsorejo waters, Banyuwangi. *J. Alchemy*, 2(2), 126-137.
- Harborne, J. B. (1996). *Phytochemical methods: A guide to modern ways of analyzing plants* (2nd ed.). Bandung: ITB.
- Hashimoto, Y. (1979). *Marine toxins and other bioactive marine metabolites*. Tokyo: Japan Scientific Soc. Press.
- Lumbessy, S. Y., Wahyuningsy, T., & Abidin, Z. (2024). Feeding tilapia (*Oreochromis niloticus*) with the addition of seaweed flour ethanol extract (*Kappaphycus alvarezii*) in a comparison of different solvent concentrations. *Jurnal Perikanan Pantura (JPP)*, 7(2), 513-516.
- Mansour, A. T., Mahboub, H. H., Elshopakey, G. E., Aziz, E. K., Alhajji, A. H. M., Rayan, G., Ghazzawy, H. S., & El-Houseiny, W. (2022). Physiological performance, antioxidant and immune status, columnaris resistance, and growth of tilapia that received *Alchemilla vulgaris*-supplemented diets. *Antioxidants*, 11(8). <https://doi.org/10.3390/antiox11081494>
- Munglue, P. (2014). Effects of dietary *Nelumbo nucifera* (Lotus) peduncle extract on growth performance of tilapia (*Oreochromis niloticus*), 279–282.
- Podungge, A., Damongilala, L. J., & Mewengkang, H. W. (2017). Antioxidant content in *Eucheuma spinosum* seaweed extracted with methanol and ethanol. *Fishery Product Technology Media*, 6(1), 1-5.
- Purba, N. E., Suhendra, L., & Wartini, N. M. (2019). Effect of temperature and extraction time by maceration method on the characteristics of dye from red algae extract (*Gracilaria sp.*). *Journal of Agro-Industrial Engineering and Management*, 7(4), 551-560.
- Purba, R. D. (2001). Analysis of alkaloid composition of handeuleun leaves (*Graptophyllum pictum* (Linn, Griff)) cultivated with different nitrogen levels. (Thesis). Bogor Agricultural University, Bogor.
- Robinson, T. (1995). *Organic compound content of higher plants* (translated by K. Padmawinata). Bandung: ITB.
- Saidi, N., Ginting, B. I., Murniana, & Mustanir. (2018). *Secondary metabolite analysis* (1st ed.). Banda Aceh: Syiah Kuala University Press.
- Sangi, M., Runtuwene, M. R. J., Simbala, H. E. I., & Makang, V. M. A. (2008). Phytochemical analysis of medicinal plants

- in North Minahasa regency. *Chemistry Progress*, 1(1), 47-53.
- Sari, B. L., Susanti, N., & Sutanto. (2015). Phytochemical screening and antioxidant activity of red algae ethanol fraction *Eucheuma spinosum*. *Pharmacy Study Program, Faculty of Mathematics and Natural Sciences, Pakuan University. Pharm Sci Res*, 2(2), 59-67.
- Serkedjieva, J. (2004). Antiviral activity of the red marine alga *Ceramium rubrum*. *Phytotherapy Research*, 18(6), 480–483.
- Setyowati, W. A. E., Ariani, S. R. D., Ashadi, M. B., & Rahmawati, C. P. (2014). Phytochemical screening and identification of main components of methanol extract of durian peel (*Durio zibethinus* Murr.) Petruk variety. In *National Seminar on Chemistry and Chemical Education VI* (pp. 271-280).
- Simaremare, S. A. (2014). Screening phytochemistry of ethanol extract of itchy leaves (*Laportea decumana* Roxb.). *Jurnal Pharmacy*, 11(1), 1-10.
- Sjahid, L. R. (2008). Isolation and identification of flavonoids from dewandaru leaves (*Eugenia uniflora* L.) (Thesis). Surakarta: Muhammadiyah University of Surakarta.
- Soamole, H., Sanger, G., Harikedua, S., Dotulong, V., Mewengkang, H., & Montolalu, R. (2018). Phytochemical content ethanol extract of fresh seaweed (*Turbinaria* sp., *Gracilaria* sp., and *Halimeda macroloba*). *Journal of Fisheries Product Technology Media*, 6(3), 287-291.
- Titis, M. B., Fachriyah, E., & Kusriani, D. (2013). Isolation, identification and activity test of alkaloid compounds of binahong leaves (*Anredera cordifolia* (Tenore) Steenis). *Chem. Info*, 1(1), 196-201.
- Vallinayagam, K., Rajasekaran, A., Kannan, R., Thirumaran, G., & Anantharaman, P. (2009). Antibacterial activity of seaweeds of Pudumadam coast. *Global Journal of Pharmacology*, 3(1), 50-52.
- Vinayak, R. C., Sabu, A. S., & Chatterji, A. (2010). Bio-prospecting of a few brown seaweeds for their cytotoxic and antioxidant activities. *Evidence-Based Complementary and Alternative Medicine*, 1-9.
- Yainahu, J. L., Mile, S., Pratama, & Suherman. (2022). Yield analysis and phytochemical screening of fresh and dried red seaweed (*Eucheuma spinosum*) extracts. *Jambura Fish Processing Journal*, 5(2), 126–132.
- Zahara, A. Z., Abidin, Z., & Lumbessy, S. Y. (2024). Addition of ethyl acetate extract of *Kappaphycus alvarezii* seaweed flour in commercial feed for tilapia (*Oreochromis niloticus*). *Jurnal Vokasi Ilmu-Ilmu Perikanan (JVIP)*, 5(1), 01-08.
- Zhai, S. W., & Liu, S. L. (2013). Effects of dietary quercetin on growth performance, serum lipid levels and body composition of tilapia (*Oreochromis niloticus*). *Italian Journal of Animal Science*, 12(4), 523–527.
<https://doi.org/10.4081/ijas.2013.e85>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125979>