



Sonographic Assessment of Maternal Portal Vein Diameter in Healthy Pregnancy in South-South Nigeria

Peter Chibuzor Oriji^a, Enefia Kelvin Kiridi^{b,c*},
Emily Gabriel Enefia Kiridi^d, Obiora Chibundu^{e,f},
Johnpatrick Uchenna Ugwoegbu^g,
Akaninyene Esemé Ubom^{h,i}, Panebi Yao Bosrotsi^j,
Abednigo Ojanerohan Addah^k, Isaac Joel Abasi^k
and Adedotun Daniel Adesina^{l,m}

^a Department of Obstetrics and Gynaecology, Federal Medical Centre, Yenagoa, Bayelsa State, Nigeria.

^b Department of Radiology, Niger Delta University Teaching Hospital, Okolobiri, Bayelsa State, Nigeria.

^c Silhouette Radiodiagnostic Consultants, Yenagoa, Bayelsa State, Nigeria.

^d Department of Physiology, Niger Delta University, Wilberforce Island, Amassoma, Bayelsa State, Nigeria.

^e Department of Obstetrics and Gynaecology, Asokoro District Hospital, Abuja, Nigeria.

^f Department of Obstetrics and Gynaecology, Nile University of Nigeria, Abuja, Nigeria.

^g Department of Radiology, Federal Medical Centre, Owerri, Imo State, Nigeria.

^h Department of Obstetrics, Gynaecology and Perinatology, Obafemi Awolowo University Teaching Hospitals Complex, Ile-Ife, Osun State, Nigeria.

ⁱ International Federation of Gynaecology and Obstetrics (FIGO), Committee on Childbirth and Postpartum Haemorrhage, Nigeria.

^j Department of Obstetrics and Gynaecology, Diète Koki Memorial Hospital, Yenagoa, Bayelsa State, Nigeria.

^k Department of Obstetrics and Gynaecology, Niger Delta University Teaching Hospital, Okolobiri, Bayelsa State, Nigeria.

^l Department of Medical Services, Nigerian Law School, Yenagoa Campus, Yenagoa, Bayelsa State, Nigeria.

^m Oasis Public Health Consulting Ltd, Yenagoa, Bayelsa State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author PCO designed the study, collated data, wrote the introduction, methodology and discussion. Author EKK conceptualised the study, carried out obstetric ultrasound scans and collected data. Authors EGEK and OC managed literature search. Author JUJ did obstetric ultrasound scans. Author AEU wrote the abstract. Authors PCO, PYB, AOA and IJA recruited patients from the antenatal clinics. Author ADA analysed data and wrote the results. All the authors read and approved the final manuscript.

*Corresponding author: E-mail: kiridienefia@gmail.com;

Article Information

DOI: 10.9734/AJMAH/2023/v21i2794

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96489>

Original Research Article

Received: 01/01/2023

Accepted: 06/02/2023

Published: 09/02/2023

ABSTRACT

Background: The portal vein is one of the main sources of blood supply of the liver. About three quarters of the liver blood flow is from the portal vein.

Objectives: To assess the mean normal portal vein diameter in healthy pregnant women, and correlate our observation with maternal age, parity and other baseline characteristics.

Subjects and Methods: This descriptive, cross-sectional study was conducted at the Obstetrics and Radiology Units of the two tertiary health facilities, one secondary facility and one radio-diagnostic facility in Bayelsa State, South-South Nigeria, between April, 2022 and December, 2022. Data analysis was done using Statistical Product and Service Solutions for Windows® version 25, SPSS Inc.; Chicago, USA. Descriptive statistics (mean, standard deviation, frequency, and percentages) and Pearson product moment correlation were used for the analysis. The level of significance was set at $p < 0.05$.

Results: Mean portal vein diameter was 10.43 ± 1.58 mm, and ranged from 7.5 mm to 14.0 mm. Maternal age ($r = 0.51$; $p = 0.001$) and parity ($r = 0.47$; $p = 0.001$) correlated fairly strong and significantly with portal vein diameter. Gestational age ($r = 0.21$; $p = 0.001$) and estimated foetal weight ($r = 0.16$; $p = 0.001$) were significantly related to the portal vein diameter.

Conclusion: This research has established baseline values for normal range of PV diameter in healthy pregnant women in the South-South region of Nigeria, and also revealed significant correlation of PV diameter with age, parity, gestational age and estimated foetal weight.

Keywords: Portal vein diameter; liver; pregnant women; maternal age; parity; gestational age.

1. INTRODUCTION

The portal vein (PV) and hepatic artery are the main sources of blood supply of the liver. About three quarters of the liver blood flow is from the portal vein, while the remaining one quarter comes from the hepatic artery [1]. The superior mesenteric vein and splenic vein meet at the level of the second lumbar vertebra, behind the pancreatic neck, to form the portal vein (PV).[1] An important tool for making diagnosis of portal hypertension is the sonographic measurement of the portal vein diameter. The complex interaction between the liver and the portal vein supports the body's homeostasis.[1]

There is a paucity of published literature on portal vein diameter in pregnancy. What abounds in the literature is portal vein diameter in non-pregnant individuals. Portal hypertension is a

major abnormality of the portal venous system. It usually occurs due to an increase in portal venous pressure, which subsequently leads to resistance of blood flow through the portal vein into the hepatic circulation [2–5]. Liver disease in pregnancy is uncommon, but could be a serious illness when it occurs.

Portal hypertension leads to splenomegaly, portal vein enlargement, and the development of portal systemic collaterals at various sites. Due to the fact that it is the most frequent complication and the main cause of death among patients with chronic liver disease, it results in significant mortality and morbidity [2,6,7]. The normal portal vein diameter varies between 7 mm and 15 mm.[8] Some authors have reported 13 mm as the upper limit of the portal vein diameter, and a value greater than that is suggestive of portal hypertension [4,5,9,10]. The

normal portal venous pressure varies between 5 mmHg and 10 mmHg [8]. A portal venous pressure of greater than 15 mmHg (30 cmH₂O) may be suggestive portal hypertension [8].

The causes of portal hypertension could either be pre-hepatic, hepatic, or post-hepatic. The most common causes of portal hypertension are cirrhosis in developed countries, [6,10,11] schistosomiasis in endemic areas [11,12] and hepatic vascular abnormalities [5]. These conditions and other risk factors like hepatitis, alcohol abuse cause scarring of the liver, which in turn causes cirrhosis [11]. Diagnostic imaging techniques such as arteriography, splenoportography, and portal venography have been used to assess patients suspected of having portal hypertension; however, these procedures are invasive, costly, time-consuming, and risky for the patient. Computed tomography and magnetic resonance imaging, on the other hand, have the advantage of providing better cross-sectional images, but they are both costly, and the former exposes the patient to high doses of ionizing radiation [13,14].

Sonography is a good diagnostic tool that plays a significant role in the diagnosis and follow-up of patients with portal hypertension due to its accessibility, non-invasive nature, mobility, low cost, and capacity to complete tasks quickly. Sonography also uses non-ionizing radiation. Therefore, the objective of this study was to assess the mean normal portal vein diameter in healthy pregnant women, and correlate our observation with maternal age, parity and other baseline characteristics.

2. MATERIALS AND METHODS

Study design and setting: This descriptive, cross-sectional study recruited and enrolled women with normal pregnancy in their second and third trimesters at the Radiology and Obstetrics units of the Niger Delta University Teaching Hospital, Okolobiri, Federal Medical Centre, Yenagoa, Silhouette Radiodiagnostic Consultants, Yenagoa and Diète Koki Memorial Hospital, Yenagoa, all in Bayelsa State, Nigeria. The study was conducted between April, 2022 and December, 2022.

Sample size calculation: This was calculated using the formula: $n = Z\alpha^2 \times \sigma^2 / \delta^2$ [15,16].

Where: $Z\alpha$ = 95% CI, which is 1.96, σ = mean of 10.65 mm from a previous study [11] δ = level of precision for our study ($\sigma/\sqrt{63}$).

Calculation:

$$\begin{aligned}n &= (1.96)^2 \times 10.65^2 / \sigma/\sqrt{63} \\n &= 3.8416 \times 113.42 / 1.34 \\n &= 432.81 / 1.34 \\n &= 322.993 \\n &= 323\end{aligned}$$

Considering attrition of 5% (16.15), n was adjusted to 339.

For this study, 339 consecutive pregnant women were enrolled. The study included consecutive patients who visited our Obstetric Unit.

Inclusion criteria: Women with normal singleton pregnancies.

Exclusion criteria: Women with liver disease and other medical conditions in pregnancy.

After counseling, written informed consent to participate was obtained from all the women enrolled in the study. Baseline information and any presenting complaints were obtained. With the patient standing on the Frankfort plane, the height of the patient was measured using a wall-mounted stadiometer. A weighing scale was used to determine weight. Patients were asked to take off their bulky outerwear and shoes and stand in the middle of the scale to evenly distribute their weight across both feet. Body mass index (BMI) was determined as the product of height (m) squared and weight (kg). The last normal menstrual period, which corresponded with their first trimester ultrasound scan, was used to determine the gestational age. Urinalysis, liver function tests and serum electrolytes, urea and creatinine, were done for the women, and if these were normal, they were then referred to the Radiology Units of the study centres for ultrasound scan.

Procedure: All ultrasound examinations were performed transabdominally by consultant radiologists, using a 2012 Philips HD11 device with a 3.5 MHz curvilinear probe. Before data collection commenced, the consultant radiologists discussed, assessed for interobserver and interobserver variability and reliability, and reached an agreement on the standard operative procedure of ultrasonography to ensure data quality. After an overnight fast, the individuals were placed in the supine and right anterior oblique positions for the ultrasound examination. When the main portal vein could be seen best, subjects were exposed from the xiphisternum to the pelvic brim, ultrasound gel

was applied to the right upper quadrants of the abdomen, and the transducer was placed in the epigastrium in both the transverse and longitudinal planes. Measurements were taken at the location where the portal vein crosses anterior to the inferior vena cava, with the calipers placed between the inner margins of the echogenic walls of the vessel at the location where the portal vein crosses prior to the inferior vena cava (Fig. 1).

Data analysis: Data capture sheet was used to record all the measurements obtained. Data analysis was done using Statistical Product and Service Solutions for Windows® version 25, SPSS Inc.; Chicago, USA. Descriptive statistics (mean, standard deviation, frequency, and percentages) and Pearson product moment correlation were used for the analysis. Interobserver and intraobserver variations were calculated with the use of the intraclass correlation coefficient (ICC) and documented. Statistical significance was considered at $P < 0.05$.

3. RESULTS

3.1 Maternal Baseline Characteristics

There were 339 pregnant women who participated in this study. Their mean age was 30.3 ± 5.1 years. Majority (60.2%) of the participants were in the fourth decade of life, while slightly above a third (36.6%) were aged between 20 and 29 years. The mean weight, height and body mass index of the pregnant women were 69.2 ± 11.3 kg, 1.62 ± 0.04 m and 26.3 ± 4.6 kg/m², respectively. One hundred and fifty-nine (46.9%) pregnant women were overweight, 136 (40.1%) had normal weight and 44 (13.0%) were obese. Half (50.1%) of the participants were multiparous/grand-multiparous, while 14.7% were primiparous. Median parity was 1, with a range between 0 and 5 (Table 1).

3.2 Relationship between Portal vein Diameter and Maternal/Foetal Characteristics

Portal vein diameter was 10.43 ± 1.58 mm (Fig. 2), with a range of 7.5 mm to 14.0 mm. Maternal age ($r = 0.51$; $p = 0.001$) and parity ($r = 0.47$; $p = 0.001$) correlated fairly strong and significantly with portal vein diameter. Maternal weight, height and body mass index in the pregnant population

did not show a significant relationship ($p > 0.05$) with portal vein diameter. Gestational age ($r = 0.21$; $p = 0.001$) and estimated foetal weight ($r = 0.16$; $p = 0.001$) were significantly related (weak) to the portal vein diameter (Table 2). Table 3 further demonstrated the relationship between portal vein diameter and age/parity of participants. There was a gradual increase in the portal vein diameter of participants aged 15 – 19 years to those aged ≥ 40 years. The difference in the mean portal vein diameter between these age groups was significant ($f\text{-stat} = 40.06$; $p = 0.001$). A similar trend was observed with parity (Table 3). Table 4 is a nomogram showing the value of portal vein diameter at different estimated gestational ages and estimated foetal weights, while Table 5 reports the interobserver and intraobserver intraclass correlation coefficient results.

4. DISCUSSION

This study was conducted among healthy pregnant women in Bayelsa State, South-South Nigeria. To the best of our knowledge, this is the first study to assess the portal vein diameter in healthy pregnant women. Therefore, the findings from this study will be compared with those conducted in healthy adults. Our study revealed a mean PV diameter of 10.43 ± 1.58 mm (with a range of 7.5 mm to 14.0 mm). This observation is in tandem with the mean PV diameter of 11.5 ± 0.15 mm reported by Anakwue et al., in South-East Nigeria,[2] 10.9 ± 0.81 mm reported by Usman et al., in Maiduguri, North-East Nigeria, [11] 11.0 ± 2.6 mm reported by Tasu et al., in France,[17] 11.7 ± 0.3 mm reported by Cosar et al. in Turkey[18] and 10.6 ± 1.8 mm reported by Geleto et al., in South-West Ethiopia.[8] A plausible reason for these similarities may be that ethnic and racial variations do not significantly influence PV dimensions. Since these studies were conducted in non-pregnant women, it is also possible that pregnancy does not affect the diameter of the portal vein. More researches are recommended to further assess portal vein diameter in pregnancy. Our value was, however, slightly higher than the 9.83 ± 0.95 mm reported by Akanni et al., in Parakou, Benin, [19] 9.6 ± 1.9 mm reported by Rokni-Yazdi and Sotouden in Iran [20] and 7.9 ± 2.0 mm reported by Hawaz et al., in Addis Ababa, Ethiopia [6]. This may be due to the difference in sample size and methodology of the various studies.




Fig. 1. Longitudinal view of the abdomen showing the levels of measurement of the portal vein (green dotted lines)

Table 1. Maternal baseline characteristics

Characteristics	Frequency, n = 339	Percent (%)
Age group (years)		
15 – 19	9	2.7
20 – 29	124	36.6
30 – 39	195	57.5
≥ 40	11	3.2
Age in years – Mean ± SD	30.3 ± 5.1	
Anthropometric measures		
Weight in kg – Mean ± SD	69.2 ± 11.3	
Height in metres – Mean ± SD	1.62 ± 0.04	
Body mass index in kg/m ² – Mean ± SD	26.3 ± 4.6	
Weight		
Normal weight	136	40.1
Overweight	159	46.9
Obese	44	13.0
Parity		
Nulliparity	119	35.1
Primiparous	50	14.7
Multiparous	145	42.8
Grand-multiparous	25	7.4
Parity – Median (range)	1 (0 – 5)	




Fig. 2. Histogram showing the distribution of portal vein diameter among pregnant women

Table 2. Correlation between portal vein diameter and age, parity and other baseline characteristics

Characteristics	Correlation coefficient - r (p-value)
Maternal features	
Age	0.51 (0.001*)
Parity	0.47 (0.001*)
Weight	0.05 (0.382)
Height	-0.09 (0.097)
Body mass index	0.08 (0.166)
Foetal features	
Gestational age	0.21 (0.001*)
Estimated foetal weight	0.16 (0.001*)

*Statistically significant

Table 3. Mean portal vein diameter in the different age-groups and parity

Characteristics	Frequency	Portal vein diameter Mean ± SD	F-stat (p-value)
Total population	339	10.43 ± 1.58	
Age group (years)			
15 – 19	9	8.87 ± 0.55	40.06 (0.001*)
20 – 29	124	9.54 ± 0.89	
30 – 39	195	10.11 ± 1.26	
≥ 40	11	10.99 ± 1.70	
Parity			
Nulliparity (0)	119	9.21 ± 0.76	75.67 (0.001*)
Primiparous (1)	50	9.35 ± 1.75	
Multiparous (2)	11	9.61 ± 1.08	
Multiparous (3)	56	10.76 ± 0.36	
Multiparous (4)	78	11.94 ± 1.12	
Grand-multiparous (≥5)	25	12.97 ± 1.51	

*Statistically significant

Table 4. Nomogram showing the relationship between PV diameter and gestational age/estimated foetal weight

Gestational age (weeks)	Percentiles in mm					Estimated foetal weight (kg)	Percentiles in mm				
	5th	10th	50th	90th	95th		5th	10th	50th	90th	95th
16	7.4	7.4	7.5	7.6	7.7	0.50	7.3	7.4	7.5	7.6	7.6
17	7.5	7.6	7.7	7.8	7.9	0.60	7.4	7.4	7.5	7.6	7.7
18	8.7	8.7	8.7	8.7	8.7	0.70	7.5	7.5	7.6	7.7	7.9
19	8.7	8.7	8.7	8.7	8.7	0.80	8.5	8.5	8.7	8.8	8.8
20	9.2	9.2	9.3	9.4	9.5	0.90	9.1	9.2	9.2	9.2	9.2
21	9.2	9.2	9.5	9.7	9.9	1.00	9.1	9.2	9.4	9.7	9.7
22	9.0	9.4	10.8	11.0	11.0	1.20	9.2	9.2	10.0	10.4	10.5
23	9.0	9.8	9.10	9.15	9.30	1.30	10.1	10.1	10.3	10.5	10.5
24	9.6	9.8	9.10	9.15	9.30	1.50	10.2	10.4	10.7	10.7	10.8
25	9.8	9.10	9.10	9.15	9.30	1.60	10.8	10.8	10.9	11.0	11.0
26	10.1	10.1	10.1	10.1	10.1	1.80	11.0	11.0	11.2	11.2	11.3
27	10.1	10.1	10.1	10.2	10.3	2.20	11.0	11.1	11.3	11.4	11.5
28	10.1	10.1	10.1	10.3	10.4	2.35	11.0	11.0	11.4	11.4	11.6
29	10.3	10.3	10.6	11.0	11.0	2.37	11.1	11.2	11.3	11.3	11.3
30	11.0	11.0	11.0	11.1	11.1	2.41	11.2	11.3	11.3	11.3	11.3
31	11.0	11.1	11.1	11.2	11.2	2.43	11.3	11.3	11.5	11.7	11.9
32	11.3	11.3	11.4	11.4	11.4	2.60	11.4	11.5	11.5	11.7	11.9
33	11.3	11.3	11.4	11.4	11.5	3.00	11.5	11.5	11.6	11.8	12.0
34	11.4	11.4	11.4	11.5	11.6	3.10	11.6	11.7	11.8	11.9	12.1
35	11.6	11.7	11.7	11.8	11.8	3.40	11.7	11.7	11.8	11.9	12.3
36	11.7	11.7	11.8	11.8	11.9	3.49	12.5	12.7	12.7	12.8	12.8
37	11.7	11.7	11.9	11.9	11.9	3.50	13.2	13.2	13.8	14.0	14.0
38	12.2	12.2	12.3	12.3	12.5	3.70	13.5	13.7	14.0	14.1	14.1
39	13.1	13.1	13.6	14.0	14.0	3.73	14.0	14.1	14.1	14.3	14.3
40	14.0	14.1	14.1	14.3	14.3						

Table 5. Interobserver and intraobserver intraclass correlation coefficient results

Ultrasound parameter	Intraclass correlation coefficient	
	Interobserver	Intraobserver
Portal vein diameter	0.99 (95% CI 0.53–0.99)	0.98 (95% CI 0.57–0.99)

In this study, maternal age correlated significantly ($r = 0.51$; $p = 0.001$) with portal vein diameter. The values of PV diameter increased with increase in the age of the women. This observation is in consonance with the reports of Anakwue et al. [2] Hawaz et al., [6] Usman et al., [11] Geleto et al. [8] and Shikha et al. [21] Weinreb et al. [4] Adeyekun and Tsebi, [14] and Cosar et al., [18] however, did not observe any correlation between PV diameter and age. This may have also resulted from the difference in sample size and methodology of the studies. In our study, there was no correlation between PV diameter and height, weight, and body mass index. This is in agreement with the findings of Usman et al. [11] in North-East Nigeria and Moriyasu et al. [22] who studied PV diameter in Caucasians. However, Gareeballah et al. [23] in

Sudan, and Saha et al. [24] and Lal et al. [25] both in India and Akanni et al. [19] in Parakou, Benin reported that weight and height were associated with PV diameter. The reason for this variable correlation with PV diameter in different studies is not readily understood. However, the relatively small sample size of these studies and the measuring techniques for PV diameter may have contributed. We could not compare our reports on parity, gestational age and estimated foetal weight with those of other authors, because studies on PV diameter assessment in healthy pregnancy did not exist in the literature as at the time of writing this discussion.

To reduce interobserver and intraobserver variability for the measurements of portal vein diameter in our study, the ICC was used. The

consistency of measurements of the same parameter is evaluated by the ICC [26]. The ICC considers the variance of all measurements and variations between the observers [26,27]. The normal range is 0 – 1, and a number greater than 0.8 suggests almost perfect agreement [28,29]. The inter- and intra-observer variance values in this research were 0.99 and 0.98, respectively, indicating an almost perfect agreement.

The strength of this research lies in the fact that it was a multicentre study where women with normal pregnancies were recruited. This removed confounding variables, like liver pathologies or other medical conditions in pregnancy, which would have affected the measurements of the portal vein diameter. This study is limited by the fact that it is a hospital-based study. Therefore, it may not reflect what is obtainable in the general population of pregnant women. Another limitation is that there are presently no studies that have assessed portal vein diameter in healthy pregnant women, which we would have used to compare the results from our study.

5. CONCLUSION

This research has established baseline values for normal range of PV diameter in healthy pregnant women in the South-South region of Nigeria, and also revealed significant correlation of PV diameter with age, parity, gestational age and estimated foetal weight. A nomogram showing the value of portal vein diameter at different estimated gestational ages and estimated foetal weights was produced. More researches in pregnant women are therefore recommended.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

The protocol for this study was approved by the Research and Ethics Committee of the Federal Medical Centre Yenagoa, Bayelsa State, Nigeria (FMCY/REC/ECC/2023/685).

ACKNOWLEDGEMENT

The authors appreciate all the patients and staff of the four health institutions for the various roles they played in making this research possible.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Luntsi G, Sani M, Zira JD, Ivor NC, Garba SH. Sonographic assessment of the portal vein diameter in apparently healthy adults in a Northern Nigerian population. *Afr Health Sci.* 2016;16(4):1163-1168. DOI:10.4314/ahs.v16i4.35
2. Anakwue AM, Anakwue R, Agwu KK, Idigo F, Ugwu A, Nwogu U. Sonographic Evaluation of Normal Portal Vein Diameter in Nigerians. *Eur J Sci Res.* 2009;36(1):114-117.
3. Ong JP, Sands M, Younossi ZM. Transjugular intrahepatic portosystemic shunts (TIPS): a decade later. *J Clin Gastroenterol.* 2000;30(1):14-28. DOI:10.1097/00004836-200001000-00005
4. Weinreb J, Kumari S, Phillips G, Pochaczewsky R. Portal vein measurements by real-time sonography. *AJR Am J Roentgenol.* 1982;139(3):497-499. DOI:10.2214/ajr.139.3.497
5. Al-Nakshabandi NA. The role of ultrasonography in portal hypertension. *Saudi J Gastroenterol Off J Saudi Gastroenterol Assoc.* 2006;12(3):111-117. DOI:10.4103/1319-3767.29750
6. Hawaz Y, Admassie D, Kebede T. Ultrasound Assessment of Normal Portal Vein Diameter in Ethiopians Done at Tikur Anbessa Specialized Hospital. *East Cent Afr J Surg.* 2012;17(1):90-93. DOI:10.4314/ecajs.v17i1
7. Uppalapati S, Lokesh S. Correlation of portal vein diameter with the presence of oesophageal varices in chronic liver disease: A prospective study. *Int J Adv Med.* 2018;5(4):859-864. DOI:10.18203/2349-3933.ijam20182529
8. Geleto G, Getnet W, Tewelde T. Mean Normal Portal Vein Diameter Using Sonography among Clients Coming to Radiology Department of Jimma University Hospital, Southwest Ethiopia. *Ethiop J Health Sci.* 2016;26(3):237-242.
9. Pinto-Silva RA, Queiroz LC, Azeredo LM, Silva LC, Lambertucci JR. Ultrasound in

- schistosomiasis mansoni. Mem Inst Oswaldo Cruz. 2010;105(4):479-484. DOI:10.1590/s0074-02762010000400021
10. Mandal L, Mandal SK, Bandyopadhyay D, Datta S. Correlation of portal vein diameter and splenic size with gastro-oesophageal varices in cirrhosis of liver. JIACM. 2011;12(4):266-270.
 11. Usman AU, Ibinaiye P, Ahidjo A, Tahir A, Sa'ad ST, Mustapha Z, et al. Determination of normal portal vein diameter on ultrasound scan among adults in northeastern Nigeria. Arch Int Surg. 2015;5(3):143. DOI:10.4103/2278-9596.167507
 12. Ohmae H, Tanaka M, Hayashi M, Matsuzaki Y, Kurosaki Y, Blas BL, et al. Improvement of ultrasonographic and serologic changes in Schistosoma japonicum-infected patients after treatment with praziquantel. Am J Trop Med Hyg. 1992;46(1):99-104. DOI:10.4269/ajtmh.1992.46.99
 13. Usman A, Tahir A, Sa'ad S, Tahir N, Ibinaiye P, Ahidjo A, et al. Ultrasound Determination of Portal Vein Diameter in Adult Patients with Chronic Liver Disease in North-Eastern Nigeria. Sub-Sahar Afr J Med. 2015;2(2):57-63. DOI:10.4103/2384-5147.157419
 14. Adeyekun AA, Tsebi HB. Grey-scale sonographic evaluation of portal vein diameter in healthy Nigerian adults. J Med Biomed Res. 2014;13(1):17-24. doi:10.4314/jmbr.v13i1
 15. Bolarinwa OA. Sample size estimation for health and social science researchers: The principles and considerations for different study designs. Niger Postgrad Med J. 2020;27(2):67-75. DOI:10.4103/npmj.npmj_19_20
 16. LaMorte WW. Sample Size for One Sample, Continuous Outcome. Boston University School of Public Health. Accessed October 26, 2022. Available: https://sphweb.bumc.bu.edu/otlt/mph-modules/bs/bs704_power/BS704_Power3.html
 17. Tasu JP, Rocher L, PEletier G, Kuoch V, Kulh E, Miquel A, et al. Hepatic venous pressure gradients measured by duplex ultrasound. Clin Radiol. 2002;57(8):746-752. DOI:10.1053/crad.2002.0951
 18. Cosar S, Oktar SO, Cosar B, Yücel C, Ozdemir H. Doppler and gray-scale ultrasound evaluation of morphological and hemodynamic changes in liver vasculature in alcoholic patients. Eur J Radiol. 2005; 54(3):393-399. DOI:10.1016/j.ejrad.2004.07.015
 19. Akanni D, Alassan KS, Kiki M, Djohoun B, Tove KMS de, Sehonou J. Ultrasound Diameter of the Portal Vein to Healthy Adult in Parakou (Benin). Open J Med Imaging. 2021;11(04):145-152. DOI:10.4236/ojmi.2021.114013
 20. Rokni-Yazdi H, Sotoudeh H. Assessment of Normal Doppler Parameters of Portal Vein and Hepatic Artery in 37 Healthy Iranian Volunteers. Iran J Radiol. 2006;3(4):213-216.
 21. Singh S, Pankaj AK, Rani A, Sharma PK, Chauhan P. Gender Linked Metric Analysis of Portal Vein: A Sonographic Appraisal. J Clin Diagn Res JCDR. 2017;11(3):AC13-AC15. DOI:10.7860/JCDR/2017/22851.9461
 22. Moriyasu F, Nishida O, Ban N, Nakamura T, Sakai M, Miyake T, et al. "Congestion index" of the portal vein. AJR Am J Roentgenol. 1986;146(4):35-39. DOI:10.2214/ajr.146.4.735
 23. Gareeballah A, Hassan L, Elzaki M, Ibraheem S, Abelwahab B, Siddig A, et al. Measurement of Normal Portal Vein Diameter in Sudanese Using Ultrasonography. Glob Adv Res J Med Med Sci. 2017;6(12):336-340.
 24. Saha N, Sarkar R, Singh M. Portal vein diameter in a tertiary care centre in North-East India. IOSR J Dent Med Sci. 2016;14(12):110-113. DOI: 10.9790/0853-14121110113
 25. Lal N, Lal V, Majumdar S, Moitra S. Anthropometric correlates of sonographically-determined normal portal vein diameter: results from a study conducted in Rajasthan, India. Int J Anat Res. 2018;6(3.3):5588-5592. DOI:10.16965/ijar.2018.208
 26. Shrout PE, Fleiss JL. Intraclass correlations: Uses in assessing rater reliability. Psychol Bull. 1979;86: 420-428. DOI:10.1037/0033-2909.86.2.420
 27. Figueras F, Fernández S, Hernández-Andrade E, Gratacós E. Umbilical venous blood flow measurement: accuracy and reproducibility. Ultrasound Obstet Gynecol. 2008;32(4):587-591. DOI:10.1002/uog.5306
 28. Costa-Santos C, Bernardes J, Ayres-de-Campos D, Costa A, Costa C. The limits of

agreement and the intraclass correlation coefficient may be inconsistent in the interpretation of agreement. J Clin Epidemiol. 2011;64(3): 264-269.
DOI:10.1016/j.jclinepi.2009.11.010

29. Fernandez S, Figueras F, Gomez O, Martinez JM, Eixarch E, Comas M, et al. Intra- and interobserver reliability of umbilical vein blood flow. Prenat Diagn. 2008;28(11):999-1003.
DOI:10.1002/pd.2092

© 2023 Orijji et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96489>